arXiv:0908.1363v1 [hep-ph] 10 Aug 2009

UNIVERSITY OF CALIFORNIA
SANTA CRUZ

PHENOMENOLOGY OF THE BASIS-INDEPENDENT CP-VIOLATING TWO-HIGGS DOUBLET MODEL

A dissertation submitted in partial satisfaction of the
requirements for the degree of

DOCTOR OF PHILOSOPHY

in

PHYSICS

by

Deva A. O'Neil

June 2009

The Dissertation of Deva A. O'Neil
is approved:

Professor Howard Haber, Chair

Professor Michael Dine

Professor Jason Nielsen

Dean Lisa Sloan
Vice Provost and Dean of Graduate Studies

Copyright © by
Deva A. O'Neil
2009

Table of Contents

List of Figures **v**

List of Tables **vi**

Abstract **viii**

Acknowledgments **ix**

1 Introduction **1**

1.1 Electroweak Symmetry Breaking in the Standard Model 1

1.1.1 The Higgs Mechanism 2

1.1.2 The Scalar Lagrangian 3

1.1.3 Generating Fermion Masses 4

1.1.4 Constraints on the Standard Model Higgs Mass 5

1.1.5 Results of Higgs Searches 8

1.1.6 Future of Higgs Searches 9

1.2 Beyond the Standard Model 9

2 The CP-Violating Two-Higgs Doublet Model **11**

2.1 Motivation 11

2.2 The Basis-Independent Formalism 12

2.3 The Higgs Bases 15

2.4 The Physical Higgs Mass-Eigenstates 18

2.5 Higgs Couplings to Bosons 24

2.6 Higgs Couplings to Fermions 28

3 Special Limits: CP Conservation, The 2HDM with $Z_6 = 0$, and Custodial Symmetry **32**

3.1 The CP-Conserving 2HDM 33

3.1.1 Basis-Independent Analysis of the CP-Conserving Limit 33

3.1.2 The CP-Conserving Limit in the Real Basis 38

3.2 The 2HDM with $Z_6 = 0$ 41

3.2.1 CP Conservation with $Z_6 = 0$ 42

3.2.2 $Z_6 = 0$ with Degenerate Neutral Scalars 45

3.2.3 Special Case: The CP-Conserving Limit when $Z_6 = 0$ and $Z_7 = 0$ 47
3.3 The Custodial Limit of the 2HDM 48
3.3.1 The Basis-Independent Condition for Custodial Symmetry in the Scalar Sector 53
3.3.2 Degeneracy in the Custodial Limit 54
3.3.3 Basis-Dependent Formulations of Custodial Symmetry in the Higgs-Quark Sector 55
3.3.4 The Basis-Independent Custodially-Symmetric Higgs-Quark Lagrangian 57

4 Phenomenology of the 2HDM 58
4.1 The Oblique Parameters S, T and U 58
4.1.1 The Parameter T and the Custodial Limit 60
4.1.2 S, T, and U in the CP-Conserving Limit 61
4.2 Numerical Analysis 62
4.2.1 Splitting Between Neutral and Charged Higgs Masses 66
4.3 Measuring $tan(\beta)$ 66

5 Conclusion 72

A The 2HDM scalar potential in a generic basis 74

B The neutral Higgs boson squared-mass matrix in a generic basis 75

C Explicit formulae for the neutral Higgs masses and mixing angles 79

D The Decoupling Limit of the 2HDM 84
D.1 The Decoupling Condition in the Basis-Independent Formalism 84
D.2 Dimension 6 Operators in the Decoupling Limit 86
D.3 The Lack of CP-Violating Effects in Dimension 6 Operators 89

E Derivation of Basis-Independent Conditions for CP-invariance of Scalar Potential 91
E.1 Calculation of S, T and U 92

F Derivation of Tree-Level Unitarity Limits 101

Bibliography 103

List of Figures

1.1 Bounds on the Standard Model Higgs mass from ref. [77], using $m_t = 173$ GeV and $\alpha_3(M_Z) = .118$. The lowest bound is from vacuum metastability, the middle bound from vacuum stability, and the upper (dotted) bounds are from perturbativity requiring $\lambda(\Lambda) < 3, 6$ (see [70]). 7

4.1 Scatterplots for T as a function of S, with $m_1 = 117 \pm 15$ GeV. The ellipse, representing the 2 σ contour, is adapted from [6]. The second plot shows a close-up of the allowed region. 64

4.2 S and T shown as a function of $m_{H^\pm}$ (solid line), with Y_2 fixed to be m_W^2. The mass scale is in GeV. The unitarity bound (dashed line) applies to Z_3; all other parameters were picked so as to undersaturate unitarity. The 1σ and 2σ contours (shaded regions) are obtained from eq. (4.21). . 65

4.3 S and T shown as a function of the mass splitting between the charged Higgs and the heaviest neutral Higgs. The mass scale is in GeV. The central values (dashed lines) and the 1σ and 2σ contours (shaded regions) are obtained from eq. (4.21). Splitting of more than 100 GeV is disfavored by the constraints from T. 67

List of Tables

2.1 The U(2)-invariant quantities $q_{k\ell}$ are functions of the the neutral Higgs mixing angles θ_{12} and θ_{13}, where $c_{ij} \equiv \cos\theta_{ij}$ and $s_{ij} \equiv \sin\theta_{ij}$. 22

3.1 The U(2)-invariant quantities $q_{k\ell}$ in the CP-conserving limit. Case I: $s_{13} = \mathrm{Im}(Z_6\, e^{-i\theta_{23}}) = \sin\phi = 0$. G^0 and h_3 are CP-odd; h_1 and h_2 are CP-even. 35
3.2 The U(2)-invariant quantities $q_{k\ell}$ in the CP-conserving limit. Case IIa: $s_{12} = \mathrm{Re}(Z_6\, e^{-i\theta_{23}}) = \cos\phi = 0$. G^0 and h_2 are CP-odd; h_1 and h_3 are CP-even. 36
3.3 The U(2)-invariant quantities $q_{k\ell}$ in the CP-conserving limit. Case IIb: $c_{12} = \mathrm{Re}(Z_6\, e^{-i\theta_{23}}) = \cos\phi = 0$. G^0 and h_1 are CP-odd; h_2 and h_3 are CP-even. 36
3.4 Values of $q_{k\ell}$ in the CP-conserving limit with $Z_6 = 0$. Case (i): $s_{13} = c_{12} = \mathrm{Im}(Z_5\, e^{-2i\theta_{23}}) = 0$. The CP quantum numbers are shown for $\mathrm{Im}(Z_7\, e^{-i\theta_{23}}) = \mathrm{Im}(\rho^Q e^{i\bar\theta_{23}}) = 0$ (upper sign), and $\mathrm{Re}(Z_7\, e^{-i\theta_{23}}) = \mathrm{Re}(\rho^Q e^{i\bar\theta_{23}}) = 0$ (lower sign). 44
3.5 Values of $q_{k\ell}$ in the CP-conserving limit with $Z_6 = 0$. Case (ii): $s_{13} = s_{12} = \mathrm{Im}(Z_5\, e^{-2i\theta_{23}}) = 0$. The CP quantum numbers are shown for $\mathrm{Im}(Z_7\, e^{-i\theta_{23}}) = \mathrm{Im}(\rho^Q e^{i\bar\theta_{23}}) = 0$ (upper sign), and $\mathrm{Re}(Z_7\, e^{-i\theta_{23}}) = \mathrm{Re}(\rho^Q e^{i\bar\theta_{23}}) = 0$ (lower sign). 45
3.6 Values of $q_{k\ell}$ in the CP-conserving limit with $Z_6 = 0$. Case (iii): $c_{13} = \mathrm{Im}(Z_5\, e^{2i(\theta_{12}-\theta_{23})})$. The CP quantum numbers are shown for $\mathrm{Im}(Z_7\, e^{-i\theta_{23}}) = \mathrm{Im}(\rho^Q e^{i\bar\theta_{23}}) = 0$ (upper sign), and $\mathrm{Re}(Z_7\, e^{-i\theta_{23}}) = \mathrm{Re}(\rho^Q e^{i\bar\theta_{23}}) = 0$ (lower sign). 45
3.7 Values of $q_{k\ell}$ with $Z_6 = 0$. Case (iv): $s_{13} = \mathrm{Im}(Z_5\, e^{-2i\theta_{23}}) = Z_1 - A^2/v^2 - \mathrm{Re}(Z_5\, e^{-2i\theta_{23}}) = 0$. 46
3.8 Values of $q_{k\ell}$ with $Z_6 = 0$. Case (v): $c_{12} = \mathrm{Im}(Z_5\, e^{-2i\theta_{23}}) = Z_1 - A^2/v^2 = 0$. 46
3.9 Values of $q_{k\ell}$ with $Z_6 = 0$. Case (vi): $s_{12} = \mathrm{Im}(Z_5\, e^{-2i\theta_{23}}) = Z_1 - A^2/v^2 = 0$. 46
3.10 The CP quantum numbers of the neutral scalars with $Z_6 = 0$. Case (iv): $s_{13} = \mathrm{Im}(Z_5\, e^{-2i\theta_{23}}) = Z_1 - A^2/v^2 - \mathrm{Re}(Z_5\, e^{-2i\theta_{23}}) = 0$. 47

3.11 The CP quantum numbers of the neutral scalars with $Z_6 = 0$. Case (v): $c_{12} = \mathrm{Im}(Z_5\, e^{-2i\theta_{23}}) = Z_1 - A^2/v^2 = 0$. 47
3.12 The CP quantum numbers of the neutral scalars with $Z_6 = 0$. Case (vi): $s_{12} = \mathrm{Im}(Z_5\, e^{-2i\theta_{23}}) = Z_1 - A^2/v^2 = 0$. 47

D.1 The U(2)-invariant quantities $q_{k\ell}$ in the exact decoupling limit. 85

E.1 Feynman rules used in the calculation of the oblique parameters. 95
E.2 Diagrams representing the 2HDM contributions to S, part 1. 96
E.3 Diagrams representing the 2HDM contributions to S, part 2. 96
E.4 Diagrams representing the 2HDM contributions to T, part 1. 97
E.5 Diagrams representing the 2HDM contributions to T, part 2. 98
E.6 Diagrams representing the 2HDM contributions to $S + U$. 99
E.7 Standard Model contributions to the oblique parameters. 100

F.1 Calculation of tree-level unitarity limits on the CP-conserving quartic couplings. Combinatoric factors are included to take into account identical particles. 102

Abstract

Phenomenology of the Basis-Independent CP-Violating Two-Higgs Doublet Model

by

Deva A. O'Neil

The Two-Higgs Doublet Model (2HDM) is a model of low-energy particle interactions that is identical to the Standard Model except for the addition of an extra Higgs doublet. This extended Higgs sector would appear in experiments as the presence of multiple Higgs particles, both neutral and charged. The neutral states may either be eigenstates of CP (in the CP-conserving 2HDM), or be mixtures of CP eigenstates (in the CP-violating 2HDM). In order to understand how to measure the couplings of these new particles, this document presents the theory of the CP-violating 2HDM in a basis-independent formalism and explicitly identifies the physical parameters of the model, including a discussion of $\tan\beta$-like parameters. The CP-conserving limit, decoupling limit, and the custodial limit of the model are presented.

In addition, phenomenological constraints from the oblique parameters (S, T, and U) are discussed. A survey of the parameter space of this model shows that the 2HDM is consistent with a large range of possible values for T. Our results also suggest that the 2HDM favors a slightly positive value of S and a value of U within .02 of zero, which is consistent with present data within the statistical error. In a scenario in which the heaviest scalar particle is the charged Higgs boson, we find that the measured value of T puts an upper limit on the mass difference between the charged Higgs boson and the heaviest neutral Higgs boson.

Acknowledgments

The text of this dissertation includes a reprint of the following previously published material:

Howard E. Haber, Deva O'Neil.*Basis-Independent Methods for the Two-Higgs-Doublet Model II: The Significance of* $\tan\beta$*.* Phys.Rev.D74:015018,2006. [hep-ph/0602242] The co-author listed in this publication directed and supervised the research which forms the basis for the dissertation.

This effort could not have been completed without the strong support and guidance from the members of my committee, Howard Haber, Michael Dine, and Jason Nielsen. In particular, I would especially like to thank Dr. Haber for serving as my advisor and for the time and effort he spent guiding this research, and Dr. Dine for his advice and mentoring. In completing this dissertation, I have also benefited from conversations with Dr. Nielsen and Dr. John Mason. I am deeply grateful to have had the privilege of working with these individuals, some of the foremost leaders in this field.

Chapter 1

Introduction

The nature of electroweak symmetry breaking is one of the most important remaining puzzles in particle physics today. The Standard Model of particle physics contains a mechanism for electroweak symmetry breaking (EWSB), but experimental confirmation for it has not yet materialized. Many other models of EWSB have also been proposed. Since the Large Hadron Collider, which will start running in fall 2009, is optimized to make discoveries at the electroweak scale, refining experimental predictions and constraints for these models is a pressing research goal. The purpose of this document is to explore electroweak symmetry breaking through the Two-Higgs Doublet Model (2HDM), an extension of the Standard Model. The 2HDM presents interesting theoretical possibilities, such as CP-violation in the scalar sector of the theory. It also presents challenges, since its phenomena are constrained by electroweak precision data. In this document a basis-independent version of the 2HDM will be presented in both CP-conserving and CP-violating scenarios. In addition to the formalism of the model, identification of the observable parameters of the 2HDM will be emphasized. New insights into custodial symmetry in the context of this model will also be discussed. Finally, the phenomenology of the CP-violating 2HDM will be explored, making use of the "oblique" parameters S, T, and U; and the well-known parameter $\tan\beta$.

1.1 Electroweak Symmetry Breaking in the Standard Model

The Standard Model of particle physics is constructed by applying gauge symmetries to the interactions of fundamental particles. At high energies (above the electroweak scale) the gauge group is SU(3)×SU(2)×U(1). Here we will focus on the electroweak sector of the theory [SU(2)×U(1)], introduced by Glashow, Weinberg, and Salam [48, 49]. A gauge symmetry alone requires a massless vector field corresponding to each generator of the symmetry group. However, spontaneous breaking of the gauge symmetry produces masses for the vector fields. The breaking of SU(2)×U(1) to $\mathrm{U(1)_{EM}}$ leads to one gauge boson remaining massless (which we identify as the photon) and three gauge bosons acquiring mass, which reproduces the pattern observed in nature in the electroweak interactions.

This spontaneous symmetry breaking occurs when a scalar field acquires a non-zero vacuum expectation value (the "Higgs Mechanism"). Once one postulates the existence of such a scalar field (the Higgs field), it may then be employed to give masses to fermions. The details of this mechanism are discussed below.

1.1.1 The Higgs Mechanism

Using the notation of [102], we will represent this hypothetical scalar field as ϕ, and take its U(1) charge to be $+\frac{1}{2}$. It also has an SU(2) spinor structure. Then under the SU(2)×U(1) gauge group, the field transforms as

$$\phi \rightarrow e^{i\alpha^a \tau^a} e^{i\beta/2}\phi, \tag{1.1}$$

where τ^a are the generator matrices for SU(2), ie $\tau^a = \sigma^a/2$. By convention, the vev of this field is taken to have the form

$$<\phi>= \frac{1}{\sqrt{2}}\begin{pmatrix} 0 \\ v \end{pmatrix}. \tag{1.2}$$

The kinetic term of the scalar Lagrangian is written

$$\mathcal{L}_{KE} = |D_\mu \phi|^2, \tag{1.3}$$

where D_μ is the covariant derivative $\delta_\mu - igA^a_\mu \tau^a - i\frac{1}{2}g' B_\mu$. The mass terms for the gauge bosons appear when the vacuum expectation value of eq. (1.3) is taken:

$$\begin{aligned}
\mathcal{L}_{KE} \quad &= \tfrac{1}{2}\begin{pmatrix} 0 & v \end{pmatrix} \left(gA^a_\mu \tau^a + \tfrac{1}{2}g' B_\mu \mathrm{I}\right) \left(gA^{\mu b}\tau^b + \tfrac{1}{2}g' B^\mu \mathrm{I}\right) \begin{pmatrix} 0 \\ v \end{pmatrix} \\
&= \tfrac{1}{2}v^2 \begin{pmatrix} 0 & 1 \end{pmatrix} \left(\tfrac{g}{2}A^a_\mu \sigma^a + \tfrac{g'}{2}B_\mu \mathrm{I}\right) \left(\tfrac{g}{2}A^{\mu b}\sigma^b + \tfrac{g'}{2}B^\mu \mathrm{I}\right) \begin{pmatrix} 0 \\ 1 \end{pmatrix} \\
&= \tfrac{v^2}{8}\left[g^2 (A^1_\mu)^2 + g^2 (A^2_\mu)^2 + \left(g' B_\mu - gA^3_\mu\right)^2\right].
\end{aligned} \tag{1.4}$$

The first two terms in eq. (1.4) gives the mass of the charged field $W^\pm_\mu = \frac{1}{\sqrt{2}}(A^1_\mu \mp iA^2_\mu)$, and the second gives the mass of the neutral field $Z_\mu = \frac{1}{\sqrt{g^2+g'^2}}(gA^3_\mu - g' B_\mu)$. The remaining orthogonal field is $A_\mu = \frac{1}{\sqrt{g^2+g'^2}}(gA^3_\mu + g' B_\mu)$. Thus, one can rewrite eq. (1.4) in terms of observable fields as follows:

$$\mathcal{L}_{KE} = \frac{v^2}{4}g^2 W^+_\mu W^{\mu -} + \frac{v^2}{8}(g^2 + g'^2) Z_\mu Z^\mu. \tag{1.5}$$

Reading off the masses for the gauge bosons yields

$$\begin{aligned}
m_W &= g\frac{v}{2}, \quad m_Z = \sqrt{g^2 + g'^2}\frac{v}{2}, \\
m_A &= 0.
\end{aligned} \tag{1.6}$$

The massless field A_μ we identify as the photon.

1.1.2 The Scalar Lagrangian

In order for the Higgs field to have a non-zero vev, we take its Lagrangian to have the form

$$\mathcal{L} = |D_\mu \phi|^2 + \mu^2 \phi^\dagger \phi - \lambda(\phi^\dagger \phi)^2, \tag{1.7}$$

so that the potential energy has a minimum at

$$v = \sqrt{\mu^2/\lambda}. \tag{1.8}$$

It is conventional to expand ϕ around its vev:

$$\phi = \frac{1}{\sqrt{2}} \begin{pmatrix} \sqrt{2}\, G^+ \\ v + h(x) + iG^0 \end{pmatrix}, \qquad \phi^* = \frac{1}{\sqrt{2}} \begin{pmatrix} \sqrt{2}\, G^- \\ v + h(x) - iG^0 \end{pmatrix}. \tag{1.9}$$

The G^0, $G^\pm$ are Goldstone bosons; The field h(x) is a neutral scalar field of zero vev, whose excitations give rise to a scalar particle, the Higgs boson. We will work in unitary gauge, absorbing G^0 and $G^\pm$ into the gauge potential terms, so that $h(x)$ is a real-valued field. In this gauge,

$$\phi = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ v + h(x) \end{pmatrix}. \tag{1.10}$$

The Higgs boson's potential energy density can be found from the latter terms in eq. (1.7):

$$V = \mu^2 h^2 + \lambda v h^3 + \frac{1}{4}\lambda h^4. \tag{1.11}$$

One can then read off the mass of the Higgs boson, $m_h = \sqrt{2}\mu$. Comparing to eq. (1.8), one obtains

$$m_h = \sqrt{2\lambda}v. \tag{1.12}$$

Thus, without knowing the value of the Higgs self-coupling λ, one cannot predict the value of m_h, even though v is known from measurements of the W mass to be 246 GeV.

Although the Goldstone fields do not explicitly appear in the Lagrangian, and thus do not correspond to observable particles, the degrees of freedom that they represent are still present in the theory after electroweak symmetry breaking. Three of the original four massless vector fields have acquired mass, which adds three degrees of freedom to the gauge boson sector. More precisely, the Goldstone bosons $G^0, G^\pm$ become the longitudinally polarized states of Z^0 and $W^\pm$, respectively, which is reflected in the equivalence theorem [12],[30],[33],[90]: A high energy process ($q \gg m_W$) involving longitudinal gauge bosons has the same amplitude as one in which they are replaced by the corresponding Goldstone bosons [$Z^0 \to G^0$ and $W^\pm \to G^\pm$], up to $\mathcal{O}(m_W/q)$.

The mixing between the A^3_μ and B_μ that produces the physical Z boson and photon fields may be represented in terms of the "weak mixing angle" θ_W, as follows:

$$\begin{pmatrix} Z^0 \\ A \end{pmatrix} = \begin{pmatrix} \cos\theta_W & -\sin\theta_W \\ \sin\theta_W & \cos\theta_W \end{pmatrix} \begin{pmatrix} A^3 \\ B \end{pmatrix}, \tag{1.13}$$

where $\cos\theta_W = \frac{g}{\sqrt{g^2+g'^2}}$ and $\sin\theta_W = \frac{g'}{\sqrt{g^2+g'^2}}$. (Note that there is a simple relation between the masses of the Z and W, $m_W = m_Z \cos\theta_W$.) Then one can rewrite the covariant derivative as

$$D_\mu = \delta_\mu - i\frac{g}{\sqrt{2}}(W_\mu^+ T^+ + W_\mu^- T^-) - i\frac{g}{\cos\theta_W} Z_\mu (T^3 - \sin^2\theta_W Q) - ieA_\mu Q, \tag{1.14}$$

where the generators in this basis are $T^\pm = \frac{1}{2}(\sigma^1 \pm i\sigma^2)$ and $Q = T^3 + Y$, and e is the electric charge, related to g via $g = \frac{e}{\sin\theta_W}$.

We have thus shown how our original gauge symmetry $SU(2)\times U(1)$ with generators T^a and Y has been broken, leaving an unbroken symmetry $U(1)_{EM}$ whose generator is Q and gauge boson is the photon. The other 3 gauge bosons have gained mass through the Higgs mechanism. In the Standard Model, this mechanism also implies the existence of a fundamental massive scalar, the Higgs boson.

1.1.3 Generating Fermion Masses

To explore the effect of electroweak symmetry breaking on the fermion sector, let us first concentrate on leptons. We will later generalize this discussion to quarks. In the Lagrangian, left-handed leptons appear in SU(2) doublets $E_L = \begin{pmatrix} \nu_e \\ e^- \end{pmatrix}_L$. For the first generation,

$$\mathscr{L} = \bar{E}_L(i\not{D})E_L + \bar{e}_R(i\not{D})e_R + \text{quark terms}. \tag{1.15}$$

The right- and left-handed fermion fields are defined the usual way; $\psi_{R,L} \equiv P_{R,L}\psi$, where $P_{R,L} \equiv \frac{1}{2}(1 \pm \gamma_5)$. Explicit lepton mass terms cannot be added to this model, since left-handed fermions appear as SU(2) doublets and the right-handed fermions are singlets. Hence, terms such as $m_e\bar{e}_R e_L + m_e\bar{e}_L e_R$ are forbidden. Thus, to generate mass terms, one requires a scalar SU(2) doublet ϕ that interacts with right- and left-handed leptons, which one can parametrize in terms of a dimensional coupling constant η^E as

$$-\mathscr{L}_Y = \eta^E \bar{E}_L \cdot \phi e_R + h.c. \tag{1.16}$$

After ϕ acquires a vev, a mass term is generated:

$$-\mathscr{L}_Y = \frac{v}{\sqrt{2}}\eta^E \bar{e}_L e_R + h.c., \tag{1.17}$$

from which one can identify $m_e = \frac{1}{\sqrt{2}}\eta^E v$.

An analogous calculation can be made for quarks; in a one-generation model, masses would be generated in the form $m_d = \frac{v}{\sqrt{2}}\eta^D$ and $m_u = \frac{v}{\sqrt{2}}\eta^U$ from Yukawa interactions similar to eq. (1.16). However, when the model is expanded to 3 generations, additional complications arise from the mixing of quark generations. Writing the SU(2) doublets as $Q_L^i = \begin{pmatrix} U^i \\ D^i \end{pmatrix}_L$, one has

$$-\mathscr{L}_{\mathrm{Y}} = \overline{Q_L^0} \cdot \widetilde{\phi}\, \eta^U\, U_R^0 + \overline{Q}_L^0 \cdot \phi\, (\eta^D)^\dagger\, D_R^0 + \text{h.c.}\,, \tag{1.18}$$

where $\widetilde{\phi} \equiv i\sigma_2 \phi^*$, and the flavor indices have been suppressed. Here, Q_L^0, U_R^0, D_R^0 denote the interaction basis quark fields, which are vectors in the quark flavor space. In this basis, the couplings $\eta^{Q,0}$ ($Q = U\,,\ D$) are two non-diagonal 3×3 matrices. To identify the quark mass-eigenstates, one applies unitary transformations of the left- and right-handed U^0 and D^0 fields to diagonalize the matrices η^Q, i.e.:

$$\begin{aligned} P_L U &= V_L^U P_L U^0\,, \qquad & P_R U &= V_R^U P_R U^0\,, \\ P_L D &= V_L^D P_L D^0\,, \qquad & P_R D &= V_R^D P_R D^0\,, \end{aligned} \tag{1.19}$$

and the Cabibbo-Kobayashi-Maskawa (CKM) matrix is defined by $K \equiv V_L^U V_L^{D\,\dagger}$. The Yukawa coupling matrices in this basis are as follows:

$$\eta^U \equiv V_L^U\, \eta^{U,0}\, V_R^{U\,\dagger}\,, \qquad\qquad \eta^D \equiv V_R^D\, \eta^{D,0}\, V_L^{D\,\dagger}\,. \tag{1.20}$$

[Note the different ordering of V_L^Q and V_R^Q in the definitions of η^Q for $Q = U,\ D$.] In terms of the mass-eigenstates, eq. (1.18) becomes

$$-\mathscr{L}_{\rm Y} = \overline{U}_L \cdot \widetilde{\phi}^0\, \eta^U U_R + \overline{D}_L K^\dagger \cdot \phi^-\, \eta^{U\,\dagger} U_R + \overline{U}_L K \cdot \widetilde{\phi}^+\, \eta^{D,\dagger} D_R + \overline{D}_L \cdot \phi^0\, \eta^{D\,\dagger} D_R + \text{h.c.} \tag{1.21}$$

One could write eq. (1.21) more compactly by defining $\mathcal{U} \equiv K^\dagger U$, and $\mathcal{Q}_L = \binom{\mathcal{U}}{D}_L$. Then the Yukawa Lagrangian in the mass-eigenstate basis can be written

$$-\mathscr{L}_{\rm Y} = \bar{\mathcal{Q}}_L \cdot \widetilde{\phi}\, \eta^U \mathcal{U}_R + \bar{\mathcal{Q}}_L \cdot \phi\, \eta^{D\,\dagger} D_R + \text{h.c.} \tag{1.22}$$

One can now obtain the mass matrices M_U and M_D by taking the vev of ϕ in eq. (1.21), which yields

$$M_U = \frac{v}{\sqrt{2}} \eta^U \quad = \quad \text{diag}(m_u\,,\, m_c\,,\, m_t) = V_L^U M_U^0 V_R^{U\,\dagger}\,, \tag{1.23}$$

$$M_D = \frac{v}{\sqrt{2}} \eta^{D\,\dagger} \quad = \quad \text{diag}(m_d\,,\, m_s\,,\, m_b) = V_L^D M_D^0 V_R^{D\,\dagger}\,. \tag{1.24}$$

By generating masses for the fermions and gauge bosons, the Higgs mechanism allows the Standard Model to reproduce the phenomena observed in nature. In order to confirm that this is how electroweak symmetry breaking is implemented in real life, we would have to observe the production of the Higgs boson in particle accelerators. Certain theoretical constraints may be used to indicate the likely range of the Higgs mass, as discussed in the following section.

1.1.4 Constraints on the Standard Model Higgs Mass

1.1.4.1 Bounds on the Higgs Mass from Finiteness and Vacuum Stability

The Standard Model cannot be a valid description of nature at all energy scales. It must be superseded by a theory that incorporates gravitational interactions near the Planck scale (10^{19} GeV). It may be that the scale at which new physics beyond the

Standard Model emerges, Λ, is high (far above the TeV scale), in which case the mass of the SM Higgs must be fairly light, lest the Higgs self-coupling become divergent at a scale below Λ. For $\Lambda \sim M_{Pl}$, the resulting upper bound from the two-loop renormalization group equation (RGE) is $m_h < 180$ GeV [70], in rough agreement with more recent (two-loop) calculations, which have placed the upper bound at 174 GeV [75] and 161.3 ± 20.6 GeV [107]. Since the only case in which λ would remain finite at all energy scales would be in the non-interacting (or "trivial") theory, in which $\lambda = 0$, this is sometimes called a "triviality" argument.

On the other hand, if new physics enters at a low scale (on the order of 1 TeV), the Higgs boson can be heavier. In a pure scalar (ϕ^4) theory, the Higgs mass can be as much as 1 TeV before λ is driven to infinity below the cut-off scale [91]. Cabbibo et al. [24] derive a stringent upper bound on m_h by extending this analysis to include Yukawa interactions with the top quark. For the RGE of λ they exhibit

$$16\pi^2 \frac{d\lambda}{dt} = 12\lambda^2 + 6\lambda y_t^2 - 3y_t^4 + \mathcal{O}(\alpha), \tag{1.25}$$

where $t = \ln(q^2/v^2)$ and y_t is the Yukawa coupling for the quark, $y_t = \sqrt{2}m_t/v$. Requiring that the Higgs coupling $\lambda(q)$ be finite (up to some high energy scale where new physics sets in, such as m_{GUT}), they numerically calculate the maximum value for $\lambda(v)$. (This upper bound for $\lambda(v)$ is dependent on the mass of the top quark, which was not then known.) The coupling $\lambda(v)$ can then be related to the Higgs mass via eq. (1.12). For $m_t = 175\,\mathrm{GeV}$, this corresponded roughly to

$$m_h \lesssim 200\,\mathrm{GeV}. \tag{1.26}$$

This calculation can be repeated for two loops, but theoretical uncertainties are significant [70].

To bound the Higgs boson mass from below, one considers vacuum stability. This condition specifies that $V(v) \leq V(\phi)$ for all $|\phi| < \Lambda$, where $V(v)$ is the value of the scalar potential at the electroweak minimum. For small values of the running coupling $\lambda(v)$, the top quark contribution to the RGE can produce a negative value of $\lambda(q)$, so that the radiatively-corrected effective scalar potential would be unbounded from below [92, 113, 5, 28][1]. This occurs because while $V_{tree} \sim \lambda\phi^4$, for large ϕ, $V_{eff} \sim \lambda(t)\phi^4$, where $t = \ln(\phi^2/M^2)$ and M is the renormalization scale [112]. Thus at some scale t, $\lambda(t)$ becomes negative.

One subtlety that appears in the literature on vacuum stability is that the scalar potential at large $|\phi|$ can have a region deeper than the electroweak vacuum provided that the decay of the "false" electroweak vacuum is suppressed [7, 76, 9, 41, 112, 114]. By requiring that the lifetime of the metastable electroweak vacuum is less than the age of the universe, one derives a lower bound on m_h that is not as strict as the one from stability. The most recent calculation of this metastable region is given in ref. [77] and shown in Fig. 1.1.

---

[1] This argument is controversial. Branchina et al. [22, 21] argue that the region of instability in the scalar potential lies beyond the range of validity for the perturbative RGE. This absence of vacuum instability is confirmed by lattice results [43]. The results of refs. [22] and [21] are disputed by [40].



Figure 1.1: Bounds on the Standard Model Higgs mass from ref. [77], using $m_t = 173$ GeV and $\alpha_3(M_Z) = .118$. The lowest bound is from vacuum metastability, the middle bound from vacuum stability, and the upper (dotted) bounds are from perturbativity requiring $\lambda(\Lambda) < 3, 6$ (see [70]).

Because these RGEs arise from perturbative calculations, one might worry that they would not be valid for large λ. Lattice calculations have been used in order to produce a non-perturbative analysis, which would be valid even for high values of the coupling. The non-perturbative limit on the Higgs mass is found to be $m_h < 9\, m_W \approx 700$ GeV [86]. This result is similar to that derived by Lunscher and Weisz [95], $m_h < 9.6\, m_W$. This consistency suggests that the perturbative calculations may be roughly accurate.

1.1.4.2 Higgs Mass Bounds from Unitarity

In a theory of the electroweak interactions without a Higgs boson, tree-level perturbative calculations of scattering amplitudes (such as $W^+W^- \to W^+W^-$) have terms proportional to $\frac{s}{m_W^2}$, which lead to amplitudes greater than unity at high values of the CM energy ($s \gg m_W^2$). This violation of unitarity is fixed by the presence of the Higgs boson; once diagrams involving the Higgs boson are included, the terms of order $\frac{s}{m_W^2}$ in the tree-level scattering amplitudes cancel, leaving only $\mathcal{O}(1)$ terms. Thus, the Higgs boson has the effect of "unitarizing" gauge boson scattering. In principle, different mechanisms could unitarize $\bar{f}f \to V_L V_L$ and $V_L V_L \to V_L V_L$, respectively; in the Standard Model the Higgs boson unitarizes both [78].

Lee et al. [90][89] derived a critical value of the Higgs mass from tree-level unitarity of processes involving longitudinally polarized gauge bosons, with the condition

$|a_0| \leq 1$, where a_0 is the amplitude of the zeroeth partial wave for $W^+W^- \to W^+W^-$. A more restrictive version of this condition, $|\text{Re } a_0| \leq \frac{1}{2}$ [4, 47, 94], produces a stricter bound of

$$m_h^2 \leq \frac{4\pi\sqrt{2}}{3G_F} \approx (700\,\text{GeV})^2, \tag{1.27}$$

as derived in ref. [97].

This is a tree-level result, which represents the maximum value of m_h for which a perturbative analysis of the scattering amplitude is reliable at all energy scales. If the Higgs mass is above this value, it would indicate that the weak interactions become strongly coupled at high energies, or that additional scalar particles not described by the SM are present to stabilize the amplitudes.

This analysis has been extended to two-loops by Durand et al. [38]. Refinements of their calculation in [111] give results that are very similar to that of eq. (1.27).

Attempts have been made to analyze unitarity using non-perturbative approaches, which would be valid for large λ. In particular, one can analyze SU(N)×U(1) theories in the large N limit [39]. Although the numerical results would not necessarily be valid for $N = 2$, this approach may yield a conceptual understanding of the strong coupling regime. Furthermore, it can be used to validate the perturbative approach in the small λ limit. To next-to-leading order in $1/N$, the non-perturbative analysis of $f\bar{f} \to h \to VV$ and $f\bar{f} \to h \to f\bar{f}$ scattering in the large N limit matches the NNLO perturbative results for $m_h < 800 - 900$ GeV [18]. It is also found that above $m_h = 1$ TeV, the Higgs mass no longer increases as the coupling increases, which violates eq. (1.12). This analysis suggests that if the Standard Model is correct, the Higgs particle will be found below about 1 TeV even if the weak interaction becomes strongly coupled.

1.1.5 Results of Higgs Searches

The experimental lower bound on the Higgs mass is currently determined by data from LEP-II, the electron-positron collider at CERN which reached maximum CM energies of 209 GeV. Higgs searches focused on the "Higgsstrahlung" channel, $e^+e^- \to ZH$. No definite discovery of the Higgs was made by time LEP was decommissioned, which put a limit on the SM Higgs mass of $m_h > 114.4$ GeV [13].

At the time of this writing, Higgs searches are proceeding at the upgraded Tevatron, a $p\bar{p}$ collider running at Fermilab with energy 1.96 TeV. Based on non-observation of the Higgs boson by winter 2009, the combined results from both detectors (CDF and D0) were sufficient to exclude a SM Higgs in the mass range $160\,\text{GeV} < \text{m}_\text{h} < 170\,\text{GeV}$ [115]. The dominant channel for Higgs production at this energy scale is gluon fusion ($gg \to h \to W^+W^-$, with final state $l\nu l\nu$, $l = e^-, \mu^-$), although $Wh \to WWW$ (Higgsstrahlung) is also being searched. The most likely channels to produce a lower energy Higgs ($m_h \leq 140\,\text{GeV}$) at the Tevatron are $Wh \to l\nu b\bar{b}$, $Zh \to llb\bar{b}$, and $Zh \to \bar{\nu}\nu b\bar{b}$. It is expected that by the end of 2010, the Tevatron will either show evidence of Higgs producation (up to 3σ level), or be able to exclude the existence of the Higgs over a large energy range ($145\,\mathrm{GeV} < \mathrm{m_h} < 185\,\mathrm{GeV}$) [109].

1.1.6 Future of Higgs Searches

With an energy of 14 TeV, the Large Hadron Collider at CERN, a proton-proton collider, will be powerful enough to produce hundreds of thousands of Higgs particles (if they are light) or tens of thousands of Higgs particles (if they are heavy) [110]. Most of these will be the result of gluon fusion, $gg \to h$, via a top quark loop [10]. Higgs particles are also likely to be produced through $qq \to qqh$ processes (called weak boson fusion or vector boson fusion). The Higgs would then most likely decay to $\bar{b}b$[2], $\tau\tau$ and/or W^+W^-, depending on its mass [3]. Below $\sim$ 150 GeV, the branching ratio for $h \to \gamma\gamma$ is also high enough to be observable. Despite having a smaller branching ratio than $\tau\tau$ or W^+W^-, this channel is easier to distinguish from background ($q\bar{q}, gg \to \gamma\gamma$) [61].

Should an actual scalar particle be discovered in these searches, the question will be asked, "Is this particle the Standard Model Higgs Boson?" Although observation of the particle will yield measurements of its mass, electric charge, and color charge, to distinguish between different theoretical models, it is important to extract information about gauge and Yukawa couplings. What might be discovered that is not predicted by the Standard Model is the subject of the next section.

1.2 Beyond the Standard Model

Since the Higgs boson has not been discovered in experiments, it is not known whether the Higgs mechanism works as predicted by the Standard Model. It may be, for example, that breaking electroweak symmetry through one (or more) scalar fields is not what happens in nature. However, precision tests of the SM suggest that the GWS description of gauge symmetry in the electroweak sector is the correct one at low energies, requiring some mechanism of electroweak symmetry breaking to generate masses for vector and fermion particles. An example of the experimental evidence for postulating that the particles observed so far are part of a spontaneously broken gauge theory is the universality of coupling constants in gauge boson interactions. In particular, the GWS theory predicts that the coupling constant g is the same for cubic and quartic interactions of charged gauge bosons ($W^\pm$). Description of the neutral gauge boson interactions requires only one additional parameter (g' or $\sin\theta_W$).

Another observable consequence of the GWS gauge symmetry is in the different behavior of left- and right-handed fields. The SU(2) gauge group applies charges $T^3 = \pm\frac{1}{2}$ to left-handed leptons (and their corresponding right-handed antiparticles) and $T^3 = 0$ to right-handed leptons (and left-handed antileptons). These charges appear in the Zl^+l^- couplings, which are proportional to $T^3 - \sin^2\theta_W Q$, and thus can be measured in $Z \rightarrow l^+l^-$ decays. For example, the branching ratio of $Z \rightarrow q\bar{q}$ ($q = u, d, c, s, b$) to $Z \rightarrow l^+l^-$ was calculated from Standard Model fits to be $R_\ell \equiv \Gamma_{had}/\Gamma_{\ell\ell} = 20.744$, which agrees with the value of 20.767 ± 0.025 measured at LEP [117]. This asymmetry between left- and right-handed fields also results in a net polarization of the decay products in $Z \rightarrow f\bar{f}$ (ie, an excess of $f_L\bar{f}_R$ over $f_R\bar{f}_L$). This "polarization asymmetry" is parametrized in

---

[2] Although decay rates to $\bar{b}b$ would be high, this channel is not useful for Higgs discovery due to high background rates.

$$\mathcal{A}_{LR} = \frac{\sigma_L - \sigma_R}{\sigma_L + \sigma_R}, \tag{1.28}$$

where σ_L (σ_R) is the e^+e^- cross-section for Z production from left-handed (right-handed) electrons. The measured value of this asymmetry has been found to be $\mathcal{A}_{LR} = .15138 \pm .00216$ [1], which is within 2 σ of the Standard Model fit $\mathcal{A}_{LR} = .1473 \pm .0011$ [6].

Although these (and other) precision tests of the electroweak sector confirm the GWS model of gauge interactions, and in particular the different SU(2) charge assignments for right- and left-handed fermion fields, the electroweak symmetry is not necessarily broken through a single scalar doublet, as in the SM. In constructing other theories, it is common to build on the Standard Model's particle content, so as to preserve the success of the SM fits to precision electroweak data.[3] One can extend the Standard Model's scalar sector by adding additional singlets, doublets, and/or higher multiplets, for example. Although the exact content of the scalar sector is unknown, the relation $m_w = m_z \cos\theta_W$ must be preserved (up to loop corrections), which arises naturally from postulating an unbroken global SU(2) symmetry. This "custodial" symmetry will be discussed in detail in Chapter 3.

---

[3] There are also "higgless" theories such as models with extra dimensions, which provide a Goldstone boson through mechanisms other than interacting scalar fields.

Chapter 2

The CP-Violating Two-Higgs Doublet Model

2.1 Motivation

The Standard Model by itself is not expected to be a complete description of nature. As discussed in section 1.1.4, the SM is considered to be a low-energy effective theory, which ceases to be valid above some energy scale Λ. Even at low energies, the SM with its single Higgs doublet may not be the correct theory. Extensions of the SM have been proposed as solutions to observations such as dark matter and various theoretical problems (grand unification, the "naturalness" problem, the strong CP problem, and insufficient CP violation to account for the matter/anti-matter imbalance). The most popular of these, supersymmetry, produces both coupling constant unification and possible dark matter candidates. The scalar sector of supersymmetry (in its simplest implementation, the Minimal Supersymmetric Standard Model, or MSSM) has two Higgs doublets. Independently of supersymmetry, the Two-Higgs Doublet Model (2HDM) is an extension of the SM, which is identical to the SM except for the one extra Higgs doublet. The 2HDM may be interesting on its own as a potential theory of nature, since the extended Higgs Sector allows for CP violation beyond what is produced by the SM. It is also useful for gaining insight into the scalar sector of supersymmetry, and other models that contain similar scalar content.

A theory with two Higgs doublets has the potential to produce dangerous Higgs-mediated flavor-changing neutral currents (FCNCs) unless the off-diagonal couplings of the neutral Higgs bosons to quarks are absent (or sufficiently small). It is common to apply a discrete symmetry that restricts the Higgs scalar potential so as to eliminate these off-diagonal couplings [35, 37, 63, 69, 87]. The general 2HDM discussed here will not have any symmetries imposed, and thus the FCNCs will be assumed to be suppressed by fine-tuning or heavy scalar masses. The degree to which the Higgs-fermion couplings are constrained by measurements of flavor-changing rates will be left to future work.

If evidence of multiple Higgs bosons is discovered experimentally, it will be necessary to know how to connect the experimentally observed quantities with the physical parameters of the model. Since one would not know in advance what symmetries are present that constrain the scalar sector, the definition of the physical parameters of the Higgs sector should be defined from the most generic implementation of the 2HDM. In a generic 2HDM, an example of an "unphysical" parameter is the common construction

$$\tan\beta \equiv \frac{\langle\Phi_2^0\rangle}{\langle\Phi_1^0\rangle}\,, \tag{2.1}$$

where Φ_1^0 and Φ_2^0 are the neutral components of the two Higgs doublets [34]. As defined, this parameter is ambiguous in a general 2HDM because it depends on the choice of basis for the Higgs fields. The two identical hypercharge-one fields can be redefined by a global 2×2 unitary transformation. The goal of this work is to construct *physical* parameters, which must be basis-independent. The physical parameter that replaces eq. (2.1) was developed in ref. [34] for the CP-conserving 2HDM. The analog for the CP-violating 2HDM will be discussed in Chapter 4.3.

The goal of this chapter is to derive the scalar, gauge boson and Yukawa couplings of the CP-violating 2HDM. (These were presented for the CP-conserving case in ref. [34].) A Two-Higgs Doublet Model is CP-violating if there is no basis in which the couplings of the scalar Lagrangian are real-valued. Complex-valued Higgs couplings can lead to the mixing of the CP-even and CP-odd eigenstates to produce Higgs fields that have indefinite CP. To develop this model in the basis-independent formalism, I will start by presenting my work with H. Haber in ref. [66]. We begin by reviewing basis-independence in section 2.2. In section 2.3, we introduce the Higgs basis (defined to be a basis in which one of the two neutral scalar fields has zero vacuum expectation value), which possesses some invariant features. We review the construction of the Higgs basis and use the basis-independent formalism to highlight the invariant qualities of this basis choice. Ultimately, we are interested in the Higgs mass-eigenstates. In the most general CP-violating 2HDM, three neutral Higgs states mix to form mass-eigenstates that are not eigenstates of CP. In section 2.4, we demonstrate how to define basis-independent Higgs mixing parameters that are crucial for deriving an invariant form for the Higgs couplings. In section 2.5 and section 2.6 we provide the explicit basis-independent forms for the Higgs couplings to bosons (gauge bosons and Higgs boson self-couplings) and fermions (quarks and leptons), respectively.

2.2 The Basis-Independent Formalism

The fields of the two-Higgs-doublet model (2HDM) consist of two identical complex hypercharge-one, $\mathrm{SU(2)_L}$ doublet scalar fields $\Phi_a(x) \equiv (\Phi_a^+(x)\,,\,\Phi_a^0(x))$, where $a = 1, 2$ labels the two Higgs doublet fields, and will be referred to as the Higgs "flavor" index. The Higgs doublet fields can always be redefined by an arbitrary non-singular complex transformation $\Phi_a \to B_{ab}\Phi_b$, where the matrix B depends on eight real parameters. However, four of these parameters can be used to transform the scalar field kinetic energy terms into canonical form.[1] The most general redefinition of the scalar fields [which leaves invariant the form of the canonical kinetic energy terms $\mathscr{L}_{\rm KE}=(D_\mu\Phi)^\dagger_{\bar a}(D^\mu\Phi)_a$] corresponds to a global U(2) transformation, $\Phi_a\to U_{a\bar b}\Phi_b$ [and $\Phi^\dagger_{\bar a}\to\Phi^\dagger_{\bar b}U^\dagger_{b\bar a}$], where the 2×2 unitary matrix U satisfies $U^\dagger_{b\bar a}U_{a\bar c}=\delta_{b\bar c}$. In our index conventions, replacing an unbarred index with a barred index is equivalent to complex conjugation. We only allow sums over barred–unbarred index pairs, which are performed by employing the U(2)-invariant tensor $\delta_{a\bar b}$. The basis-independent formalism consists of writing all equations involving the Higgs sector fields in a U(2)-covariant form. Basis-independent quantities can then be identified as U(2)-invariant scalars, which are easily identified as products of tensor quantities with all barred–unbarred index pairs summed with no Higgs flavor indices left over.

We begin with the most general 2HDM scalar potential. An explicit form for the scalar potential in a generic basis is given in Appendix A. Following refs. [44] and [34], the scalar potential can be written in U(2)-covariant form:

$$\mathcal{V}=Y_{a\bar b}\Phi^\dagger_{\bar a}\Phi_b+\tfrac{1}{2}Z_{a\bar b c\bar d}(\Phi^\dagger_{\bar a}\Phi_b)(\Phi^\dagger_{\bar c}\Phi_d)\,, \tag{2.2}$$

where the indices a, $\bar b$, c and $\bar d$ are labels with respect to the two-dimensional Higgs flavor space and $Z_{a\bar b c\bar d}=Z_{c\bar d a\bar b}$. The hermiticity of $\mathcal{V}$ yields $Y_{a\bar b}=(Y_{b\bar a})^*$ and $Z_{a\bar b c\bar d}=(Z_{b\bar a d\bar c})^*$. Under a U(2) transformation, the tensors $Y_{a\bar b}$ and $Z_{a\bar b c\bar d}$ transform covariantly: $Y_{a\bar b}\to U_{a\bar c}Y_{c\bar d}U^\dagger_{d\bar b}$ and $Z_{a\bar b c\bar d}\to U_{a\bar e}U^\dagger_{f\bar b}U_{c\bar g}U^\dagger_{h\bar d}Z_{e\bar f g\bar h}$. Thus, the scalar potential $\mathcal{V}$ is a U(2)-scalar. The interpretation of these results is simple. Global U(2)-flavor transformations of the two Higgs doublet fields do not change the functional form of the scalar potential. However, the coefficients of each term of the potential depends on the choice of basis. The transformation of these coefficients under a U(2) basis change are precisely the transformation laws of Y and Z given above.

We shall assume that the vacuum of the theory respects the electromagnetic $\mathrm{U}(1)_{\rm EM}$ gauge symmetry. In this case, the non-zero vacuum expectation values of Φ_a must be aligned. The standard convention is to make a gauge-$\mathrm{SU}(2)_{\rm L}$ transformation (if necessary) such that the lower (or second) component of the doublet fields correspond to electric charge $Q=0$. In this case, the most general $\mathrm{U}(1)_{\rm EM}$-conserving vacuum expectation values are:

$$\langle\Phi_a\rangle=\frac{v}{\sqrt{2}}\left(\begin{array}{c}0\\ \widehat v_a\end{array}\right)\,,\qquad\text{with}\qquad \widehat v_a\equiv e^{i\eta}\left(\begin{array}{c}c_\beta\\ s_\beta\, e^{i\xi}\end{array}\right)\,, \tag{2.3}$$

where $v\equiv 2m_W/g=246$ GeV and $\widehat v_a$ is a vector of unit norm. The overall phase η is arbitrary. By convention, we take $0\leq\beta\leq\pi/2$ and $0\leq\xi<2\pi$. Taking the derivative of eq. (2.2) with respect to Φ_b, and setting $\langle\Phi^0_a\rangle=v_a/\sqrt{2}$, we find the covariant form for the scalar potential minimum conditions:

$$v\,\widehat v^*_{\bar a}\,[Y_{a\bar b}+\tfrac{1}{2}v^2Z_{a\bar b c\bar d}\,\widehat v^*_{\bar c}\,\widehat v_d]=0\,. \tag{2.4}$$

---

[1] That is, starting from $\mathscr{L}_{\rm KE}=a\,(D_\mu\Phi_1)^\dagger(D_\mu\Phi_1)+b\,(D_\mu\Phi_2)^\dagger(D_\mu\Phi_2)+\left[c\,(D_\mu\Phi_1)^\dagger(D_\mu\Phi_2)+\text{h.c.}\right]$, where a and b are real and c is complex, one can always find a (non-unitary) transformation B that removes the four real degrees of freedom corresponding to a, b and c and sets $a=b=1$ and $c=0$. Mathematically, such a transformation is an element of the coset space $\mathrm{GL}(2,\mathbb{C})/\mathrm{U}(2)$.

Before proceeding, let us consider the most general global-U(2) transformation (see p. 5 of ref. [99]):

$$U = e^{i\psi} \begin{pmatrix} e^{i\gamma} \cos\theta & e^{-i\zeta} \sin\theta \\ -e^{i\zeta} \sin\theta & e^{-i\gamma} \cos\theta \end{pmatrix} , \tag{2.5}$$

where $-\pi \leq \theta\,, \psi < \pi$ and $-\pi/2 \leq \zeta\,, \gamma \leq \pi/2$ defines the closed and bounded U(2) parameter space. The matrices U with $\psi = 0$ span an SU(2) matrix subgroup of U(2). The factor of $\{e^{i\psi}\}$ constitutes a U(1) subgroup of U(2). More precisely, $\mathrm{U}(2) \cong \mathrm{SU}(2)\times\mathrm{U}(1)/\mathbb{Z}_2$. In the scalar sector, this U(1) coincides with global hypercharge $\mathrm{U}(1)_{\mathrm{Y}}$. However, the former U(1) is distinguished from hypercharge by the fact that it has no effect on the other fields of the Standard Model.

Because the scalar potential is *invariant* under $\mathrm{U}(1)_{\mathrm{Y}}$ hypercharge transformations,[2] it follows that Y and Z are invariant under U(1)-flavor transformations. Thus, from the standpoint of the Lagrangian, only SU(2)-flavor transformations correspond to a change of basis. Nevertheless, the vacuum expectation value $\widehat{v}$ does change by an overall phase under flavor-U(1) transformations. Thus, it is convenient to expand our definition of the basis to include the phase of $\widehat{v}$. In this convention, all U(2)-flavor transformations correspond to a change of basis. The reason for this choice is that it permits us to expand our potential list of basis-independent quantities to include quantities that depend on $\widehat{v}$. Since $\Phi_a \to U_{a\bar{b}}\Phi_b$ it follows that $\widehat{v}_a \to U_{a\bar{b}}\widehat{v}_b$, and the covariance properties of quantities that depend on $\widehat{v}$ are easily discerned.

The unit vector $\widehat{v}_a$ can also be regarded as an eigenvector of unit norm of the Hermitian matrix $V_{a\bar{b}} \equiv \widehat{v}_a\widehat{v}^*_{\bar{b}}$. The overall phase of $\hat{v}_a$ is not determined in this definition, but as noted above different phase choices are related by U(1)-flavor transformations. Since $V_{a\bar{b}}$ is hermitian, it possesses a second eigenvector of unit norm that is orthogonal to $\widehat{v}_a$. We denote this eigenvector by $\widehat{w}_a$, which satisfies:

$$\widehat{v}^*_{\bar{b}}\widehat{w}_b = 0\,. \tag{2.6}$$

The most general solution to eq. (2.6), up to an overall multiplicative phase factor, is:

$$\widehat{w}_b \equiv \widehat{v}^*_{\bar{a}}\epsilon_{ab} = e^{-i\eta} \begin{pmatrix} -s_\beta\, e^{-i\xi} \\ c_\beta \end{pmatrix} . \tag{2.7}$$

That is, we have chosen a convention in which $\widehat{w}_b \equiv e^{i\chi}\widehat{v}^*_{\bar{a}}\epsilon_{ab}$, where $\chi = 0$. Of course, χ is not fixed by eq. (2.6); the existence of this phase choice is reflected in the non-uniqueness of the Higgs basis, as discussed in section 2.3.

The inverse relation to eq. (2.7) is easily obtained: $\widehat{v}^*_{\bar{a}} = \epsilon_{\bar{a}\bar{b}}\,\widehat{w}_b$. Above, we have introduced two Levi-Civita tensors with $\epsilon_{12} = -\epsilon_{21} = 1$ and $\epsilon_{11} = \epsilon_{22} = 0$. However, ϵ_{ab} and $\epsilon_{\bar{a}\bar{b}}$ are not proper tensors with respect to the full flavor-U(2) group (although these are invariant SU(2)-tensors). Consequently, $\widehat{w}_a$ does not transform covariantly with respect to the full flavor-U(2) group. If we write $U = e^{i\psi}\widehat{U}$, with $\det \widehat{U} = 1$ (and $\det U = e^{2i\psi}$), it is simple to check that under a U(2) transformation

---

[2] The $\mathrm{SU}(2)_{\mathrm{L}}\times\mathrm{U}(1)_{\mathrm{Y}}$ gauge transformations act on the fields of the Standard Model, but do *not* transform the coefficients of the terms appearing in the Lagrangian.

$$\widehat{v}_a \to U_{a\bar{b}}\widehat{v}_b \qquad \text{implies that} \qquad \widehat{w}_a \to (\det\, U)^{-1}\, U_{a\bar{b}}\, \widehat{w}_b\,. \tag{2.8}$$

Henceforth, we shall define a pseudotensor[3] as a tensor that transform covariantly with respect to the flavor-SU(2) subgroup but whose transformation law with respect to the full flavor-U(2) group is only covariant modulo an overall non-trivial phase equal to some integer power of $\det U$. Thus, $\widehat{w}_a$ is a pseudovector. However, we can use $\widehat{w}_a$ to construct proper tensors. For example, the Hermitian matrix $W_{a\bar{b}} \equiv \widehat{w}_a \widehat{w}^*_{\bar{b}} = \delta_{a\bar{b}} - V_{a\bar{b}}$ is a proper second-ranked tensor.

Likewise, a pseudoscalar (henceforth referred to as a pseudo-invariant) is defined as a quantity that transforms under U(2) by multiplication by some integer power of $\det U$. We reiterate that pseudo-invariants *cannot* be physical observables as the latter must be true U(2)-invariants.

2.3 The Higgs Bases

Once the scalar potential minimum is determined, which defines $\widehat{v}_a$, one class of basis choices is uniquely selected. Suppose we begin in a generic Φ_1–Φ_2 basis. We define new Higgs doublet fields:

$$H_1 = (H_1^+\,,\, H_1^0) \equiv \widehat{v}^*_{\bar{a}}\Phi_a\,, \qquad\qquad H_2 = (H_2^+\,,\, H_2^0) \equiv \widehat{w}^*_{\bar{a}}\Phi_a = \epsilon_{\bar{b}\bar{a}}\widehat{v}_b\Phi_a\,. \tag{2.9}$$

The transformation between the generic basis and the Higgs basis, $H_a = \widehat{U}_{a\bar{b}}\Phi_b$, is given by the following flavor-SU(2) matrix:

$$\widehat{U} = \begin{pmatrix} \widehat{v}^*_1 & \widehat{v}^*_2 \\ \widehat{w}^*_1 & \widehat{w}^*_2 \end{pmatrix} = \begin{pmatrix} \widehat{v}^*_1 & \widehat{v}^*_2 \\ -\widehat{v}_2 & \widehat{v}_1 \end{pmatrix}\,. \tag{2.10}$$

This defines a particular Higgs basis.

Inverting eq. (2.9) yields:

$$\Phi_a = H_1\widehat{v}_a + H_2\widehat{w}_a = H_1\widehat{v}_a + H_2\widehat{v}^*_{\bar{b}}\epsilon_{ba}\,. \tag{2.11}$$

The definitions of H_1 and H_2 imply that

$$\langle H_1^0\rangle = \frac{v}{\sqrt{2}}\,, \qquad\qquad \langle H_2^0\rangle = 0\,, \tag{2.12}$$

where we have used eq. (2.6) and the fact that $\widehat{v}^*_{\bar{a}}\,\widehat{v}_a = 1$.

The Higgs basis is not unique. Suppose one begins in a generic Φ'_1–Φ'_2 basis, where $\Phi'_a = V_{a\bar{b}}\Phi_b$ and $\det V \equiv e^{i\chi} \neq 1$. If we now define:

$$H'_1 \equiv \widehat{v}^*_{\bar{a}}\Phi'_a\,, \qquad\qquad H'_2 \equiv \widehat{w}^*_{\bar{a}}\Phi'_a\,, \tag{2.13}$$

---

[3] In tensor calculus, analogous quantities are usually referred to as tensor densities or relative tensors [116].

then

$$H_1' = H_1\,, \qquad\qquad H_2' = (\det V)H_2 = e^{i\chi}H_2\,. \tag{2.14}$$

That is, H_1 is an invariant field, whereas H_2 is pseudo-invariant with respect to arbitrary U(2) transformations. In particular, the unitary matrix

$$U_D \equiv \begin{pmatrix} 1 & 0 \\ 0 & e^{i\chi} \end{pmatrix} \tag{2.15}$$

transforms from the unprimed Higgs basis to the primed Higgs basis. The phase angle χ parameterizes the class of Higgs bases. From the definition of H_2 given in eq. (2.9), this phase freedom can be attributed to the choice of an overall phase in the definition of $\widehat{w}$ as discussed in section 2.2. This phase freedom will be reflected by the appearance of pseudo-invariants in the study of the Higgs basis. However, pseudo-invariants are useful in that they can be combined to create true invariants, which are candidates for observable quantities.

It is now a simple matter to insert eq. (2.11) into eq. (2.2) to obtain:

$$\begin{aligned}\mathcal{V} &= Y_1 H_1^\dagger H_1 + Y_2 H_2^\dagger H_2 + [Y_3 H_1^\dagger H_2 + \text{h.c.}] \\ &\quad + \tfrac{1}{2}Z_1(H_1^\dagger H_1)^2 + \tfrac{1}{2}Z_2(H_2^\dagger H_2)^2 + Z_3(H_1^\dagger H_1)(H_2^\dagger H_2) + Z_4(H_1^\dagger H_2)(H_2^\dagger H_1) \\ &\quad + \left\{\tfrac{1}{2}Z_5(H_1^\dagger H_2)^2 + \big[Z_6(H_1^\dagger H_1) + Z_7(H_2^\dagger H_2)\big]H_1^\dagger H_2 + \text{h.c.}\right\}\,,\end{aligned} \tag{2.16}$$

where Y_1, Y_2 and $Z_{1,2,3,4}$ are U(2)-invariant quantities and Y_3 and $Z_{5,6,7}$ are pseudo-invariants. The explicit forms for the Higgs basis coefficients have been given in ref. [34]. The invariant coefficients are conveniently expressed in terms of the second-ranked tensors $V_{a\bar{b}}$ and $W_{a\bar{b}}$ introduced in section 2.2:

$$\begin{aligned} Y_1 &\equiv \operatorname{Tr}(YV)\,, & Y_2 &\equiv \operatorname{Tr}(YW)\,, \\ Z_1 &\equiv Z_{a\bar{b}c\bar{d}}\,V_{b\bar{a}}V_{d\bar{c}}\,, & Z_2 &\equiv Z_{a\bar{b}c\bar{d}}\,W_{b\bar{a}}W_{d\bar{c}}\,, \\ Z_3 &\equiv Z_{a\bar{b}c\bar{d}}\,V_{b\bar{a}}W_{d\bar{c}}\,, & Z_4 &\equiv Z_{a\bar{b}c\bar{d}}\,V_{b\bar{c}}W_{d\bar{a}}\,, \end{aligned} \tag{2.17}$$

whereas the pseudo-invariant coefficients are given by:

$$\begin{aligned} Y_3 &\equiv Y_{a\bar{b}}\,\widehat{v}_{\bar{a}}^*\,\widehat{w}_b\,, & Z_5 &\equiv Z_{a\bar{b}c\bar{d}}\,\widehat{v}_{\bar{a}}^*\,\widehat{w}_b\,\widehat{v}_{\bar{c}}^*\,\widehat{w}_d\,, \\ Z_6 &\equiv Z_{a\bar{b}c\bar{d}}\,\widehat{v}_{\bar{a}}^*\,\widehat{v}_b\,\widehat{v}_{\bar{c}}^*\,\widehat{w}_d\,, & Z_7 &\equiv Z_{a\bar{b}c\bar{d}}\,\widehat{v}_{\bar{a}}^*\,\widehat{w}_b\,\widehat{w}_{\bar{c}}^*\,\widehat{w}_d\,. \end{aligned} \tag{2.18}$$

The invariant coefficients are manifestly real, whereas the pseudo-invariant coefficients are potentially complex.

Using eq. (2.8), it follows that under a flavor-U(2) transformation specified by the matrix U, the pseudo-invariants transform as:

$$[Y_3, Z_6, Z_7] \to (\det U)^{-1}[Y_3, Z_6, Z_7] \qquad \text{and} \qquad Z_5 \to (\det U)^{-2}Z_5\,. \tag{2.19}$$

One can also deduce eq. (2.19) from eq. (2.16) by noting that $\mathcal{V}$ and H_1 are invariant whereas H_2 is pseudo-invariant field that is transforms as:

$$H_2 \to (\det U) H_2\,. \tag{2.20}$$

In the class of Higgs bases defined by eq. (2.14), $\widehat{v} = (1,0)$ and $\widehat{w} = (0,1)$, which are independent of the angle χ that distinguishes among different Higgs bases. That is, under the phase transformation specified by eq. (2.15), both $\widehat{v}$ and $\widehat{w}$ are unchanged. Inserting these values of $\widehat{v}$ and $\widehat{w}$ into eqs. (2.17) and (2.18) yields the coefficients of the Higgs basis scalar potential. For example, the coefficient of $H_1^\dagger H_2$ is given by $Y_{12} = Y_3$ in the unprimed Higgs basis and $Y'_{12} = Y'_3$ in the primed Higgs basis. Using eq. (2.19), it follows that $Y'_{12} = Y_{12} e^{-i\chi}$, which is consistent with the matrix transformation law $Y' = U_D Y U_D^\dagger$.

From the four complex pseudo-invariant coefficients, one can form four independent real invariants $|Y_3|$, $|Z_{5,6,7}|$ and three invariant relative phases $\arg(Y_3^2 Z_5^*)$, $\arg(Y_3 Z_6^*)$ and $\arg(Y_3 Z_7^*)$. Including the six invariants of eq. (2.17), we have therefore identified thirteen independent invariant real degrees of freedom prior to imposing the scalar potential minimum conditions. Eq. (2.4) then imposes three additional conditions on the set of thirteen invariants[4]

$$Y_1 = -\tfrac{1}{2} Z_1 v^2\,, \qquad\qquad Y_3 = -\tfrac{1}{2} Z_6 v^2\,. \tag{2.21}$$

This leaves eleven independent real degrees of freedom (one of which is the vacuum expectation value $v = 246$ GeV) that specify the 2HDM parameter space.

The doublet of scalar fields in the Higgs basis can be parameterized as follows:

$$H_1 = \begin{pmatrix} G^+ \\ \frac{1}{\sqrt{2}}\left(v + \varphi_1^0 + iG^0\right) \end{pmatrix}, \qquad H_2 = \begin{pmatrix} H^+ \\ \frac{1}{\sqrt{2}}\left(\varphi_2^0 + ia^0\right) \end{pmatrix}, \tag{2.22}$$

and the corresponding hermitian conjugated fields are likewise defined. We identify $G^\pm$ as a charged Goldstone boson pair and G^0 as the CP-odd neutral Goldstone boson.[5] In particular, the identification of $G^0 = \sqrt{2}\,\mathrm{Im}\, H_1^0$ follows from the fact that we have defined the Higgs basis [see eqs. (2.9) and (2.12)] such that $\langle H_1^0\rangle$ is real and non-negative. Of the remaining fields, φ_1^0 is a CP-even neutral scalar field, φ_2^0 and a^0 are states of indefinite CP quantum numbers,[6] and $H^\pm$ is the physical charged Higgs boson pair. If the Higgs sector is CP-violating, then φ_1^0, φ_2^0, and a^0 all mix to produce three physical neutral Higgs mass-eigenstates of indefinite CP quantum numbers.

---

[4] The second condition of eq. (2.21) is a complex equation that can be rewritten in terms of invariants: $|Y_3| = \frac{1}{2}|Z_6| v^2$ and $Y_3 Z_6^* = -\frac{1}{2}|Z_6|^2 v^2$.

[5] The definite CP property of the neutral Goldstone boson persists even if the Higgs Lagrangian is CP-violating (either explicitly or spontaneously), as shown in Chapter 3.

[6] The CP-properties of the neutral scalar fields (in the Higgs basis) can be determined by studying the pattern of gauge boson/scalar boson couplings and the scalar self-couplings in the interaction Lagrangian (see section 2.5). If the scalar potential is CP-conserving, then two orthogonal linear combinations of φ_2^0 and a^0 can be found that are eigenstates of CP. By an appropriate rephasing of H_2 (which corresponds to some particular choice among the possible Higgs bases) such that all the coefficients of the scalar potential in the Higgs basis are real, one can then identify φ_2^0 as a CP-even scalar field and a^0 as a CP-odd scalar field. See Chapter 3 for further details.

2.4 The Physical Higgs Mass-Eigenstates

To determine the Higgs mass-eigenstates, one must examine the terms of the scalar potential that are quadratic in the scalar fields (after minimizing the scalar potential and defining shifted scalar fields with zero vacuum expectation values). This procedure is carried out in Appendix B starting from a generic basis. However, there is an advantage in performing the computation in the Higgs basis since the corresponding scalar potential coefficients are invariant or pseudo-invariant quantities [eqs. (2.16)–(2.18)]. This will allow us to identify U(2)-invariants in the Higgs mass diagonalization procedure.

Thus, we proceed by inserting eq. (2.11) into eq. (2.2) and examining the terms linear and quadratic in the scalar fields. The requirement that the coefficient of the linear term vanishes corresponds to the scalar potential minimum conditions [eq. (2.21)]. These conditions are then used in the evaluation of the coefficients of the terms quadratic in the fields. One can easily check that no quadratic terms involving the Goldstone boson fields survive (as expected, since the Goldstone bosons are massless). This confirms our identification of the Goldstone fields in eq. (2.22). The charged Higgs boson mass is also easily determined:

$$m_{H^\pm}^2 = Y_2 + \tfrac{1}{2} Z_3 v^2 \,. \tag{2.23}$$

The three remaining neutral fields mix, and the resulting neutral Higgs squared-mass matrix in the φ_1^0–φ_2^0–a^0 basis is:

$$\mathcal{M} = v^2 \begin{pmatrix} Z_1 & \operatorname{Re}(Z_6) & -\operatorname{Im}(Z_6) \\ \operatorname{Re}(Z_6) & \tfrac{1}{2}\left[Z_3+Z_4+\operatorname{Re}(Z_5)\right] + Y_2/v^2 & -\tfrac{1}{2}\operatorname{Im}(Z_5) \\ -\operatorname{Im}(Z_6) & -\tfrac{1}{2}\operatorname{Im}(Z_5) & \tfrac{1}{2}\left[Z_3+Z_4-\operatorname{Re}(Z_5)\right] + Y_2/v^2 \end{pmatrix} . \tag{2.24}$$

Note that $\mathcal{M}$ depends implicitly on the choice of Higgs basis [eq. (2.14)] via the χ-dependence of the pseudo-invariants Z_5 and Z_6. Moreover, the real and imaginary parts of these pseudo-invariants mix if χ is changed. Thus, $\mathcal{M}$ does not possess simple transformation properties under arbitrary flavor-U(2) transformations. Nevertheless, we demonstrate below that the eigenvalues and normalized eigenvectors are U(2)-invariant. First, we compute the characteristic equation:

$$\det(\mathcal{M} - xI) = -x^3 + \operatorname{Tr}(\mathcal{M})\, x^2 - \tfrac{1}{2}\left[(\operatorname{Tr}\mathcal{M})^2 - \operatorname{Tr}(\mathcal{M}^2)\right] x + \det(\mathcal{M})\,, \tag{2.25}$$

where I is the 3×3 identity matrix. [The coefficient of x in eq. (2.25) is particular to 3×3 matrices (see Fact 4.9.3 of ref. [17]).] Explicitly,

$$\begin{aligned} \operatorname{Tr}(\mathcal{M}) &= 2Y_2 + (Z_1+Z_3+Z_4)v^2\,, \\ \operatorname{Tr}(\mathcal{M}^2) &= Z_1^2 v^4 + \tfrac{1}{2}v^4\left[(Z_3+Z_4)^2 + |Z_5|^2 + 4|Z_6|^2\right] + 2Y_2[Y_2 + (Z_3+Z_4)v^2]\,, \\ \det(\mathcal{M}) &= \tfrac{1}{4}\left\{Z_1 v^6[(Z_3+Z_4)^2 - |Z_5|^2] - 2v^4[2Y_2 + (Z_3+Z_4)v^2]|Z_6|^2 \right. \\ &\qquad \left. + 4Y_2 Z_1 v^2[Y_2 + (Z_3+Z_4)v^2] + 2v^6 \operatorname{Re}(Z_5^* Z_6^2)\right\} . \end{aligned} \tag{2.26}$$

Clearly, all the coefficients of the characteristic polynomial are U(2)-invariant. Since the roots of this polynomial are the squared-masses of the physical Higgs bosons, it follows that the physical Higgs masses are basis-independent as required. Since $\mathcal{M}$ is a real symmetric matrix, the eigenvalues of $\mathcal{M}$ are real. However, if any of these eigenvalues are negative, then the extremal solution of eq. (2.4) with $v \neq 0$ is *not* a minimum of the scalar potential. The requirements that $m^2_{H^\pm} > 0$ [eq. (2.23)] and the positivity of the squared-mass eigenvalues of $\mathcal{M}$ provide basis-independent conditions for the desired spontaneous symmetry breaking pattern specified by eq. (2.3).

The real symmetric squared-mass matrix $\mathcal{M}$ can be diagonalized by an orthogonal transformation

$$R\mathcal{M}R^T = \mathcal{M}_D \equiv \text{diag}\,(m_1^2\,,\, m_2^2\,,\, m_3^2)\,, \tag{2.27}$$

where $RR^T = I$ and the m_k^2 are the eigenvalues of $\mathcal{M}$ [*i.e.*, the roots of eq. (2.25)]. A convenient form for R is:

$$\begin{aligned} R = R_{12}R_{13}R_{23} &= \begin{pmatrix} c_{12} & -s_{12} & 0 \\ s_{12} & c_{12} & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} c_{13} & 0 & -s_{13} \\ 0 & 1 & 0 \\ s_{13} & 0 & c_{13} \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & c_{23} & -s_{23} \\ 0 & s_{23} & c_{23} \end{pmatrix} \\ &= \begin{pmatrix} c_{13}c_{12} & -c_{23}s_{12} - c_{12}s_{13}s_{23} & -c_{12}c_{23}s_{13} + s_{12}s_{23} \\ c_{13}s_{12} & c_{12}c_{23} - s_{12}s_{13}s_{23} & -c_{23}s_{12}s_{13} - c_{12}s_{23} \\ s_{13} & c_{13}s_{23} & c_{13}c_{23} \end{pmatrix}, \end{aligned} \tag{2.28}$$

where $c_{ij} \equiv \cos\theta_{ij}$ and $s_{ij} \equiv \sin\theta_{ij}$. Note that $\det R = 1$, although we could have chosen an orthogonal matrix with determinant equal to -1 by choosing $-R$ in place of R. In addition, if we take the range of the angles to be $-\pi \leq \theta_{12},\ \theta_{23} < \pi$ and $|\theta_{13}| \leq \pi/2$, then we cover the complete parameter space of SO(3) matrices (see p. 11 of ref. [99]). That is, we work in a convention where $c_{13} \geq 0$. However, this parameter space includes points that simply correspond to the redefinition of two of the Higgs mass-eigenstate fields by their negatives. Thus, we may reduce the parameter space further and define all Higgs mixing angles modulo π. We shall verify this assertion at the end of this section.

The neutral Higgs mass-eigenstates are denoted by h_1, h_2 and h_3:

$$\begin{pmatrix} h_1 \\ h_2 \\ h_3 \end{pmatrix} = R \begin{pmatrix} \varphi_1^0 \\ \varphi_2^0 \\ a^0 \end{pmatrix}. \tag{2.29}$$

It is often convenient to choose a convention for the mass ordering of the h_k such that $m_1 \leq m_2 \leq m_3$.

Since the mass-eigenstates h_k do not depend on the initial basis choice, they must be U(2)-invariant fields. In order to present a formal proof of this assertion, we need to determine the transformation properties of the elements of R under an arbitrary U(2) transformation. In principle, these can be determined from eq. (2.27), using the fact that the m_k^2 are invariant quantities. However, the form of $\mathcal{M}$ is not especially convenient for this purpose as noted below eq. (2.24). This can be ameliorated by introducing the unitary matrix:

$$W = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1/\sqrt{2} & 1/\sqrt{2} \\ 0 & -i/\sqrt{2} & i/\sqrt{2} \end{pmatrix}\,, \tag{2.30}$$

and rewriting eq. (2.27) as

$$(RW)(W^\dagger \mathcal{M} W)(RW)^\dagger = \mathcal{M}_D = \text{diag}\,(m_1^2\,,\, m_2^2\,,\, m_3^2)\,. \tag{2.31}$$

A straightforward calculation yields:

$$W^\dagger \mathcal{M} W \;=\; v^2 \begin{pmatrix} Z_1 & \frac{1}{\sqrt{2}} Z_6 & \frac{1}{\sqrt{2}} Z_6^* \\ \frac{1}{\sqrt{2}} Z_6^* & \frac{1}{2}(Z_3+Z_4)+Y_2/v^2 & \frac{1}{2} Z_5^* \\ \frac{1}{\sqrt{2}} Z_6 & \frac{1}{2} Z_5 & \frac{1}{2}(Z_3+Z_4)+Y_2/v^2 \end{pmatrix}, \tag{2.32}$$

$$RW \;=\; \begin{pmatrix} q_{11} & \frac{1}{\sqrt{2}} q_{12}^*\, e^{i\theta_{23}} & \frac{1}{\sqrt{2}} q_{12}\, e^{-i\theta_{23}} \\ q_{21} & \frac{1}{\sqrt{2}} q_{22}^*\, e^{i\theta_{23}} & \frac{1}{\sqrt{2}} q_{22}\, e^{-i\theta_{23}} \\ q_{31} & \frac{1}{\sqrt{2}} q_{32}^*\, e^{i\theta_{23}} & \frac{1}{\sqrt{2}} q_{32}\, e^{-i\theta_{23}} \end{pmatrix}, \tag{2.33}$$

where

$$\begin{aligned} q_{11} &= c_{13}c_{12}\,, & q_{21} &= c_{13}s_{12}\,, & q_{31} &= s_{13}\,, \\ q_{12} &= -s_{12} - ic_{12}s_{13}\,, & q_{22} &= c_{12} - is_{12}s_{13}\,, & q_{32} &= ic_{13}\,. \end{aligned} \tag{2.34}$$

The matrix RW defined in eq. (2.33) is unitary and satisfies $\det RW = i$. Evaluating this determinant yields:

$$\tfrac{1}{2} \sum_{j,k,\ell=1}^{3} \epsilon_{jk\ell}\, q_{j1} \text{Im}(q_{k2}^* q_{\ell 2}) = 1\,, \tag{2.35}$$

while unitarity implies:

$$\text{Re}\,(q_{k1}q_{\ell 1}^* + q_{k2}q_{\ell 2}^*) = \delta_{k\ell}\,, \tag{2.36}$$

$$\sum_{k=1}^{3} |q_{k1}|^2 = \tfrac{1}{2}\sum_{k=1}^{3} |q_{k2}|^2 = 1\,, \qquad \sum_{k=1}^{3} q_{k2}^2 = \sum_{k=1}^{3} q_{k1}q_{k2} = 0\,. \tag{2.37}$$

These results can be used to prove the identity [106]:

$$q_{j1} = \tfrac{1}{2} \sum_{k,\ell=1}^{3} \epsilon_{jk\ell}\, \text{Im}(q_{k2}^* q_{\ell 2})\,. \tag{2.38}$$

Since the matrix elements of $W^\dagger \mathcal{M} W$ only involve invariants and pseudo-invariants, we may use eq. (2.31) to determine the flavor-U(2) transformation properties of $q_{k\ell}$ and $e^{i\theta_{23}}$. The resulting transformation laws are:

$$q_{k\ell} \to q_{k\ell}\,, \qquad \text{and} \qquad e^{i\theta_{23}} \to (\det U)^{-1} e^{i\theta_{23}}\,, \tag{2.39}$$

under a U(2) transformation U. That is, the $q_{k\ell}$ are invariants, or equivalently θ_{12} and θ_{13} (modulo π) are U(2)-invariant angles, whereas $e^{i\theta_{23}}$ is a pseudo-invariant. Eq. (2.39) is critical for the rest of the paper. Finally, to show that the Higgs mass-eigenstates are invariant fields, we rewrite eq. (2.29) as

$$\begin{pmatrix} h_1 \\ h_2 \\ h_3 \end{pmatrix} = RW \begin{pmatrix} \sqrt{2}\,\mathrm{Re}H_1^0 - v \\ H_2^0 \\ H_2^{0\,\dagger} \end{pmatrix}\,. \tag{2.40}$$

Since the $q_{k\ell}$, H_1 and the product $e^{i\theta_{23}} H_2$ are U(2)-invariant quantities, it follows that the h_k are invariant fields.

The transformation laws given in eqs. (2.19) and (2.39) imply that the quantities $Z_5\, e^{-2i\theta_{23}}$, $Z_6\, e^{-i\theta_{23}}$ and $Z_7\, e^{-i\theta_{23}}$ are U(2)-invariant. These combinations will appear in the physical Higgs boson self-couplings of section 2.5 and in the expressions for the invariant mixing angles given in Appendix C. With this in mind, it is useful to rewrite the neutral Higgs mass diagonalization equation [eq. (2.27)] as follows. With $R \equiv R_{12} R_{13} R_{23}$ given by eq. (2.28),

$$\widetilde{\mathcal{M}} \equiv R_{23} \mathcal{M} R_{23}^T = v^2 \begin{pmatrix} Z_1 & \mathrm{Re}(Z_6\, e^{-i\theta_{23}}) & -\mathrm{Im}(Z_6\, e^{-i\theta_{23}}) \\ \mathrm{Re}(Z_6 e^{-i\theta_{23}}) & \mathrm{Re}(Z_5\, e^{-2i\theta_{23}}) + A^2/v^2 & -\tfrac{1}{2}\mathrm{Im}(Z_5\, e^{-2i\theta_{23}}) \\ -\mathrm{Im}(Z_6\, e^{-i\theta_{23}}) & -\tfrac{1}{2}\mathrm{Im}(Z_5\, e^{-2i\theta_{23}}) & A^2/v^2 \end{pmatrix}. \tag{2.41}$$

where A^2 is defined by

$$A^2 \equiv Y_2 + \tfrac{1}{2}[Z_3 + Z_4 - \mathrm{Re}(Z_5 e^{-2i\theta_{23}})]v^2\,. \tag{2.42}$$

The diagonal neutral Higgs squared-mass matrix is then given by the following:

$$\widetilde{R}\,\widetilde{\mathcal{M}}\,\widetilde{R}^T = \mathcal{M}_D = \mathrm{diag}(m_1^2\,,\, m_2^2\,,\, m_3^2)\,, \tag{2.43}$$

where the diagonalizing matrix $\widetilde{R} \equiv R_{12} R_{13}$ depends only on θ_{12} and θ_{13}:

$$\widetilde{R} = \begin{pmatrix} c_{12}c_{13} & -s_{12} & -c_{12}s_{13} \\ c_{13}s_{12} & c_{12} & -s_{12}s_{13} \\ s_{13} & 0 & c_{13} \end{pmatrix}\,. \tag{2.44}$$

Eqs. (2.41)–(2.44) provide a manifestly U(2)-invariant squared-mass matrix diagonalization, since the elements of $\widetilde{R}$ and $\widetilde{\mathcal{M}}$ are invariant quantities.

Eq. (2.40) can be conviently written as

$$h_k = \frac{1}{\sqrt{2}} \left[\overline{H}_1^{0\,\dagger} q_{k1} + H_2^{0\,\dagger} q_{k2} e^{-i\theta_{23}} + \overline{H}_1^0 q_{k1}^* + H_2^0 q_{k2}^* e^{i\theta_{23}} \right]\,, \tag{2.45}$$

where $\overline{H}_1^0 \equiv H_1^0 - v\widehat{v}_a/\sqrt{2}$. The $q_{k\ell}$, defined for $k = 1, 2, 3$ and $\ell = 1, 2$ by eq. (2.34), are displayed in Table 2.1. To account for the Goldstone boson ($k = 4$) we have also introduced: $q_{41} = i$ and $q_{42} = 0$ Note that the $q_{k\ell}$ are invariant, and $\overline{H}_1^0$ and $H_2^0 e^{i\theta_{23}}$ are invariant fields.

In this section, all computations were carried out by first transforming to the Higgs basis. The advantage of this procedure is that one can readily identify the relevant invariant and pseudo-invariant quantities involved in the determination of the Higgs mass-eigenstates. We may now combine eqs. (2.9) and (2.45) to obtain explicit expressions for the Higgs mass-eigenstate fields h_k in terms of the scalar fields in the generic basis Φ_a. Since these expressions do not depend on the Higgs basis, one could have obtained the results for the Higgs mass-eigenstates directly without reference to Higgs basis quantities. In Appendix B, we present a derivation starting from the generic basis, which produces the following expressions for the Higgs mass-eigenstates (and the Goldstone boson) in terms of the generic basis fields:

$$h_k = \frac{1}{\sqrt{2}} \left[\overline{\Phi}_{\bar{a}}^{0\,\dagger} (q_{k1} \widehat{v}_a + q_{k2} \widehat{w}_a e^{-i\theta_{23}}) + (q_{k1}^* \widehat{v}_{\bar{a}}^* + q_{k2}^* \widehat{w}_{\bar{a}}^* e^{i\theta_{23}}) \overline{\Phi}_a^0 \right] , \tag{2.46}$$

for $k = 1, \ldots, 4$, where $h_4 = G^0$. The shifted neutral fields are defined by $\overline{\Phi}_a^0 \equiv \Phi_a^0 - v\widehat{v}_a/\sqrt{2}$.

Table 2.1: The U(2)-invariant quantities $q_{k\ell}$ are functions of the the neutral Higgs mixing angles θ_{12} and θ_{13}, where $c_{ij} \equiv \cos\theta_{ij}$ and $s_{ij} \equiv \sin\theta_{ij}$.

k	q_{k1}	q_{k2}
1	$c_{12}c_{13}$	$-s_{12} - ic_{12}s_{13}$
2	$s_{12}c_{13}$	$c_{12} - is_{12}s_{13}$
3	s_{13}	ic_{13}
4	i	0

Since the $q_{k\ell}$ are U(2)-invariant and $\widehat{w}_a e^{-i\theta_{23}}$ is a *proper* vector under U(2) transformations, it follows that eq. (2.46) provides a U(2)-invariant expression for the Higgs mass-eigenstates. It is now a simple matter to invert eq. (2.46) to obtain

$$\Phi_a = \begin{pmatrix} G^+ \widehat{v}_a + H^+ \widehat{w}_a \\ \\ \dfrac{v}{\sqrt{2}} \widehat{v}_a + \dfrac{1}{\sqrt{2}} \displaystyle\sum_{k=1}^{4} \left(q_{k1} \widehat{v}_a + q_{k2} e^{-i\theta_{23}} \widehat{w}_a \right) h_k \end{pmatrix} , \tag{2.47}$$

where $h_4 \equiv G^0$. The form of the charged upper component of Φ_a is a consequence of eq. (2.11). The U(2)-covariant expression for Φ_a in terms of the Higgs mass-eigenstate scalar fields given by eq. (2.47) is one of the central results of this paper. In sections 2.5 and 2.6, we shall employ this result for Φ_a in the computation of the Higgs couplings of the 2HDM.

Finally, we return to the question of the domains of the angles θ_{ij}. We assume that $Z_6 \equiv |Z_6| e^{i\theta_6} \neq 0$ (the special case of $Z_6 = 0$ is treated at the end of Appendix C).

Since $e^{-i\theta_{23}}$ is a pseudo-invariant, we prefer to deal with the invariant angle ϕ:

$$\phi \equiv \theta_6 - \theta_{23}\,, \qquad \text{where} \qquad \theta_6 \equiv \arg Z_6\,. \tag{2.48}$$

As shown in Appendix C, the invariant angles θ_{12}, θ_{13} and ϕ are determined modulo π in terms of invariant combinations of the scalar potential parameters. This domain is smaller than the one defined by $-\pi \leq \theta_{12},\ \theta_{23} < \pi$ and $|\theta_{13}| \leq \pi/2$, which covers the parameter space of SO(3) matrices. Since the U(2)-invariant mass-eigenstate fields h_k are real, one can always choose to redefine any one of the h_k by its negative. Redefining two of the three Higgs fields h_1, h_2 and h_3 by their negatives[7] is equivalent to multiplying two of the rows of R by -1. In particular,

$$\theta_{12} \to \theta_{12} \pm \pi \quad \Longrightarrow \quad h_1 \to -h_1 \text{ and } h_2 \to -h_2\,, \tag{2.49}$$

$$\phi \to \phi \pm \pi\,, \quad \theta_{13} \to -\theta_{13}\,, \quad \theta_{12} \to \pm\pi - \theta_{12} \quad \Longrightarrow \quad h_1 \to -h_1 \text{ and } h_3 \to -h_3\,, \tag{2.50}$$

$$\theta_{13} \to \theta_{13} \pm \pi\,, \quad \theta_{12} \to -\theta_{12} \quad \Longrightarrow \quad h_1 \to -h_1 \text{ and } h_3 \to -h_3\,, \tag{2.51}$$

$$\phi \to \phi \pm \pi\,, \quad \theta_{13} \to -\theta_{13}\,, \quad \theta_{12} \to -\theta_{12} \quad \Longrightarrow \quad h_2 \to -h_2 \text{ and } h_3 \to -h_3\,, \tag{2.52}$$

$$\theta_{13} \to \theta_{13} \pm \pi\,, \quad \theta_{12} \to \pm\pi - \theta_{12} \quad \Longrightarrow \quad h_2 \to -h_2 \text{ and } h_3 \to -h_3\,. \tag{2.53}$$

This means that if we adopt a convention in which c_{12}, c_{13} and $\sin\phi$ are non-negative, with the angles defined modulo π, then the sign of the Higgs mass-eigenstate fields will be fixed.

Given a choice of the overall sign conventions of the neutral Higgs fields, the number of solutions for the invariant angles θ_{12}, θ_{13} and ϕ modulo π are in one-to-one correspondence with the possible mass orderings of the m_k (except at certain singular points of the parameter space[8]). For example, note that

$$\theta_{12} \to \theta_{12} \pm \pi/2 \Longrightarrow h_1 \to \mp h_2 \text{ and } h_2 \to \pm h_1\,. \tag{2.54}$$

That is, two solutions for θ_{12} exist modulo π. If $m_1 < m_2$, then eq. (C.25) implies that the solutions for θ_{12} and ϕ are correlated such that $s_{12}\cos\phi \geq 0$, and (for fixed ϕ) only one θ_{12} solution modulo π survives. The corresponding effects on the invariant angles that result from swapping other pairs of neutral Higgs fields are highly non-linear and cannot be simply exhibited in closed form. Nevertheless, we can use the results of Appendix C to conclude that for $m_{1,2} < m_3$ (in a convention where $\sin\phi \geq 0$), eq. (C.23) yields $s_{13} \leq 0$, and for $m_1 < m_2 < m_3$, eq. (C.21) implies that $\sin 2\theta_{56}\cos\phi \geq 0$, where $\theta_{56} \equiv -\frac{1}{2}\arg(Z_5^* Z_6^2)$.

The sign of the neutral Goldstone field is conventional, but is not affected by the choice of Higgs mixing angles. Finally, we note that the charged fields $G^\pm$ and

---

[7] In order to have an odd number of Higgs mass-eigenstates redefined by their negatives, one would have to employ an orthogonal Higgs mixing matrix with $\det\ R = -1$.

[8] At singular points of the parameter space corresponding to two (or three) mass-degenerate neutral Higgs bosons, some (or all) of the invariant Higgs mixing angles are indeterminate. An indeterminate invariant angle also arises in the case of $Z_6 = 0$ and $c_{13} = 0$ as explained at the end of Appendix C.

$H^\pm$ are complex. Eq. (2.47) implies that $G^\pm$ is an invariant field and $H^\pm$ is a pseudo-invariant field that transforms as:

$$H^\pm \to (\det U)^{\pm 1}\, H^\pm \tag{2.55}$$

with respect to U(2) transformations. That is, once the Higgs Lagrangian is written in terms the Higgs mass-eigenstates and the Goldstone bosons, one is still free to rephase the charged fields. By convention, we shall fix this phase according to eq. (2.47).

2.5 Higgs Couplings to Bosons

We begin by computing the Higgs self-couplings in terms of U(2)-invariant quantities. First, we use eq. (2.47) to obtain:

$$\begin{aligned}
\Phi^\dagger_{\bar a}\Phi_b \;=\;& \tfrac{1}{2}v^2 V_{b\bar a} + v h_k\left[V_{b\bar a}\,\mathrm{Re}\; q_{k1} + \tfrac{1}{2}\left(\widehat{v}_b \widehat{w}^*_{\bar a} q^*_{k2} e^{i\theta_{23}} + \widehat{v}^*_{\bar a}\widehat{w}_b q_{k2} e^{-i\theta_{23}}\right)\right] \\
&+\tfrac{1}{2}h_j h_k\left[V_{b\bar a}\mathrm{Re}(q^*_{j1}q_{k1}) + W_{b\bar a}\mathrm{Re}(q^*_{j2}q_{k2})\right. \\
&\quad\left. +\widehat{v}_b \widehat{w}^*_{\bar a} q^*_{j_2} q_{k1} e^{i\theta_{23}} + \widehat{v}^*_{\bar a}\widehat{w}_b q^*_{j1} q_{k2} e^{-i\theta_{23}}\right] \\
&+G^+G^- V_{b\bar a} + H^+H^- W_{b\bar a} + G^-H^+\widehat{v}^*_{\bar a}\widehat{w}_b + G^+H^-\widehat{w}^*_{\bar a}\widehat{v}_b\,,
\end{aligned} \tag{2.56}$$

where repeated indices are summed over and $j,k = 1,\ldots,4$. We then insert eq. (2.56) into eq. (2.2), and expand out the resulting expression. We shall write:

$$\mathcal{V} = \mathcal{V}_0 + \mathcal{V}_2 + \mathcal{V}_3 + \mathcal{V}_4\,, \tag{2.57}$$

where the subscript indicates the overall degree of the fields that appears in the polynomial expression. $\mathcal{V}_0$ is a constant of no significance and $\mathcal{V}_1 = 0$ by the scalar potential minimum condition. $\mathcal{V}_2$ is obtained in Appendix C. In this section, we focus on the cubic Higgs self-couplings that reside in $\mathcal{V}_3$ and the quartic Higgs self-couplings that reside in $\mathcal{V}_4$.

Using eqs. (2.17) and (2.18), one can express $\mathcal{V}_3$ and $\mathcal{V}_4$ in terms of the invariants (Y_1, Y_2 and $Z_{1,2,3,4}$) and pseudo-invariants (Y_3, $Z_{5,6,7}$). In the resulting expressions, we have eliminated Y_1 and Y_3 by the scalar potential minimum conditions [eq. (2.21)]. The cubic Higgs couplings are governed by the following terms of the scalar potential:

$$\begin{aligned}
\mathcal{V}_3 \;=\;& \tfrac{1}{2}v\, h_j h_k h_\ell\Big[q_{j1}q^*_{k1}\mathrm{Re}(q_{\ell1})Z_1 + q_{j2}q^*_{k2}\,\mathrm{Re}(q_{\ell1})(Z_3+Z_4) + \mathrm{Re}(q^*_{j1}q_{k2}q_{\ell2}Z_5\, e^{-2i\theta_{23}}) \\
&\qquad +\mathrm{Re}\left([2q_{j1}+q^*_{j1}]q^*_{k1}q_{\ell2}Z_6\, e^{-i\theta_{23}}\right) + \mathrm{Re}(q^*_{j2}q_{k2}q_{\ell2}Z_7\, e^{-i\theta_{23}})\Big] \\
&+v\, h_k G^+G^-\Big[\mathrm{Re}(q_{k1})Z_1 + \mathrm{Re}(q_{k2}\, e^{-i\theta_{23}}Z_6)\Big] \\
&+v\, h_k H^+H^-\Big[\mathrm{Re}(q_{k1})Z_3 + \mathrm{Re}(q_{k2}\, e^{-i\theta_{23}}Z_7)\Big] \\
&+\tfrac{1}{2}v\, h_k\Big\{G^-H^+\, e^{i\theta_{23}}\left[q^*_{k2}Z_4 + q_{k2}\, e^{-2i\theta_{23}}Z_5 + 2\mathrm{Re}(q_{k1})Z_6\, e^{-i\theta_{23}}\right] + \text{h.c.}\Big\}\,,
\end{aligned} \tag{2.58}$$

where there is an implicit sum over the repeated indices[9] j, k, $\ell = 1,2,3,4$. Since the neutral Goldstone boson field is denoted by $h_4 \equiv G^0$, we can extract the cubic couplings of G^0 by using $q_{41} = i$ and $q_{42} = 0$. The only cubic Higgs–G^0 couplings that survive are:

$$\begin{aligned}
\mathcal{V}_{3G} \;&=\; \tfrac{1}{2}v\sum_{k=1}^{3}\sum_{\ell=1}^{3} G^0 h_k h_\ell\Big[\mathrm{Im}(q_{k2}q_{\ell 2}Z_5\,e^{-2i\theta_{23}}) + 2q_{k1}\,\mathrm{Im}\left(q_{\ell 2}Z_6\,e^{-i\theta_{23}}\right)\Big] \\
&\quad +\tfrac{1}{2}v\sum_{\ell=1}^{3} G^0G^0 h_\ell\Big[q_{\ell 1}Z_1 + \mathrm{Re}(q_{\ell 2}Z_6 e^{-i\theta_{23}})\Big]\,, \qquad (2.59)
\end{aligned}$$

where we have used the fact that q_{j1} is real for $j = 1,2,3$.

At the end of the section 2.4, we noted that H^+ is a pseudo-invariant field. However $e^{i\theta_{23}}H^+$ is a U(2)-invariant field [see eqs. (2.39) and (2.55)], and it is precisely this combination that shows up in eq. (2.58). Moreover, as shown in section 2.4, the $q_{k\ell}$ and the quantities $Z_5\,e^{-2i\theta_{23}}$, $Z_6\,e^{-i\theta_{23}}$ and $Z_7\,e^{-i\theta_{23}}$ are also invariant with respect to flavor-U(2) transformations. Thus, we conclude that eq. (2.58) is U(2)-invariant as required.

The quartic Higgs couplings are governed by the following terms of the scalar potential:

$$\begin{aligned}
\mathcal{V}_4 = \tfrac{1}{8}h_j h_k h_l h_m\Big[&q_{j1}q_{k1}q^*_{\ell 1}q^*_{m1}Z_1 + q_{j2}q_{k2}q^*_{\ell 2}q^*_{m2}Z_2 \\
&+2q_{j1}q^*_{k1}q_{\ell 2}q^*_{m2}(Z_3+Z_4) + 2\mathrm{Re}(q^*_{j1}q^*_{k1}q_{\ell 2}q_{m2}Z_5\,e^{-2i\theta_{23}}) \\
&+4\mathrm{Re}(q_{j1}q^*_{k1}q^*_{\ell 1}q_{m2}Z_6\,e^{-i\theta_{23}}) + 4\mathrm{Re}(q^*_{j1}q_{k2}q_{\ell 2}q^*_{m2}Z_7\,e^{-i\theta_{23}})\Big] \\
&+\tfrac{1}{2}h_j h_k G^+G^-\Big[q_{j1}q^*_{k1}Z_1 + q_{j2}q^*_{k2}Z_3 + 2\mathrm{Re}(q_{j1}q_{k2}Z_6\,e^{-i\theta_{23}})\Big] \\
&+\tfrac{1}{2}h_j h_k H^+H^-\Big[q_{j2}q^*_{k2}Z_2 + q_{j1}q^*_{k1}Z_3 + 2\mathrm{Re}(q_{j1}q_{k2}Z_7\,e^{-i\theta_{23}})\Big] \\
&+\tfrac{1}{2}h_j h_k\Big\{G^-H^+\,e^{i\theta_{23}}\Big[q_{j1}q^*_{k2}Z_4 + q^*_{j1}q_{k2}Z_5\,e^{-2i\theta_{23}} + q_{j1}q^*_{k1}Z_6\,e^{-i\theta_{23}} \\
&\qquad +q_{j2}q^*_{k2}Z_7\,e^{-i\theta_{23}}\Big] + \mathrm{h.c.}\Big\} + \tfrac{1}{2}Z_1G^+G^-G^+G^- + \tfrac{1}{2}Z_2H^+H^-H^+H^- \\
&+(Z_3+Z_4)G^+G^-H^+H^- + \tfrac{1}{2}Z_5H^+H^+G^-G^- + \tfrac{1}{2}Z^*_5H^-H^-G^+G^+ \\
&+G^+G^-(Z_6H^+G^- + Z^*_6H^-G^+) + H^+H^-(Z_7H^+G^- + Z^*_7H^-G^+)\,, \qquad (2.60)
\end{aligned}$$

---

[9] Note that the sum over repeated indices can be rewritten by appropriately symmetrizing the relevant coefficients. For example, $\sum_{jk\ell} g_{jk\ell}\,h_j h_k h_\ell = \sum_{j\leq k\leq \ell} h_j h_k h_\ell\,[g_{jk\ell} + \mathrm{perm}]$, where "perm" is an instruction to add additional terms (as needed) such that the indices j, k and ℓ appear in all possible *distinct* permutations.

where there is an implicit sum over the repeated indices j, k, ℓ, $m = 1, 2, 3, 4$. One can check the U(2)-invariance of $\mathcal{V}_4$ by noting that $Z_5 H^+ H^+$, $Z_6 H^+$ and $Z_7 H^+$ are U(2)-invariant combinations.[10] It is again straightforward to isolate the quartic couplings of the neutral Goldstone boson ($h_4 \equiv G^0$):

$$\begin{aligned}
\mathcal{V}_{4G} = {} & \tfrac{1}{8} q_{11}^4 Z_1 G^0 G^0 G^0 G^0 + \tfrac{1}{2} \mathrm{Im}(q_{m2} Z_6 \, e^{-i\theta_{23}}) \, G^0 G^0 G^0 h_m \\
& + \tfrac{1}{4} G^0 G^0 h_\ell h_m \bigg[q_{\ell 1} q_{m1} Z_1 + q_{\ell 2} q_{m2}^* (Z_3 + Z_4) - \mathrm{Re}(q_{\ell 2} q_{m2} Z_5 e^{-2i\theta_{23}}) \\
& \quad + 2 q_{\ell 1} \mathrm{Re}(q_{m2} Z_6 e^{-i\theta_{23}}) \bigg] + \tfrac{1}{2} G^0 h_k h_\ell h_m \bigg[q_{k1} \mathrm{Re}(q_{\ell 2} q_{m2} Z_5 e^{-2i\theta_{23}}) \\
& \quad + q_{k1} q_{\ell 1} \mathrm{Re}(q_{m2} Z_6 e^{-i\theta_{23}}) + \mathrm{Re}(q_{k2} q_{\ell 2} q_{m2}^* Z_7 e^{-i\theta_{23}}) \bigg] \\
& - \mathrm{Im}(q_{m2} Z_6 \, e^{-i\theta_{23}}) \, G^+ G^- G^0 h_m - \mathrm{Im}(q_{m2} Z_7 \, e^{-i\theta_{23}}) \, H^+ H^- G^0 h_m \\
& + \tfrac{1}{2} i \, G^0 h_m \bigg\{ G^- H^+ e^{i\theta_{23}} \left[q_{m2}^* Z_4 - q_{m2} Z_5 e^{-2i\theta_{23}} \right] + \text{h.c.} \bigg\} \\
& + \tfrac{1}{2} Z_1 G^0 G^0 G^+ G^- + \tfrac{1}{2} Z_3 G^0 G^0 H^+ H^- \\
& + \tfrac{1}{2} Z_6 G^0 G^0 G^- H^+ + \tfrac{1}{2} Z_6^* G^0 G^0 G^+ H^- ,
\end{aligned} \tag{2.61}$$

where the repeated indices k, ℓ, $m = 1, 2, 3$ are summed over.

The Feynman rules are obtained by multiplying the relevant terms of the scalar potential by $-iS$, where the symmetry factor $S = \prod_i n_i!$ for an the interaction term that possesses n_i identical particles of type i. Explicit forms for the $q_{k\ell}$ in terms of the invariant mixing angles θ_{12} and θ_{13} are displayed in Table 2.1. For example, the Feynman rule for the cubic self-coupling of the lightest neutral Higgs boson is given by $ig(h_1 h_1 h_1)$ where

$$\begin{aligned}
g(h_1 h_1 h_1) = {} & -3v \bigg[Z_1 c_{12}^3 c_{13}^3 + (Z_3 + Z_4) c_{12} c_{13} |s_{123}|^2 + c_{12} c_{13} \, \mathrm{Re}(s_{123}^2 Z_5 \, e^{-2i\theta_{23}}) \\
& - 3 c_{12}^2 c_{13}^2 \, \mathrm{Re}(s_{123} Z_6 \, e^{-i\theta_{23}}) - |s_{123}|^2 \, \mathrm{Re}(s_{123} Z_7 \, e^{-i\theta_{23}}) \bigg] ,
\end{aligned} \tag{2.62}$$

where $s_{123} \equiv s_{12} + i c_{12} s_{13}$. Similarly, the Feynman rule for the quartic self-coupling of the lightest neutral Higgs boson is given by $ig(h_1 h_1 h_1 h_1)$ where

$$\begin{aligned}
g(h_1 h_1 h_1 h_1) = {} & -3 \bigg[Z_1 c_{12}^4 c_{13}^4 + Z_2 |s_{123}|^4 + 2(Z_3 + Z_4) c_{12}^2 c_{13}^2 |s_{123}|^2 \\
& + 2 c_{12}^2 c_{13}^2 \, \mathrm{Re}(s_{123}^2 Z_5 \, e^{-2i\theta_{23}}) - 4 c_{12}^3 c_{13}^3 \, \mathrm{Re}(s_{123} Z_6 \, e^{-i\theta_{23}}) \\
& - 4 c_{12} c_{13} |s_{123}|^2 \, \mathrm{Re}(s_{123} Z_7 \, e^{-i\theta_{23}}) \bigg] .
\end{aligned} \tag{2.63}$$

---

[10] It is instructive to write, *e.g.*, $Z_6 H^+ = (Z_6 \, e^{-i\theta_{23}})(H^+ \, e^{i\theta_{23}})$, *etc.* to exhibit the well-known U(2)-invariant combinations.

We turn next to the coupling of the Higgs bosons to the gauge bosons. These arise from the Higgs boson kinetic energy terms when the partial derivatives are replaced by the gauge covariant derivatives: $\mathscr{L}_{\rm KE} = D^\mu \Phi^\dagger_{\bar a} D_\mu \Phi_a$. In the $\mathrm{SU(2)_L}\times\mathrm{U(1)}$ electroweak gauge theory,

$$D_\mu \Phi_a = \begin{pmatrix} \partial_\mu \Phi_a^+ + \left[\dfrac{ig}{c_W}\left(\tfrac{1}{2}-s_W^2\right)Z_\mu + ieA_\mu\right]\Phi_a^+ + \dfrac{ig}{\sqrt{2}}W_\mu^+\Phi_a^0 \\ \partial_\mu \Phi_a^0 - \dfrac{ig}{2c_W}Z_\mu\Phi_a^0 + \dfrac{ig}{\sqrt{2}}W_\mu^-\Phi_a^+ \end{pmatrix}, \tag{2.64}$$

where $s_W \equiv \sin\theta_W$ and $c_W \equiv \cos\theta_W$. Inserting eq. (2.64) into $\mathscr{L}_{\rm KE}$ yields the Higgs boson–gauge boson interactions in the generic basis. Finally, we use eq. (2.47) to obtain the interaction Lagrangian of the gauge bosons with the physical Higgs boson mass-eigenstates. The resulting interaction terms are:

$$\begin{aligned}
\mathscr{L}_{VVH} \;=\;& \left(gm_W W_\mu^+ W^{\mu\,-} + \frac{g}{2c_W} m_Z Z_\mu Z^\mu\right)\mathrm{Re}(q_{k1})h_k \\
&+ em_W A^\mu (W_\mu^+ G^- + W_\mu^- G^+) - g m_Z s_W^2 Z^\mu (W_\mu^+ G^- + W_\mu^- G^+)\,,
\end{aligned} \tag{2.65}$$

$$\begin{aligned}
\mathscr{L}_{VVHH} \;=\;& \left[\tfrac{1}{4}g^2 W_\mu^+ W^{\mu\,-} + \frac{g^2}{8c_W^2} Z_\mu Z^\mu\right]\mathrm{Re}(q_{j1}^* q_{k1} + q_{j2}^* q_{k2})\, h_j h_k \\
&+ \Bigg[\tfrac{1}{2}g^2 W_\mu^+ W^{\mu\,-} + e^2 A_\mu A^\mu + \frac{g^2}{c_W^2}\left(\tfrac{1}{2}-s_W^2\right)^2 Z_\mu Z^\mu \\
&\qquad + \frac{2ge}{c_W}\left(\tfrac{1}{2}-s_W^2\right) A_\mu Z^\mu\Bigg](G^+G^- + H^+H^-) \\
&+ \left\{\left(\tfrac{1}{2}egA^\mu W_\mu^+ - \frac{g^2 s_W^2}{2c_W} Z^\mu W_\mu^+\right)(q_{k1}G^- + q_{k2}\, e^{-i\theta_{23}} H^-)h_k + \text{h.c.}\right\},
\end{aligned} \tag{2.66}$$

and

$$\begin{aligned}
\mathscr{L}_{VHH} =\;& \frac{g}{4c_W}\,\mathrm{Im}(q_{j1}q_{k1}^* + q_{j2}q_{k2}^*) Z^\mu h_j \overleftrightarrow{\partial}_\mu\, h_k \\
&- \tfrac{1}{2}g\left\{iW_\mu^+\left[q_{k1}G^- \overleftrightarrow{\partial}^\mu\, h_k + q_{k2}e^{-i\theta_{23}} H^- \overleftrightarrow{\partial}^\mu\, h_k\right] + \text{h.c.}\right\} \\
&+ \left[ieA^\mu + \frac{ig}{c_W}\left(\tfrac{1}{2}-s_W^2\right)Z^\mu\right](G^+ \overleftrightarrow{\partial}_\mu\, G^- + H^+ \overleftrightarrow{\partial}_\mu\, H^-)\,,
\end{aligned} \tag{2.67}$$

where the repeated indices $j,k = 1,\ldots,4$ are summed over. The neutral Goldstone boson interaction terms can be ascertained by taking $h_4 \equiv G^0$:

$$\begin{aligned}\mathscr{L}_{VG} \;=\;& \left[\tfrac{1}{4}g^2 W^+_\mu W^{\mu -} + \frac{g^2}{8c^2_W} Z_\mu Z^\mu\right] G^0 G^0 \\ &+\left\{\tfrac{1}{2}iegA^\mu W^+_\mu G^- G^0 - \frac{ig^2 s^2_W}{2c_W} Z^\mu W^+_\mu G^- G^0 + \text{h.c.}\right\} \\ &+\frac{g}{2c_W}\text{Re}(q_{k1}) Z^\mu G^0 \overleftrightarrow{\partial}_\mu\, h_k + \tfrac{1}{2}g\left(W^+_\mu G^- \overleftrightarrow{\partial}^\mu\, G^0 + W^-_\mu G^+ \overleftrightarrow{\partial}^\mu\, G^0\right) . \end{aligned} \tag{2.68}$$

Once again, we can verify by inspection that the Higgs boson–vector boson interactions are U(2)-invariant. Moreover, one can derive numerous relations among these couplings using the properties of the $q_{k\ell}$. In particular, eqs. (2.36)–(2.38) imply the following relations among the Higgs boson–vector boson couplings [56, 55, 106]:

$$g(ZZh_j) = m_Z \sum_{k,\ell=1}^{3} \epsilon_{jk\ell}\, g(Zh_k h_\ell)\,, \qquad (j = 1, 2, 3)\,, \tag{2.69}$$

$$\sum_{k=1}^{3} [g(VVh_k)]^2 = \frac{g^2 m_V^4}{m_W^2}\,, \qquad V = W^\pm \text{ or } Z\,, \tag{2.70}$$

$$\sum_{1\leq j<k\leq 3} [g(Zh_j h_k)]^2 = \frac{g^2}{4c_W^2}\,, \tag{2.71}$$

$$g(ZZh_j)g(ZZh_k) + 4m_Z^2 \sum_{\ell=1}^{3} g(Zh_j h_\ell) g(Zh_k h_\ell) = \frac{g^2 m_Z^2}{c_W^2}\,\delta_{jk}\,, \tag{2.72}$$

where the Feynman rules for the VVh_k and Zh_jh_k vertices are given by $ig^{\mu\nu}\, g(VVh_k)$ and $(p_k - p_j)^\mu\, g(Zh_jh_k)$, respectively, and the four-momenta p_j, p_k of the neutral Higgs bosons h_j, h_k point into the vertex.[11] Note that eq. (2.72) holds for $j, k = 1, 2, 3, 4$.

2.6 Higgs Couplings to Fermions

The most general Yukawa couplings of Higgs bosons to fermions yield neutral Higgs-mediated flavor-changing neutral currents at tree-level [50, 45, 100]. Typically, these couplings are in conflict with the experimental bounds on FCNC processes. Thus, most model builders impose restrictions on the structure of the Higgs fermion couplings to avoid the potential for phenomenological disaster. However, even in the case of the most general Higgs-fermion couplings, parameter regimes exist where FCNC effects are sufficiently under control. In the absence of new physics beyond the 2HDM, such parameter regimes are unnatural (but can be arranged with fine-tuning). In models such as the minimal supersymmetric extension of the Standard Model (MSSM), supersymmetry-breaking effects generate all possible Higgs-fermion Yukawa couplings allowed by electroweak gauge invariance. Nevertheless, the FCNC effects are one-loop suppressed and hence phenomenologically acceptable.

---

[11] The Feynman rule for the ZZh_k vertex includes a factor of two relative to the coefficient of the corresponding term in $i\mathscr{L}_{VVH}$ due to the identical Z bosons. The Feynman rule for the Zh_jh_k vertex is given by $\frac{1}{2}(g/c_W)\text{Im}[q_{j1}q^*_{k1} + q_{j2}q^*_{k2}](p_k - p_j)^\mu$. Here, the factor of two relative to the corresponding term in eq. (2.67) arises from the implicit double sum over j and k in the Lagrangian. Note that the rule for the Zh_jh_k vertex does not depend on the ordering of j and k.

In this section, we will study the basis-independent description of the Higgs-fermion interaction. In a generic basis, the so-called type-III model [11, 29, 34] of Higgs fermion interactions is governed by the following interaction Lagrangian:

$$-\mathscr{L}_{\rm Y} = \overline{Q^0_L}\,\widetilde{\Phi}_1\eta_1^{U,0}\,U^0_R + \overline{Q}^0_L\,\Phi_1(\eta_1^{D,0})^\dagger\,D^0_R + \overline{Q^0_L}\,\widetilde{\Phi}_2\eta_2^{U,0}\,U^0_R + \overline{Q^0_L}\,\Phi_2(\eta_2^{D,0})^\dagger\,D^0_R + \text{h.c.}\,, \tag{2.73}$$

where $\Phi_{1,2}$ are the Higgs doublets and $\widetilde{\Phi}_i \equiv i\sigma_2\Phi_i^*$. As in section 1.1.3, Q^0_L, U^0_R, D^0_R denote the interaction basis quark fields, which are vectors in the quark flavor space, and $\eta_1^{Q,0}$ and $\eta_2^{Q,0}$ ($Q = U\,,\,D$) are four 3×3 matrices in quark flavor space. We have omitted the leptonic couplings in eq. (2.73); these are obtained from eq. (2.73) with the obvious substitutions $Q^0_L \to L^0_L$ and $D^0_R \to E^0_R$. (In the absence of right-handed neutrinos, there is no analog of U^0_R.)

The derivation of the couplings of the physical Higgs bosons with the quark mass-eigenstates was given in ref. [34] in the case of a CP-conserving Higgs sector. Here, we generalize that discussion to the more general case of a CP-violating Higgs sector. The first step is to identify the quark mass-eigenstates. This is accomplished by setting the scalar fields to their vacuum expectation values and performing unitary transformations of the left and right-handed up and down quark multiplets such that the resulting quark mass matrices are diagonal with non-negative entries. In more detail, we define left-handed and right-handed quark mass-eigenstate fields according to eq. (1.19), with the CKM matrix $K \equiv V_L^U V_L^{D\,\dagger}$ as before. In addition, we introduce "rotated" Yukawa coupling matrices:

$$\eta_a^U \equiv V_L^U\,\eta_a^{U,0}\,V_R^{U\,\dagger}\,, \qquad\qquad \eta_a^D \equiv V_R^D\,\eta_a^{D,0}\,V_L^{D\,\dagger}\,. \tag{2.74}$$

We then rewrite eq. (2.73) in terms of the quark mass-eigenstate fields and the transformed couplings:

$$-\mathscr{L}_{\rm Y} = \overline{U}_L\widetilde{\Phi}_{\bar a}\eta_a^U U_R + \overline{D}_L K^\dagger\widetilde{\Phi}_{\bar a}\eta_a^U U_R + \overline{U}_L K\Phi_a\eta_{\bar a}^{D\,\dagger}D_R + \overline{D}_L\Phi_a\eta_{\bar a}^{D\,\dagger}D_R + \text{h.c.}\,, \tag{2.75}$$

where $\eta_a^Q \equiv (\eta_1^Q\,,\,\eta_2^Q)$ is a (basis-dependent) vector in U(2) space. If we assume that there is no basis in which can write either $\eta_2^U = \eta_2^D = 0$ (Type I) or $\eta_1^U = \eta_2^D = 0$ (Type II), then this "Type III" 2HDM Yukawa Lagrangian. One could also write it in a more compact form:

$$-\mathscr{L}_{\rm Y} = \overline{\mathcal{Q}_L}\widetilde{\Phi}_{\bar a}\eta_a^U\mathcal{U}_R + \overline{\mathcal{Q}_L}\Phi_a\eta_{\bar a}^{D\,\dagger}D_R + \text{h.c.}\,, \tag{2.76}$$

where $\mathcal{U} \equiv K^\dagger U$, and $\mathcal{Q}_L = \binom{\mathcal{U}}{D}_L$.

Under a U(2)-transformation of the scalar fields, $\eta^Q_a \to U_{a\bar{b}}\eta^Q_b$ and $\eta^{Q\,\dagger}_{\bar{a}} \to \eta^{Q\,\dagger}_{\bar{b}} U^\dagger_{b\bar{a}}$. Hence, the Higgs–quark Lagrangian is U(2)-invariant. We can construct basis-independent couplings following the strategy of section 2.3 by transforming to the Higgs basis. Using eq. (2.11), we can rewrite eq. (2.75) in terms of Higgs basis scalar fields:

$$-\mathscr{L}_{\rm Y} = \overline{\mathcal{Q}_L}(\tilde{H}_1\kappa^U + \tilde{H}_2\rho^U)\mathcal{U}_R + \overline{\mathcal{Q}_L}(H_1\kappa^{D\,\dagger} + H_2\rho^{D\,\dagger})D_R + \text{h.c.}\,, \tag{2.77}$$

where

$$\kappa^Q \equiv \widehat{v}^*_{\bar{a}}\,\eta^Q_a\,, \qquad\qquad \rho^Q \equiv \widehat{w}^*_{\bar{a}}\,\eta^Q_a\,. \tag{2.78}$$

Inverting eq. (2.78) yields:

$$\eta^Q_a = \kappa^Q\widehat{v}_a + \rho^Q\widehat{w}_a\,. \tag{2.79}$$

Under a U(2) transformation, κ^Q is invariant, whereas ρ^Q is a pseudo-invariant that transforms as:

$$\rho^Q \to (\det U)\rho^Q\,. \tag{2.80}$$

By construction, κ^U and κ^D are proportional to the (real non-negative) diagonal quark mass matrices M_U and M_D, respectively. In particular, the M_Q are obtained by inserting eq. (2.12) into eq. (2.77). As in the SM (see eqs. (1.23) and (1.24)), we find:

$$M_U = \frac{v}{\sqrt{2}}\kappa^U \quad = \quad \text{diag}(m_u\,,\, m_c\,,\, m_t) = V^U_L M^0_U V^{U\,\dagger}_R\,, \tag{2.81}$$

$$M_D = \frac{v}{\sqrt{2}}\kappa^{D\,\dagger} \quad = \quad \text{diag}(m_d\,,\, m_s\,,\, m_b) = V^D_L M^0_D V^{D\,\dagger}_R\,, \tag{2.82}$$

where $M^0_U \equiv (v/\sqrt{2})\widehat{v}^*_{\bar{a}}\,\eta^{U,0}_a$ and $M^0_D \equiv (v/\sqrt{2})\widehat{v}_a\,\eta^{D,0\,\dagger}_{\bar{a}}$. That is, we have chosen the unitary matrices V^U_L, V^U_R, V^D_L and V^D_R such that M_D and M_U are diagonal matrices with real non-negative entries.[12] In contrast, the ρ^Q are independent complex 3×3 matrices.

In order to obtain the interactions of the physical Higgs bosons with the quark mass-eigenstates, we do not require the intermediate step involving the Higgs basis. Instead, we insert eq. (2.47) into eq. (2.75) and obtain:

$$\begin{aligned}
-\mathscr{L}_Y = {} & \frac{1}{v}\overline{D}\left\{ M_D(q_{k1}P_R + q^*_{k1}P_L) + \frac{v}{\sqrt{2}}\left[q_{k2}\,[e^{i\theta_{23}}\rho^D]^\dagger P_R + q^*_{k2}\,e^{i\theta_{23}}\rho^D P_L\right]\right\}Dh_k \\
& + \frac{1}{v}\overline{U}\left\{ M_U(q_{k1}P_L + q^*_{k1}P_R) + \frac{v}{\sqrt{2}}\left[q^*_{k2}\,e^{i\theta_{23}}\rho^U P_R + q_{k2}\,[e^{i\theta_{23}}\rho^U]^\dagger P_L\right]\right\}Uh_k \\
& + \left\{\overline{U}\left[K[\rho^D]^\dagger P_R - [\rho^U]^\dagger K P_L\right] DH^+ \right. \\
& \qquad \left. + \frac{\sqrt{2}}{v}\,\overline{U}\,[K M_D P_R - M_U K P_L]\,DG^+ + \text{h.c.}\right\},
\end{aligned} \tag{2.83}$$

---

[12] This can be accomplished by the singular-value decompositions of the complex matrices M^0_U and M^0_D [73].

where $k = 1, \ldots 4$. Since $e^{i\theta_{23}}\rho^Q$ and $[\rho^Q]^\dagger H^+$ are U(2)-invariant, it follows that eq. (2.83) is a basis-independent representation of the Higgs–quark interactions.

The neutral Goldstone boson interactions ($h_4 \equiv G^0$) are easily isolated:

$$-\mathscr{L}_{YG} = \frac{i}{v}\left[\overline{D} M_D \gamma_5 D - \overline{U} M_U \gamma_5 U\right] G^0 \,. \tag{2.84}$$

In addition, since the q_{k1} are real for $k = 1, 2, 3$, it follows that the piece of the neutral Higgs–quark couplings proportional to the quark mass matrix is of the form $v^{-1}\overline{Q}\, M_Q\, q_{k1}\, Q\, h_k$.

The couplings of the neutral Higgs bosons to quark pairs are generically CP-violating as a result of the complexity of the q_{k2} and the fact that the matrices $e^{i\theta_{23}}\rho^Q$ are not generally purely real orpurely imaginary. (Invariant conditions for the CP-invariance of these couplings are given in Chapter 3.1). Eq. (2.83) also exhibits Higgs-mediated FCNCs at tree-level due to the ρ^Q not being flavor-diagonal. Thus, for a phenomenologically acceptable theory, the off-diagonal elements of ρ^Q must be small.

Chapter 3

Special Limits: CP Conservation, The 2HDM with $Z_6 = 0$, and Custodial Symmetry

In the limit of $g' \to 0$, the bosonic sector of the Standard Model has a global $SU(2)_L \times SU(2)_R$ symmetry. After electroweak symmetry breaking, this symmetry reduces to $SU(2)_{L+R} \equiv SU(2)_V$, known as the "custodial symmetry." Violations of custodial symmetry lead to corrections to the relation $m_W^2 = m_Z^2 cos^2\theta_W$. This custodial limit is a more restrictive case of CP conservation. In the 3-generation model, we know that CP violation must exist in the CKM matrix, so a 2HDM with no CP violation is unrealistic. Thus, for the purposes of this chapter, when we refer to the "CP conserving" model, we mean that the bosonic sector and the neutral Higgs-Quark couplings conserve CP.

The CP-conserving limit of the basis-independent 2HDM has been analyzed in [34] and [66][1]. In this chapter, we build on the results of previous work and define two cases of CP conservation that are distinguishable based on the parameter $Z_6 e^{-i\theta_{23}}$. In section 3.1.1, we derive basis-independent expressions for the masses and mixing angles of the Higgs particles in the CP-conserving limit. In section 3.1.2, we make contact with the existing literature by working in the real basis and relating the Higgs fields to the parameters $c_{\beta-\alpha}$ and $s_{\beta-\alpha}$.

We then analyze the scenario in which $Z_6 = 0$. We will start by deriving the mass matrix and invariant expressions for the mixing angles in section 3.2, and then discuss the effects of CP violation or conservation in the scalar couplings. We end with a discussion of the special case $Z_6 = Z_7 = 0$ in section 3.2.3.

In section 3.3, we apply these results to analyse the custodial limit of the 2HDM. The use of the basis-independent formalism allows us to clarify aspects of custodial symmetry which have been made unnecessarily complicated in the literature. We derive unambiguous conditions for custodial symmetry in the scalar sector and the

---

[1] By CP conservation, we mean that the neutral scalars are CP eigenstates. The complex phase in the CKM matrix induces CP violation in the charged Higgs interactions.

Higgs-Quark sector and discuss the resulting implications for the scalar masses.

3.1 The CP-Conserving 2HDM

3.1.1 Basis-Independent Analysis of the CP-Conserving Limit

In the CP-conserving limit, we impose CP-invariance on all bosonic couplings of the Higgs bosons and the fermionic couplings of the neutral Higgs bosons. (We will ignore, for now, the CP violation in charged Higgs-quark interactions that arises from the complexity of the CKM matrix.) The requirement of a CP-conserving bosonic sector is equivalent to the requirement that the scalar potential is explicitly CP-conserving and that the Higgs vacuum is CP-invariant (*i.e.*, there is no spontaneous CP-violation). Basis-independent conditions for a CP-conserving bosonic sector have been given in refs. [20, 34, 60, 88]. In ref. [34], these conditions were recast into the following form. The bosonic sector is CP-conserving if and only if:[2]

$$\mathrm{Im}[Z_6 Z_7^*] = \mathrm{Im}[Z_5^* Z_6^2] = \mathrm{Im}[Z_5^*(Z_6+Z_7)^2] = 0\,. \tag{3.1}$$

Eq. (3.1) is equivalent to the requirement that

$$\sin 2(\theta_5-\theta_6) = \sin 2(\theta_5-\theta_7) = \sin(\theta_6-\theta_7) = 0\,, \tag{3.2}$$

where θ_5 and θ_6 are defined in eq. (C.7) and $\theta_7 \equiv \arg Z_7$ (note that θ_5 is defined modulo π and θ_6 and θ_7 are defined modulo 2π).

One can explore the consequences of CP-invariance by studying the pattern of Higgs couplings and the structure of the neutral Higgs boson squared-mass matrix [eq. (2.24)]. The tree-level couplings of G^0 are CP-conserving, even in the general CP-violating 2HDM. In particular, the couplings $G^0G^0G^0$, $G^0G^+G^-$, $G^0H^+H^-$ and ZZG^0 are absent. Moreover, eq. (2.84) implies that G^0 possesses purely pseudoscalar couplings to the fermions. Hence, G^0 is a CP-odd scalar, independently of the structure of the scalar potential. We can therefore use the couplings of G^0 to the neutral Higgs bosons as a probe of the CP-quantum numbers of these states. The analysis of the neutral Higgs boson squared-mass matrix (which does not depend on Z_7) simplifies significantly when $\mathrm{Im}[Z_5^* Z_6^2] = 0$. One can then choose a basis where Z_5 and Z_6 are simultaneously real, in which case the scalar squared-mass matrix decomposes into diagonal block form. The upper 2×2 block can be diagonalized analytically and yields the mass-eigenstates h^0 and H^0 (with $m_{h^0} \leq m_{H^0}$). The lower 1×1 block yields the mass-eigenstate A^0. If all the conditions of eqs. (3.1) and (3.28) are satisfied, then the neutral Higgs boson mass-eigenstates are also states of definite CP quantum number. We shall demonstrate below that h^0 and H^0 are CP-even scalars and A^0 is a CP-odd scalar.

Since $Z_6 \neq 0$ by assumption, eq. (C.24) yields $\sin\phi\cos\phi = 0$, and eq. (C.21) implies that either some of the neutral Higgs boson masses are degenerate or $s_{13}s_{12}c_{12} = 0$.[3] In the case of degenerate masses, some of the invariant angles are not well defined, since any linear combination of the degenerate states is also a mass-eigenstate. Hence, the degenerate case must be treated separately. In what follows, we shall assume that all three neutral Higgs boson masses are non-degenerate. Note that if $\sin\phi = 0$, then eq. (C.23) yields $s_{13} = 0$, whereas if $\cos\phi = 0$, then eq. (C.25) yields $\sin 2\theta_{12} = 0$.[4] Thus, we shall consider separately the two cases:

---

[2] Since the scalar potential minimum conditions imply that $Y_3 = -\frac{1}{2}Z_6v^2$, no separate condition involving Y_3 is required.

[3] Since $Z_6 \neq 0$, one can use eqs. (C.9) and (C.10) to show that $c_{13} \neq 0$.

$$\begin{aligned} &\text{CP Case I}: && \sin\phi = 0 \quad\Longrightarrow\quad \mathrm{Im}(Z_5 e^{-2i\theta_{23}}) = \mathrm{Im}(Z_6 e^{-i\theta_{23}}) = 0\,, && (3.3)\\ &\text{CP Case II}: && \cos\phi = 0 \quad\Longrightarrow\quad \mathrm{Im}(Z_5 e^{-2i\theta_{23}}) = \mathrm{Re}(Z_6 e^{-i\theta_{23}}) = 0\,, && (3.4)\end{aligned}$$

where $\phi \equiv \arg(Z_6 e^{-i\theta_{23}})$ is an invariant quantity. The first case corresponds to the mass ordering $m_{A^0} > m_{H^0}$; the second case to the reverse. By comparing equations (3.3) and (3.4) with the mass matrix in eq. (2.41), one identifies the CP-odd field as the following:

$$A^0 = \begin{cases} \mathrm{Im}(e^{i\theta_{23}} H_2^0) & [\text{CP Case I}]\,,\\ \mathrm{Re}(e^{i\theta_{23}} H_2^0) & [\text{CP Case II}]\,. \end{cases} \tag{3.5}$$

This is equivalent to the statement that the fields in the Higgs basis transform as follows:

$$\begin{aligned} \begin{pmatrix} H_1 \\ e^{i\theta_{23}} H_2 \end{pmatrix} &\rightarrow \begin{pmatrix} H_1^* \\ e^{-i\theta_{23}} H_2^* \end{pmatrix} \quad [\text{Case I}]\,,\\ \begin{pmatrix} H_1 \\ i e^{i\theta_{23}} H_2 \end{pmatrix} &\rightarrow \begin{pmatrix} H_1^* \\ (i e^{i\theta_{23}} H_2)^* \end{pmatrix} \quad [\text{Case II}]\,. \end{aligned} \tag{3.6}$$

Alternatively, one can define a CP-transformation on fields in the generic basis using the following [66]:

$$\begin{pmatrix} H_1 \\ H_2 \end{pmatrix} = \begin{pmatrix} \widehat{v}_1^* & \widehat{v}_2^* \\ -\widehat{v}_2 & \widehat{v}_1 \end{pmatrix} \begin{pmatrix} \Phi_1 \\ \Phi_2 \end{pmatrix}. \tag{3.7}$$

Substituting eq. (3.7) into eq. (3.6) yields

$$\begin{pmatrix} \widehat{v}_1^* & \widehat{v}_2^* \\ -\widehat{v}_2 e^{i\theta_{23}} & \widehat{v}_1 e^{i\theta_{23}} \end{pmatrix} \begin{pmatrix} \Phi_1 \\ \Phi_2 \end{pmatrix} \rightarrow \begin{pmatrix} \widehat{v}_1 & \widehat{v}_2 \\ \mp \widehat{v}_2^* e^{-i\theta_{23}} & \pm \widehat{v}_1^* e^{-i\theta_{23}} \end{pmatrix} \begin{pmatrix} \Phi_1^* \\ \Phi_2^* \end{pmatrix}, \tag{3.8}$$

or

$$\begin{aligned} \begin{pmatrix} \Phi_1 \\ \Phi_2 \end{pmatrix} &\rightarrow \begin{pmatrix} \widehat{v}_1 & -\widehat{v}_2^* e^{-i\theta_{23}} \\ \widehat{v}_2 & \widehat{v}_1^* e^{-i\theta_{23}} \end{pmatrix} \begin{pmatrix} \widehat{v}_1 & \widehat{v}_2 \\ \mp \widehat{v}_2^* e^{-i\theta_{23}} & \pm \widehat{v}_1^* e^{-i\theta_{23}} \end{pmatrix} \begin{pmatrix} \Phi_1^* \\ \Phi_2^* \end{pmatrix} \\ &= \begin{pmatrix} \widehat{v}_1^2 \pm \widehat{v}_2^{*2} e^{-2i\theta_{23}} & \widehat{v}_1 \widehat{v}_2 \mp \widehat{v}_2^* \widehat{v}_1^* e^{-2i\theta_{23}} \\ \widehat{v}_1 \widehat{v}_2 \mp \widehat{v}_2^* \widehat{v}_1^* e^{-2i\theta_{23}} & \widehat{v}_2^2 \pm \widehat{v}_1^{*2} e^{-2i\theta_{23}} \end{pmatrix} \begin{pmatrix} \Phi_1^* \\ \Phi_2^* \end{pmatrix}. \end{aligned} \tag{3.9}$$

Hence, the "covariant" form of eq. (3.6) is

$$\Phi_a(\vec{x}, t) \rightarrow (\widehat{v}_a \widehat{v}_b \pm e^{-2i\theta_{23}} \widehat{w}_a \widehat{w}_b) \Phi_{\bar{b}}^*(-\vec{x}, t), \tag{3.10}$$

---

[4] The same constraints are obtained by imposing the requirement of CP-conserving Higgs couplings. In particular, the existence of a $G^0 h_k h_k$ coupling would imply that h_k is a state of mixed CP-even and CP-odd components. All such couplings must therefore be absent in the CP-conserving limit. Using the results of eqs. (2.59) and (3.2) one can easily check that at least one of these CP-violating couplings is present unless $s_{13} = \sin\phi = 0$ or $\cos\phi = \sin 2\theta_{12} = 0$.

with the positive (negative) solution corresponding to CP Case I (II).

As a consistency check, we note that a CP-transformation of the Higgs doublets in the generic basis takes the following form (in the notation of [60]):

$$\mathcal{CP}\ \Phi_a(\vec{x},t)\ \mathcal{CP}^{-1} = (U^{CP})_{ab}\Phi^*_{\bar{b}}(-\vec{x},t). \tag{3.11}$$

Invariance of the vacuum under CP requires [44]:

$$< \Phi_a >= (U^{CP})_{ab} < \Phi_{\bar{b}} >^* . \tag{3.12}$$

Eq. (3.10) indeed satisfies eq. (3.12), with

$$(U^{CP})_{ab} = \widehat{v}_a\widehat{v}_b \pm e^{-2i\theta_{23}}\,\widehat{w}_a\widehat{w}_b. \tag{3.13}$$

When the scalar potential is CP-conserving, there always exists a basis in which the Y and Z parameters and the scalar vacuum expectation values are all real-valued. In a generic basis, the quantities Y_3, Z_5, Z_6 and Z_7 are complex. For the CP-transformation given in eq. (3.10), requiring $\mathcal{CP}\ \mathcal{V}\ \mathcal{CP}^{-1} = \mathcal{V}$ reproduces the relations in eq. (3.1). If all three physical neutral fields couple to CP-even states, eg, $H^+H^-h_k \neq 0$ and $W^+W^-h_k \neq 0\ \forall\ k$, then the scalar sector violates CP. Otherwise, the field h_k for which all such couplings vanish is CP-odd, and the remaining two fields are CP-even. In our basis-independent notation, the quantity q_{k1} will be non-zero for the CP-even states and zero for the CP-odd state.

Assuming that the masses are non-degenerate[5] and that Z_6 is non-zero, we find that

$$\begin{aligned}
&\text{If } h_3 \text{ is CP odd}, & s_{13} &= \mathrm{Im}(Z_6\,e^{-i\theta_{23}}) = \sin\phi = 0 \quad \text{[CP Case I]}, & (3.14)\\
&\text{If } h_2 \text{ is CP odd}, & s_{12} &= \mathrm{Re}(Z_6\,e^{-i\theta_{23}}) = \cos\phi = 0 \quad \text{[CP Case IIa]}, & (3.15)\\
&\text{If } h_1 \text{ is CP odd}, & c_{12} &= \mathrm{Re}(Z_6\,e^{-i\theta_{23}}) = \cos\phi = 0 \quad \text{[CP Case IIb]}. & (3.16)
\end{aligned}$$

The values of the $q_{k\ell}$ corresponding to cases I, IIa and IIb are given in Tables 3.1—3.3.

Table 3.1: The U(2)-invariant quantities $q_{k\ell}$ in the CP-conserving limit. Case I: $s_{13} = \mathrm{Im}(Z_6\,e^{-i\theta_{23}}) = \sin\phi = 0$. G^0 and h_3 are CP-odd; h_1 and h_2 are CP-even.

k	q_{k1}	q_{k2}
1	c_{12}	$-s_{12}$
2	s_{12}	c_{12}
3	0	i
4	i	0

---

[5] One can investigate separately cases of degenerate masses, in which case not all of the mixing angles are well-defined, which we will not do here for the general case of $Z_6 \neq 0$. We analyze the degenerate cases for $Z_6 = 0$ in section 3.2.2.

Table 3.2: The U(2)-invariant quantities $q_{k\ell}$ in the CP-conserving limit. Case IIa: $s_{12} = \mathrm{Re}(Z_6\, e^{-i\theta_{23}}) = \cos\phi = 0$. G^0 and h_2 are CP-odd; h_1 and h_3 are CP-even.

k	q_{k1}	q_{k2}
1	c_{13}	$-is_{13}$
2	0	1
3	s_{13}	ic_{13}
4	i	0

Table 3.3: The U(2)-invariant quantities $q_{k\ell}$ in the CP-conserving limit. Case IIb: $c_{12} = \mathrm{Re}(Z_6\, e^{-i\theta_{23}}) = \cos\phi = 0$. G^0 and h_1 are CP-odd; h_2 and h_3 are CP-even.

k	q_{k1}	q_{k2}
1	0	1
2	$-c_{13}$	is_{13}
3	s_{13}	ic_{13}
4	i	0

In both Case I and Case II, $\widetilde{\mathcal{M}}$ assumes a block diagonal form consisting of a 2×2 block (corresponding to the the CP-even Higgs bosons) and a 1×1 block (corresponding to the CP-odd Higgs boson). The CP-odd field has mass

$$\begin{aligned} m^2_{A^0} &= \tfrac{1}{2}v^2\left[Y_2/v^2 + Z_1 + \tfrac{1}{2}(Z_3 + Z_4 - \mathrm{Re}(Z_5\, e^{-2i\theta_{23}}))\right] \qquad \text{[Case I]}\ , \\ m^2_{A^0} &= \tfrac{1}{2}v^2\left[Y_2/v^2 + Z_1 + \tfrac{1}{2}(Z_3 + Z_4 + \mathrm{Re}(Z_5\, e^{-2i\theta_{23}}))\right] \qquad \text{[Case II]}\ . \end{aligned} \tag{3.17}$$

It is possible to eliminate the explicit dependence on θ_{23} by defining a quantity ε_{56} as follows:

$$\mathrm{Re}(Z_5^* Z_6^2) = \varepsilon_{56}|Z_5|\,|Z_6|^2\,, \qquad \varepsilon_{56} \equiv \pm 1\,. \tag{3.18}$$

Note that in the CP-conserving limit,

$$\mathrm{Re}[Z_5^* Z_6^2] = \mathrm{Re}(Z_5\, e^{-2i\theta_{23}})\,\mathrm{Re}[(Z_6 e^{-i\theta_{23}})^2] = \pm\mathrm{Re}(Z_5\, e^{-2i\theta_{23}})|Z_6|^2\,, \tag{3.19}$$

where the upper (lower) sign corresponds to case I (II). Then can write $\mathrm{Re}(Z_5\, e^{-2i\theta_{23}}) = \pm\varepsilon_{56}|Z_5|$, and the masses of the neutral Higgs fields become

$$\begin{aligned} m^2_{h^0,H^0} &= \tfrac{1}{2}v^2\Big[Y_2/v^2 + Z_1 + \tfrac{1}{2}(Z_3 + Z_4 + \varepsilon_{56}|Z_5|) \\ &\qquad \mp\sqrt{\left[Y_2/v^2 - Z_1 + \tfrac{1}{2}(Z_3 + Z_4 + \varepsilon_{56}|Z_5|)\right]^2 + 4|Z_6|^2}\,\Big]\,, \\ m^2_{A^0} &= Y_2 + \tfrac{1}{2}(Z_3 + Z_4 - \varepsilon_{56}|Z_5|)v^2\,, \end{aligned} \tag{3.20}$$

where is defined by the relation ε_{56} is an invariant quantity, its value must be determined from experiment.

Additional constraints for a CP-conserving 2HDM arise when the Higgs-fermion couplings are included. Let us write the transformation of the fields h_k under CP as $h_k \to \eta_k h_k$, where $\eta_k = \pm 1$. The Higgs-Quark Lagrangian [eq. (2.83)] contains the term

$$\frac{1}{\sqrt{2}} q_{k2}^* \, \overline{D} e^{i\theta_{23}} \rho^D P_L D h_k + h.c. \tag{3.21}$$

(and a similar term with U in place of D). If CP is conserved, the quark mass-eigenstates D_i and U_i transform under CP with some phase,

$$D_i \to e^{i\theta_i^D} D_i^*, \qquad U_i \to e^{i\theta_i^U} U_i^* \,. \tag{3.22}$$

However, one can rephase D_i and U_i such that $D_i \to D_i^*$ and $U_i \to U_i^*$ under a CP transformation. Let us assume that we have done such a rephasing, defined by $D_i \to \eta_i^D D_i$ (and similarly for U_i), with $|\eta_i^Q| = 1$. This rephasing also transforms the Yukawa matrices, with the result $\rho_{ij}^Q \to \eta_i^Q \eta_j^{Q*} \rho_{ij}^Q$. Reinstituting the flavor indices, the CP transformation on eq. (3.21) (having suitably rephased the quark fields) gives

$$q_{k2}^* \, \overline{D}_i [e^{i\theta_{23}} \rho^D]_{ij} P_L D_j h_k \to \eta_k q_{k2}^* \, \overline{D}_i [e^{i\theta_{23}} \rho^D]_{ji} P_R D_j h_k \,, \tag{3.23}$$

where ρ^D has been appropriately rephased, as described above. Comparing this to the hermitian conjugate of eq. (3.21),

$$q_{k2} \, \overline{D}_i [e^{i\theta_{23}} \rho^D]_{ij}^\dagger P_R D_j h_k \,, \tag{3.24}$$

we obtain a condition for CP invariance of the neutral Higgs bosons, and with the analogous result for ρ^U. Both conditions can be summarized as

$$\eta_k q_{k2}^* [e^{i\theta_{23}} \rho^Q]_{ij} = [e^{i\theta_{23}} \rho^Q]_{ij}^* q_{k2} \,. \tag{3.25}$$

The values of q_{k2} and η_k can be obtained from Tables 3.1—3.3. One finds that eq. (3.25) is equivalent to the following:

$$e^{i\theta_{23}} \rho^Q \text{ is } \begin{cases} \text{real} & \text{in Case I}\,, \\ \text{imaginary} & \text{in Cases IIa and IIb}\,. \end{cases} \tag{3.26}$$

In both Cases I and II, the results of eqs. (3.3), (3.4) and (3.26) imply that

$$\begin{aligned} \text{Im}(Z_6 \rho^Q) &= \text{Im}(Z_6 e^{-i\theta_{23}} e^{i\theta_{23}} \rho^Q) \\ &= \text{Re}(Z_6 e^{-i\theta_{23}}) \text{Im}(e^{i\theta_{23}} \rho^Q)) - \text{Im}(Z_6 e^{-i\theta_{23}}) \text{Re}(e^{i\theta_{23}} \rho^Q)) \\ &= 0\,. \end{aligned} \tag{3.27}$$

One can prove similar conditions involving Z_7 and Z_5, so that the complete set of conditions for CP-invariance of the couplings of the neutral Higgs bosons to fermion pairs is the following:

$$\text{Im}[Z_6 \rho^Q] = \text{Im}[Z_7 \rho^Q] = \text{Im}[Z_5 (\rho^Q)^2] = 0\,. \tag{3.28}$$

Thus, if eqs. (3.1) and (3.28) are satisfied, then the neutral Higgs bosons are eigenstates of CP, and the only possible source of CP-violation in the 2HDM is the unremovable phase in the CKM matrix K that enters via the charged current interactions mediated by either $W^\pm$ or $H^\pm$ exchange[6] [see eq. (2.83)].

Invariant techniques for describing the constraints on the Higgs-fermion interaction due to CP-invariance have also been considered in refs. [20] and [44]. In these works, the authors construct invariant expressions that are both U(2)-invariant and invariant with respect to the redefinition of the quark fields. For example, the invariants denoted by J_a and J_b in ref. [20] are given by $J_a \equiv \mathrm{Im}\, J^D$ and $J_b \equiv \mathrm{Im}\, J^U$ where

$$J^Q = \mathrm{Tr}(VYT^Q)\,, \qquad T^Q_{a\bar{b}} \equiv \mathrm{Tr_f}(\eta_a^{Q,0}\eta_{\bar{b}}^{Q,0\,\dagger}) = \mathrm{Tr_f}(\eta_a^{Q}\eta_{\bar{b}}^{Q\,\dagger})\,, \tag{3.29}$$

and the trace $\mathrm{Tr_f}$ sums over the diagonal quark generation indices. Note that the trace over generation indices ensures that the resulting expression is invariant with respect to unitary redefinitions of the quark fields [eq. (1.19)]. Using eq. (B.5) [with $A = Y$], it is straightforward to re-express eq. (3.29) as:

$$J^Q = Y_1 \mathrm{Tr_f}[(\kappa^Q)^2] + Y_3 \mathrm{Tr_f}[\kappa^Q \rho^Q]\,, \tag{3.30}$$

after using eqs. (2.17), (2.18) and (2.78). Indeed, J^Q is invariant with respect to U(2) transformations since the product of pseudo-invariants $Y_3\,\rho^Q$ is a U(2)-invariant quantity. Moreover, taking the trace over the quark generation indices ensures that J^Q is invariant with respect to unitary redefinitions of the quark fields. In ref. [20], a proof is given that $\mathrm{Im}\, J^Q = 0$ is one of the invariant conditions for CP-invariance of the Higgs-fermion interactions. In our formalism, this result is easily verified. Using the scalar potential minimum conditions [eq. (2.21)], we obtain:

$$\begin{aligned} \mathrm{Im}\, J^Q &= -\frac{v}{\sqrt{2}}\,\mathrm{Im}\big[Z_6 \mathrm{Tr_f}(M_Q \rho^Q)\big]\,, \\ &= -\frac{v}{\sqrt{2}}\,\mathrm{Tr_f}\mathrm{Im}\big[M_Q Z_6 \rho^Q\big]\,. \end{aligned} \tag{3.31}$$

But, CP-invariance requires [by eq. (3.28)] that $Z_6\rho^Q$ is real. Since M_Q is a real diagonal matrix, it then immediately follows that $\mathrm{Im}\, J^Q = 0$.

3.1.2 The CP-Conserving Limit in the Real Basis

For completeness, we will now analyse the CP-conserving 2HDM in a specific basis, in order to express the masses and physical Higgs states in terms of more traditional parameters. In the standard notation of the CP-conserving 2HDM, one considers only real basis choices, in which the Higgs Lagrangian parameters and the scalar vacuum expectation values are real. We can therefore restrict basis changes to O(2) transformations [34].[7] In this context, pseudo-invariants are SO(2)-invariant quantities that change sign under an O(2) transformation with determinant equal to -1. Note that Z_5 is now an invariant with respect to O(2) transformations, but Z_6, Z_7 and $e^{-i\theta_{23}}$ are pseudo-invariants. In particular, for $Z_6 \neq 0$ in the convention where $0 \leq \phi < \pi$,

---

[6] One can also formulate a basis-independent condition (that is invariant with respect to separate redefinitions of the Higgs doublet fields and the quark fields) for the absence of CP-violation in the charged current interactions. This condition involves the Jarlskog invariant [79, 80], and can also be written as [16, 44]: $\mathrm{Tr_f}\left[H^{U,0},\, H^{D,0}\right]^3 = 0$ (summed over three quark generations), where $H^{Q,0} \equiv M^{Q,0}M^{Q,0\,\dagger}$ and the $M^{Q,0}$ are defined below eq. (1.24). Since CP-violating phenomena in the charged current interactions are observed and well described by the CKM matrix, we shall not impose this latter condition here.

$$e^{-i\theta_{23}} = e^{i\phi} e^{-i\theta_6} = \begin{cases} \varepsilon_6 & [\text{Case I}]\,, \\ i\varepsilon_6 & [\text{Case II}]\,, \end{cases} \tag{3.32}$$

where $Z_6 \equiv \varepsilon_6 |Z_6|$ in the real basis. That is, ε_6 is a pseudo-invariant quantity (in contrast, the sign of Z_5 is invariant) with respect to O(2) transformations. Note that in the real basis,

$$\varepsilon_{56} = e^{2i\theta_5} e^{-2i\theta_6} = \mathrm{sgn(Z_5)}\varepsilon_6^2 = \mathrm{sgn(Z_5)}\,, \tag{3.33}$$

so that eq. (3.20) can be written in terms of the real-basis parameters:

$$m_{A^0}^2 = Y_2 + \tfrac{1}{2} v^2 \left(Z_3 + Z_4 - Z_5\right)\,. \tag{3.34}$$

The generic real basis fields can be expressed in terms of the two neutral CP-even scalar mass-eigenstates h^0, H^0 (with $m_{h^0} \leq m_{H^0}$) and the CP-odd scalar mass-eigenstate A^0, G^0 as follows [57, 58, 61]:

$$\Phi_1^0 = \frac{1}{\sqrt{2}} \left[v\widehat{v}_1 - h^0 s_\alpha + H^0 c_\alpha + i(G^0 c_\beta - A^0 s_\beta) \right]\,, \tag{3.35}$$

$$\Phi_2^0 = \frac{1}{\sqrt{2}} \left[v\widehat{v}_2 + h^0 c_\alpha + H^0 s_\alpha + i(G^0 s_\beta + A^0 c_\beta) \right]\,, \tag{3.36}$$

with $m_{h^0} \leq m_{H^0}$, where $\widehat{v}_a = (c_\beta\,, s_\beta)$, $s_\alpha \equiv \sin\alpha$, $c_\alpha \equiv \cos\alpha$, and α is the CP-even neutral Higgs boson mixing angle. These equations can be written more compactly as

$$\Phi_a^0 = \frac{1}{\sqrt{2}} \left[(v + h^0 s_{\beta-\alpha} + H^0 c_{\beta-\alpha} + iG^0)\widehat{v}_a + (h^0 c_{\beta-\alpha} - H^0 s_{\beta-\alpha} + iA^0)\widehat{w}_a \right]\,, \tag{3.37}$$

where $s_{\beta-\alpha} \equiv \sin(\beta-\alpha)$ and $c_{\beta-\alpha} \equiv \cos(\beta-\alpha)$.

Using the results of Tables 3.1—3.3 and comparing eq. (3.37) to eq. (2.47) [with $e^{-i\theta_{23}}$ determined from eq. (3.32)], one can identify the neutral Higgs fields h_k with the eigenstates of definite CP quantum numbers, h^0, H^0 and A^0, and relate the angular factor $\beta-\alpha$ with the appropriate invariant angle:[8]

---

[7] If $Z_6 = Z_7 = \rho^Q = 0$, then the possible transformations among real bases are elements of $\mathrm{O(2)}\times\mathbb{Z}_2$. In particular, the sign of Z_5 changes when when the Higgs basis field $H_2 \to iH_2$. In this case, Z_5 is an O(2)-invariant but it is a pseudo-invariant with respect to $\mathbb{Z}_2$.

$$\begin{aligned}
&\text{Case I}: && h_1=h^0\,,\ h_2=-\varepsilon_6 H^0\,,\ h_3=\varepsilon_6 A^0\,,\ c_{12}=s_{\beta-\alpha}\ \text{and}\ s_{12}=-\varepsilon_6 c_{\beta-\alpha}\,,\\
&\text{Case IIa}: && h_1=h^0\,,\ h_2=\varepsilon_6 A^0\,,\ h_3=\varepsilon_6 H^0\,,\ c_{13}=s_{\beta-\alpha}\ \text{and}\ s_{13}=\varepsilon_6 c_{\beta-\alpha}\,,\\
&\text{Case IIb}: && h_1=\varepsilon_6 A^0\,,\ h_2=-h^0\,,\ h_3=\varepsilon_6 H^0\,,\ c_{13}=s_{\beta-\alpha}\ \text{and}\ s_{13}=\varepsilon_6 c_{\beta-\alpha}\,.
\end{aligned} \tag{3.38}$$

In the convention for the angular domain given by eq. (C.8), c_{12} and c_{13} are non-negative and therefore $s_{\beta-\alpha}\geq 0$. The appearance of the pseudo-invariant quantity ε_6 in eq. (3.38) implies that H^0, A^0 (and $H^\pm$) are pseudo-invariant fields, and $c_{\beta-\alpha}$ is a pseudo-invariant with respect to O(2) transformations.[9] In contrast, h^0 is an invariant field.

At this stage, we have not imposed any mass ordering of the three neutral scalar states. Since one can distinguish between the CP-odd and the CP-even neutral scalars, it is sufficient to require that $m_{h^0}\leq m_{H^0}$. (If one does not care about the mass ordering of A^0 relative to the CP-even states, then Cases IIa and IIb can be discarded without loss of generality.) We can compute the masses of the CP-even scalars and the angle $\beta-\alpha$ [59] in any of the three cases:

$$m^2_{h^0} = m^2_{A^0}\,c^2_{\beta-\alpha}+v^2\left[Z_1 s^2_{\beta-\alpha}+Z_5 c^2_{\beta-\alpha}+2s_{\beta-\alpha}c_{\beta-\alpha}Z_6\right]\,, \tag{3.39}$$

$$m^2_{H^0} = m^2_{A^0}\,s^2_{\beta-\alpha}+v^2\left[Z_1 c^2_{\beta-\alpha}+Z_5 s^2_{\beta-\alpha}-2s_{\beta-\alpha}c_{\beta-\alpha}Z_6\right]\,, \tag{3.40}$$

and

$$\tan[2(\beta-\alpha)]=\frac{2Z_6v^2}{m^2_{A^0}+(Z_5-Z_1)v^2}\,,\qquad \sin[2(\beta-\alpha)]=\frac{-2Z_6v^2}{m^2_{H^0}-m^2_{h^0}}\,. \tag{3.41}$$

Note that eqs. (3.39)–(3.41) are covariant with respect to O(2) transformations, since Z_6 and $c_{\beta-\alpha}$ are both pseudo-invariant quantities.

We end this section with a very brief outline of the tree-level MSSM Higgs sector. Since this model is CP-conserving, it is conventional to choose the phase conventions of the Higgs fields that yield a real basis. In the natural supersymmetric basis, the λ_i of eq. (A.1) are given by:

$$\lambda_1=\lambda_2=\tfrac14(g^2+g'^2)\,,\quad \lambda_3=\tfrac14(g^2-g'^2)\,,\quad \lambda_4=-\tfrac12 g^2\,,\quad \lambda_5=\lambda_6=\lambda_7=0\,, \tag{3.42}$$

where g and g' are the usual electroweak couplings [with $m_Z^2=\frac14(g^2+g'^2)v^2$]. From these results, one can compute the (pseudo)invariants:

$$\begin{aligned}
Z_1=Z_2 &= \tfrac14(g^2+g'^2)\cos^2 2\beta\,,\quad Z_3=Z_5+\tfrac14(g^2-g'^2)\,,\quad Z_4=Z_5-\tfrac12 g^2\,,\\
Z_5 &= \tfrac14(g^2+g'^2)\sin^2 2\beta\,,\quad Z_6=-Z_7=-\tfrac14(g^2+g'^2)\sin 2\beta\cos 2\beta\,.
\end{aligned} \tag{3.43}$$

The standard MSSM tree level Higgs sector formulae [57, 58] for the Higgs masses and $\beta-\alpha$ are easily reproduced using eq. (3.43) and the results of this section.

---

[8] The extra minus signs in the identification of $h_2=-\varepsilon_6 H^0$ in Case I and $h_2=-h^0$ in Case IIb arise due to the fact that the standard conventions of the CP-conserving 2HDM correspond to $\det R=-1$ (whereas $\det R=+1$ in Case IIa).

[9] Note that $s_{\beta-\alpha}$ is invariant with respect to O(2) transformations, which is consistent with our convention that $s_{\beta-\alpha}\geq 0$. The analogous results have been obtained in ref. [34] in a convention where $c_{\beta-\alpha}\geq 0$.

3.2 The 2HDM with $Z_6 = 0$

In this section we discuss the case where $Z_6 = 0$. For now, we will assume that all three neutral Higgs squared-masses are non-degenerate. Therefore, we require that $Z_5 \equiv |Z_5|e^{2i\theta_5} \neq 0$ in what follows,[10] and define the invariant angle $\phi_5 \equiv \theta_5 - \theta_{23}$. Once the sign conventions of the neutral Higgs fields are fixed, the invariant angles θ_{12}, θ_{13} and ϕ_5 are defined modulo π. We first note that eqs. (C.2), (C.3) and (C.11) are valid when $Z_6 = 0$. Thus, setting eq. (C.3) to zero implies that $\sin 2\theta_{13} = 0$,[11] which yields two possible solutions, $s_{13} = 0$ or $c_{13} = 0$. In the former case, eq. (C.2) yields $\mathrm{Im}(Z_5 e^{-2i\theta_{23}}) = 0$, *i.e.*, $\sin 2\phi_5 = 0$, and eq. (C.11) implies that $\sin 2\theta_{12} = 0$. In the latter case, we can use eq. (C.11) to write

$$\tan(2\theta_{12}) = -\tan(2\phi_5). \tag{3.44}$$

Thus, we can define three cases:

$$\begin{array}{lll} s_{13} = 0, \;\; \mathrm{Im}(Z_5\, e^{-2i\theta_{23}}) = c_{12} = 0 & & \text{Case } (i)\,, \\ s_{13} = 0, \;\; \mathrm{Im}(Z_5\, e^{-2i\theta_{23}}) = s_{12} = 0 & & \text{Case } (ii)\,, \\ c_{13} = 0, \;\; \mathrm{Im}[Z_5\, e^{2i(\theta_{12}-\theta_{23})}] = 0 & & \text{Case } (iii)\,. \end{array} \tag{3.45}$$

Let us start with (i). For $Z_6 = 0$, the mass matrix $\widetilde{\mathcal{M}}$ has the form [eq. (2.41)]

$$\widetilde{\mathcal{M}} \equiv R_{23}\mathcal{M}R_{23}^T = v^2 \begin{pmatrix} Z_1 & 0 & -\frac{1}{2}\mathrm{Im}(Z_5\, e^{-2i\theta_{23}}) \\ 0 & \mathrm{Re}(Z_5\, e^{-2i\theta_{23}}) + A^2/v^2 & 0 \\ 0 & -\frac{1}{2}\mathrm{Im}(Z_5\, e^{-2i\theta_{23}}) & A^2/v^2 \end{pmatrix}. \tag{3.46}$$

Applying the mass mixing matrix $\widetilde{R}\,\widetilde{\mathcal{M}}\,\widetilde{R}^T$ and setting $s_{13} = c_{12} = 0$ in $\widetilde{R}$ produces the diagonalized masses for case (i):

$$\begin{aligned} m_1^2 &= A^2 + v^2\ \mathrm{Re}(Z_5\, e^{-2i\theta_{23}}) = Y_2 + \tfrac{1}{2}[Z_3 + Z_4 + \mathrm{Re}(Z_5\, e^{-2i\theta_{23}})]\ v^2\,, \\ m_2^2 &= Z_1 v^2\,, \\ m_3^2 &= A^2 = Y_2 + \tfrac{1}{2}[Z_3 + Z_4 - \mathrm{Re}(Z_5\, e^{-2i\theta_{23}})]\ v^2\,, \end{aligned} \tag{3.47}$$

where we have used the convention $c_{13} = +1$ and $s_{12} = -1$ as required by eq. (C.8), and eliminated A^2 using eq. (2.42). Note that since $\mathrm{Im}(Z_5\, e^{-2i\theta_{23}}) = 0$, $\mathrm{Re}(Z_5\, e^{-2i\theta_{23}}) = \mathrm{sgn}(\mathrm{Z}_5)|\mathrm{Z}_5|^2$. The ambiguity in sign will be clarified in section 3.2.1.

For case (ii), one uses instead $s_{13} = s_{12} = 0$, with $c_{12} = c_{13} = +1$. Then taking $\widetilde{R}\,\widetilde{\mathcal{M}}\,\widetilde{R}^T$ yields

$$\begin{aligned} m_1^2 &= Z_1 v^2\,, \\ m_2^2 &= A^2 + v^2\ \mathrm{Re}(Z_5\, e^{-2i\theta_{23}}) = Y_2 + \tfrac{1}{2}[Z_3 + Z_4 + \mathrm{Re}(Z_5\, e^{-2i\theta_{23}})]\ v^2\,, \\ m_3^2 &= A^2 = Y_2 + \tfrac{1}{2}[Z_3 + Z_4 - \mathrm{Re}(Z_5\, e^{-2i\theta_{23}})]\ v^2\,. \end{aligned} \tag{3.48}$$

---

[10] If $Z_5 = Z_6 = 0$, then the neutral Higgs squared-mass matrix is diagonal in the Higgs basis, with two degenerate Higgs boson mass-eigenstates for $A^2 \neq Z_1 v^2$. If $A^2 = Z_1 v^2$ there are three degenerate Higgs boson mass-eigenstates.

[11] If $Z_6 = 0$ and $A^2 = Z_1 v^2$, then Eq. (C7) is automatically equal to zero. We will see in section 3.2.2 that the neutral Higgs masses are degenerate in this scenario.

The final (non-degenerate) scenario for $Z_6 = 0$ is case (iii), in which $q_{12} = ie^{i\theta_{12}}$, $q_{22v} = e^{i\theta_{12}}$, $q_{41} = i$, $q_{31} = -1$, and all other $q_{k\ell}$ vanish. Diagonalizing the mass matrix in eq. (3.46) yields

$$\begin{aligned} m_1^2 &= A^2 + v^2\mathrm{Re}(ie^{i\theta_{12}})\mathrm{Re}(ie^{i\theta_{12}}Z_5\, e^{-2i\theta_{23}})\,, \\ m_2^2 &= A^2 + v^2\mathrm{Re}(e^{i\theta_{12}})\mathrm{Re}(e^{i\theta_{12}}Z_5\, e^{-2i\theta_{23}})\,, \\ m_3^2 &= Z_1 v^2\,. \end{aligned} \tag{3.49}$$

Using $\mathrm{Im}(Z_5\, e^{2i(\theta_{12}-\theta_{23})}) = 0$ and $\mathrm{Re}[z_1, z_2] = \mathrm{Re}z_1\mathrm{Re}z_2 - \mathrm{Im}z_1\mathrm{Im}z_2$, the masses take the form

$$\begin{aligned} m_1^2 &= Y_2 + \tfrac{1}{2}[Z_3 + Z_4 - \mathrm{Re}(Z_5\, e^{2i(\theta_{12}-\theta_{23})})]\; v^2\,, \\ m_2^2 &= Y_2 + \tfrac{1}{2}[Z_3 + Z_4 + \mathrm{Re}(Z_5\, e^{2i(\theta_{12}-\theta_{23})})]\; v^2\,, \\ m_3^2 &= Z_1 v^2\,. \end{aligned} \tag{3.50}$$

Since $\mathrm{Im}(Z_5\, e^{2i(\theta_{12}-\theta_{23})})] = 0$, $\mathrm{Re}(Z_5\, e^{2i(\theta_{12}-\theta_{23})})] = \mathrm{sgn(Z_5)|Z_5|^2}$. Thus, up to a reordering of the three fields, all three cases exhibit the same masses,

$$\begin{aligned} m_1^2 &= Y_2 + \tfrac{1}{2}[Z_3 + Z_4 - \mathrm{sgn(Z_5)|Z_5|]\; v^2}\,, \\ m_2^2 &= Y_2 + \tfrac{1}{2}[Z_3 + Z_4 + \mathrm{sgn(Z_5)|Z_5|]\; v^2}\,, \\ m_3^2 &= Z_1 v^2\,. \end{aligned} \tag{3.51}$$

3.2.1 CP Conservation with $Z_6 = 0$

Since the mass matrix for $Z_6 = 0$ is broken up into 2×2 and 1×1 blocks [see eq. (3.46)], one might expect that the scalar sector is automatically CP-conserving. Indeed, the field whose mass is given by $Z_1 v^2$ in the three cases (i)–(iii) is CP even, as one can deduce from the couplings $G^0G^0h_k$. However, the coupling $H^+H^-h_k$ contains the expression $\mathrm{Re}(q_{k2}Z_7\, e^{-i\theta_{23}})$ and the $\bar{Q}Qh_k$ interactions [eq. (2.83)] contain $q_{k2}^* e^{i\theta_{23}}\rho^Q$ $(Q = U, D)$. Thus, for CP to be conserved, one requires

$$\mathrm{Im}(Z_5^* Z_7^2) = 0, \qquad \mathrm{Im}[Z_5(\rho^Q)^2] = 0, \text{ and } \mathrm{Im}[Z_7\rho^Q] = 0. \tag{3.52}$$

One observes from eqs. (3.47), (3.48) and (3.50) that in case (iii), the angle $\theta_{23} - \theta_{12}$ plays the same role as θ_{23} in cases (i) and (ii). Thus, it is convenient to define

$$\bar{\theta}_{23} \equiv \begin{cases} \theta_{23} & \text{cases } (i),(ii)\,; \\ \theta_{23} - \theta_{12} & \text{case } (iii)\,. \end{cases} \tag{3.53}$$

Since $\mathrm{Im}(Z_5\, e^{-2i\bar{\theta}_{23}}) = 0$ [see eq. (3.45)] and $\mathrm{Im}[Z_7\rho^Q] = 0$, one can write

$$\begin{aligned} \mathrm{Im}[Z_5^* Z_7^2] &= \mathrm{Re}(Z_5\, e^{-2i\bar{\theta}_{23}})\mathrm{Im}(Z_7^2\, e^{-2i\bar{\theta}_{23}}) \\ &= \mathrm{Re}(Z_5\, e^{-2i\bar{\theta}_{23}})2\mathrm{Im}(Z_7\, e^{-i\bar{\theta}_{23}})\mathrm{Re}(Z_7\, e^{-i\bar{\theta}_{23}}) = 0\,, \end{aligned} \tag{3.54}$$

$$\begin{aligned} \mathrm{Im}[Z_7\, e^{-i\bar{\theta}_{23}} e^{i\bar{\theta}_{23}}\rho^Q] &= \mathrm{Re}(Z_7\, e^{-i\bar{\theta}_{23}})\mathrm{Im}(\rho^Q\, e^{i\bar{\theta}_{23}}) \\ &\quad + \mathrm{Im}(Z_7\, e^{-i\bar{\theta}_{23}})\mathrm{Re}(\rho^Q\, e^{i\bar{\theta}_{23}}) = 0\,. \end{aligned} \tag{3.55}$$

Eqs. (3.54) and (3.55) have two solutions,

$$\begin{aligned}
(a) \qquad \mathrm{Im}(Z_7 e^{-i\bar{\theta}_{23}}) \quad &= \quad \mathrm{Im}[\rho^Q e^{i\bar{\theta}_{23}}] \;=\; 0\,, && (3.56)\\
(b) \qquad \mathrm{Re}(Z_7 e^{-i\bar{\theta}_{23}}) \quad &= \quad \mathrm{Re}[\rho^Q e^{i\bar{\theta}_{23}}] \;=\; 0\,. && (3.57)
\end{aligned}$$

Note that eq. (3.10) correctly defines the CP transformation for $Z_6 = 0$ provided that one replace θ_{23} by $\bar{\theta}_{23}$. The transformation with the positive sign in eq. (3.10) corresponds to solution (a) [eq. (3.56)] and the one with the negative sign to solution (b) [eq. (3.57)].

If neither eq. (3.56) nor eq. (3.57) holds, then CP is violated, and the remaining two neutral scalar fields will have indefinite CP quantum numbers. If the conditions of eq. (3.52) hold, all neutral scalars will be CP eigenstates, with CP quantum numbers displayed in Tables 3.4–3.5. Comparing the results of the tables with eqs. (3.47), (3.48) and (3.50), one finds that in all three cases,

$$\begin{aligned}
m_{A^0}^2 \;&=\; Y_2 + \tfrac{1}{2}[Z_3 + Z_4 - \mathrm{Re}(Z_5\, e^{-2i\bar{\theta}_{23}})]\; v^2\,, \text{ for } \mathrm{Im}(\mathrm{Z_7\, e^{-i\bar{\theta}_{23}}}) = \mathrm{Im}(\rho^{\mathrm{Q}}\, \mathrm{e}^{\mathrm{i}\bar{\theta}_{23}}) = 0\,,\\
m_{A^0}^2 \;&=\; Y_2 + \tfrac{1}{2}[Z_3 + Z_4 + \mathrm{Re}(Z_5\, e^{-2i\bar{\theta}_{23}})]\; v^2\,, \text{ for } \mathrm{Re}(\mathrm{Z_7\, e^{-i\bar{\theta}_{23}}}) = \mathrm{Re}(\rho^{\mathrm{Q}}\, \mathrm{e}^{\mathrm{i}\bar{\theta}_{23}}) = 0\,.
\end{aligned} \tag{3.58}$$

$$\begin{aligned}
m_{H^0}^2 \;&=\; Y_2 + \tfrac{1}{2}[Z_3 + Z_4 + \mathrm{Re}(Z_5\, e^{-2i\bar{\theta}_{23}})]\; v^2\,, \text{ for } \mathrm{Im}(\mathrm{Z_7\, e^{-i\bar{\theta}_{23}}}) = \mathrm{Im}(\rho^{\mathrm{Q}}\, \mathrm{e}^{\mathrm{i}\bar{\theta}_{23}}) = 0\,,\\
m_{H^0}^2 \;&=\; Y_2 + \tfrac{1}{2}[Z_3 + Z_4 - \mathrm{Re}(Z_5\, e^{-2i\bar{\theta}_{23}})]\; v^2\,, \text{ for } \mathrm{Re}(\mathrm{Z_7\, e^{-i\bar{\theta}_{23}}}) = \mathrm{Re}(\rho^{\mathrm{Q}}\, \mathrm{e}^{\mathrm{i}\bar{\theta}_{23}}) = 0\,.
\end{aligned} \tag{3.59}$$

One can condense these results by defining the symbol ε_{57}:

$$\mathrm{Re}(Z_5^* Z_7^2) = \varepsilon_{57} |Z_5||Z_7|^2\,, \qquad \varepsilon_{57} = \pm 1\,. \tag{3.60}$$

The quantity ε_{57} is independent of basis. Note that

$$\begin{aligned}
\mathrm{Re}(Z_5^* e^{2i\theta_{23}}) \quad &= \quad \frac{\mathrm{Re}(Z_5^* |Z_7|^2 e^{2i\theta_{23}})}{|Z_7|^2}\,,\\
&= \quad \frac{\mathrm{Re}(Z_5^* Z_7^2)\mathrm{Re}[(Z_7^*)^2 e^{2i\theta_{23}}]}{|Z_7|^2}\,.
\end{aligned} \tag{3.61}$$

Then applying the definition of ϵ_{57} in eq. (3.60),

$$\begin{aligned}
\mathrm{Re}(Z_7\, e^{-i\theta_{23}}) = 0 \quad &\Rightarrow \quad \varepsilon_{57}|Z_5| = -\mathrm{Re}(Z_5\, e^{-2i\theta_{23}})\,,\\
\mathrm{Im}(Z_7\, e^{-i\theta_{23}}) = 0 \quad &\Rightarrow \quad \varepsilon_{57}|Z_5| = +\mathrm{Re}(Z_5\, e^{-2i\theta_{23}})\,.
\end{aligned} \tag{3.62}$$

Thus, equations (3.58) can be expressed as a single equation, so the mass of the CP-odd field is given by

$$m_{A^0}^2 = Y_2 + \tfrac{1}{2}[Z_3 + Z_4 - \varepsilon_{57}|Z_5|]\; v^2\,, \tag{3.63}$$

and the second CP-even field has mass

$$m_{H^0}^2 = Y_2 + \tfrac{1}{2}[Z_3 + Z_4 + \varepsilon_{57}|Z_5|]\; v^2\,. \tag{3.64}$$

If $Z_7 = 0$ but $\rho^Q \neq 0$, one can derive the analog of ϵ_{57} as follows: From the conditions of CP symmetry in eq. (3.52),

$$\mathrm{Tr}_f \mathrm{Im}[Z_5(\rho^Q)^2] = 0\,, \qquad \Rightarrow \qquad \mathrm{Im}\{Z_5 \mathrm{Tr}_f[(\rho^Q)^2]\} = 0\,. \tag{3.65}$$

Any 2×2 matrix A satisfies its characteristic equation, $A^2 - A\mathrm{Tr}A + \det A = 0$. Taking the trace of this equation yields the identity $\mathrm{Tr}(A^2) - (\mathrm{Tr}A)^2 + 2\det A = 0$ Thus, the condition in eq. (3.65) is equivalent to

$$\mathrm{Im}[Z_5(\mathrm{Tr}_f\rho^Q)^2 - 2Z_5\det(\rho^Q)] = 0\,, \qquad \Rightarrow \qquad \mathrm{Im}\{Z_5 \mathrm{Tr}_f[(\rho^Q)^2]\} = 0\,. \tag{3.66}$$

Note that $\rho^Q e^{i\theta_{23}}$ is purely real or imaginary [see eqs. (3.56) and (3.57)], so

$$\mathrm{Im}[Z_5\det(\rho^Q)] = \mathrm{Re}(Z_5\, e^{-2i\theta_{23}})\mathrm{Im}[\det(\rho^Q e^{-i\theta_{23}})] = 0\,. \tag{3.67}$$

Then eq. (3.66) becomes

$$\mathrm{Im}[Z_5(\mathrm{Tr}_f\rho^Q)^2] = 0\,. \tag{3.68}$$

Now we can define

$$\mathrm{Re}[Z_5(\mathrm{Tr}_f\rho^Q)^2] \equiv \varepsilon_{5Q}|Z_5||(\mathrm{Tr}\rho^Q)^2|\,, \qquad \varepsilon_{5Q} = \pm 1\,. \tag{3.69}$$

Using a similar calculation as in the case with $Z_7 \neq 0$, one finds

$$\begin{aligned} \mathrm{Re}(\rho^Q\, e^{-i\theta_{23}}) = 0 \quad &\Rightarrow \quad \varepsilon_{5Q}|Z_5| = -\mathrm{Re}(Z_5\, e^{-2i\theta_{23}})\,, \\ \mathrm{Im}(\rho^Q\, e^{-i\theta_{23}}) = 0 \quad &\Rightarrow \quad \varepsilon_{5Q}|Z_5| = +\mathrm{Re}(Z_5\, e^{-2i\theta_{23}})\,. \end{aligned} \tag{3.70}$$

Thus, for $Z_6 = Z_7 = 0$, one replaces ε_{57} by ε_{5Q} in eqs. (3.63) and (3.64).

To summarize, we find that for $Z_6 = 0$ there is one neutral Higgs field that is always CP-even, with mass squared equal to $Z_1 v^2$. Unless $Z_5^* Z_7^2$, $Z_7\rho^Q$, and $Z_5(\rho^Q)^2$ are all real-valued, the remaining neutral fields are mixtures of CP eigenstates, even though the mass matrix has a block diagonal form. In the CP conserving case, the two solutions represented by eq. (3.56) and eq. (3.57) correspond to different possibilities for the CP quantum numbers of those 2 remaining fields. An overview of the three cases and the associated CP quantum numbers for the CP-conserving case are given in Tables 3.4–3.6. The preceding results are valid for the CP conserving case with $Z_6 = 0$ as long either ρ^Q or Z_7 is non-vanishing. If $Z_6 = Z_7 = \rho^Q = 0$, the model has some extra features which will be described in section 3.2.3.

Table 3.4: Values of $q_{k\ell}$ in the CP-conserving limit with $Z_6 = 0$. Case (i): $s_{13} = c_{12} = \mathrm{Im}(Z_5\, e^{-2i\theta_{23}}) = 0$. The CP quantum numbers are shown for $\mathrm{Im}(Z_7\, e^{-i\theta_{23}}) = \mathrm{Im}(\rho^Q e^{i\bar{\theta}_{23}}) = 0$ (upper sign), and $\mathrm{Re}(Z_7\, e^{-i\theta_{23}}) = \mathrm{Re}(\rho^Q e^{i\bar{\theta}_{23}}) = 0$ (lower sign).

k	q_{k1}	q_{k2}	CP
1	0	1	± 1
2	-1	0	$+1$
3	0	i	∓ 1
4	i	0	-1

Table 3.5: Values of $q_{k\ell}$ in the CP-conserving limit with $Z_6 = 0$. Case (*ii*): $s_{13} = s_{12} = \mathrm{Im}(Z_5\, e^{-2i\theta_{23}}) = 0$. The CP quantum numbers are shown for $\mathrm{Im}(Z_7\, e^{-i\theta_{23}}) = \mathrm{Im}(\rho^Q e^{i\bar{\theta}_{23}}) = 0$ (upper sign), and $\mathrm{Re}(Z_7\, e^{-i\theta_{23}}) = \mathrm{Re}(\rho^Q e^{i\bar{\theta}_{23}}) = 0$ (lower sign).

k	q_{k1}	q_{k2}	CP
1	1	0	$+1$
2	0	1	± 1
3	0	i	∓ 1
4	i	0	-1

Table 3.6: Values of $q_{k\ell}$ in the CP-conserving limit with $Z_6 = 0$. Case (*iii*): $c_{13} = \mathrm{Im}(Z_5\, e^{2i(\theta_{12}-\theta_{23})})$. The CP quantum numbers are shown for $\mathrm{Im}(Z_7\, e^{-i\theta_{23}}) = \mathrm{Im}(\rho^Q e^{i\bar{\theta}_{23}}) = 0$ (upper sign), and $\mathrm{Re}(Z_7\, e^{-i\theta_{23}}) = \mathrm{Re}(\rho^Q e^{i\bar{\theta}_{23}}) = 0$ (lower sign).

k	q_{k1}	q_{k2}	CP
1	0	$ie^{i\theta_{12}}$	∓ 1
2	0	$e^{i\theta_{12}}$	± 1
3	-1	0	$+1$
4	i	0	-1

3.2.2 $Z_6 = 0$ with Degenerate Neutral Scalars

In the previous discussion we assumed that none of the of neutral scalar masses were degenerate. If we relax that requirement, three additional ways of satisfying eqs. (C.2), (C.3) and (C.11) appear:

$$\begin{array}{lll}
\mathrm{Im}(Z_5\, e^{-2i\theta_{23}}) = s_{13} = 0, & Z_1 v^2 = Y_2 + \tfrac{1}{2}[Z_3 + Z_4 + \mathrm{Re}(Z_5\, e^{-2i\theta_{23}})]v^2\,, & \text{Case } (iv)\,, \\
\mathrm{Im}(Z_5\, e^{-2i\theta_{23}}) = c_{12} = 0, & Z_1 v^2 = Y_2 + \tfrac{1}{2}[Z_3 + Z_4 - \mathrm{Re}(Z_5\, e^{-2i\theta_{23}})]v^2\,, & \text{Case } (v)\,, \\
\mathrm{Im}(Z_5\, e^{-2i\theta_{23}}) = s_{12} = 0, & Z_1 v^2 = Y_2 + \tfrac{1}{2}[Z_3 + Z_4 - \mathrm{Re}(Z_5\, e^{-2i\theta_{23}})]v^2\,. & \text{Case } (vi)\,.
\end{array}$$

These cases are presented in Tables 3.7–3.9.

For case (*iv*), diagonalizing the mass matrix yields

$$\begin{aligned}
m_1^2 &= s_{12}^2 A^2 + c_{12}^2 Z_1 v^2 + v^2 s_{12}^2 \mathrm{Re}(Z_5\, e^{-2i\theta_{23}}) = Z_1 v^2\,, \\
m_2^2 &= c_{12}^2 A^2 + s_{12}^2 Z_1 v^2 + v^2 c_{12}^2 \mathrm{Re}(Z_5\, e^{-2i\theta_{23}}) = Z_1 v^2\,, \\
m_3^2 &= A^2 = Y_2 + \tfrac{1}{2} v^2 [Z_3 + Z_4 - \mathrm{Re}(Z_5\, e^{-2i\theta_{23}})]\,, \qquad (3.71)
\end{aligned}$$

where A^2 has been replaced by $Z_1 v^2 - v^2 \mathrm{Re}(Z_5\, e^{-2i\theta_{23}})$. The analogous calculation for case (*v*) gives

$$\begin{aligned}
m_1^2 &= A^2 + v^2 \mathrm{Re}(Z_5\, e^{-2i\theta_{23}}) = Y_2 + \tfrac{1}{2} v^2 [Z_3 + Z_4 + \mathrm{Re}(Z_5\, e^{-2i\theta_{23}})]\,, \\
m_2^2 &= s_{13}^2 A^2 + v^2 c_{13}^2 Z_1 = Z_1 v^2\,, \\
m_3^2 &= c_{13}^2 A^2 + v^2 s_{13}^2 Z_1 = Z_1 v^2\,. \qquad (3.72)
\end{aligned}$$

Similarly, for case (vi) one finds

$$\begin{aligned} m_1^2 &= Z_1 v^2\,, \\ m_2^2 &= v^2[Z_1 + \mathrm{Re}(Z_5\, e^{-2i\theta_{23}})] = Y_2 + \tfrac{1}{2}v^2[Z_3 + Z_4 + \mathrm{Re}(Z_5\, e^{-2i\theta_{23}})]\,, \\ m_3^2 &= Z_1 v^2\,. \end{aligned} \tag{3.73}$$

One observes that the degenerate fields have mass squared equal to $Z_1 v^2$ in all cases. If $\mathrm{Im}[Z_5^* Z_7^2] = 0$, $\mathrm{Im}[Z_5(\rho^Q)^2] = 0$, and $\mathrm{Im}[Z_7\rho^Q] = 0$, the scalar sector will be CP-conserving. In case (iv), the interaction $\mathrm{Re}(q_{k2} Z_7 e^{-i\theta_{23}}) h_k H^+ H^-$ indicates that the non-degenerate field, h_3, is CP-odd if $\mathrm{Im}(Z_7\, e^{-i\theta_{23}}) = 0$ and CP-even if $\mathrm{Re}(Z_7\, e^{-i\theta_{23}}) = 0$. One also deduces that the combination $s_{12}h_1 + c_{12}h_2$ is CP-even if $\mathrm{Im}(Z_7\, e^{-i\theta_{23}}) = 0$ and CP-odd if $\mathrm{Re}(Z_7\, e^{-i\theta_{23}}) = 0$. [One obtains analogous results using the $\bar{Q}Qh_k$ interaction.] The orthogonal combination is always CP-even. These results are summarized in Table 3.10. A similar analysis for cases (v) and (vi) gives the results shown in Tables 3.11 and 3.12, respectively. Using Tables 3.10–3.12 and the equations for the masses in eqs. (3.71), (3.72) and (3.73), one can write expressions for the mass of the CP-odd field that apply in all three cases, namely

Table 3.7: Values of $q_{k\ell}$ with $Z_6 = 0$. Case (iv): $s_{13} = \mathrm{Im}(Z_5\, e^{-2i\theta_{23}}) = Z_1 - A^2/v^2 - \mathrm{Re}(Z_5\, e^{-2i\theta_{23}}) = 0$.

k	q_{k1}	q_{k2}
1	c_{12}	$-s_{12}$
2	s_{12}	c_{12}
3	0	i
4	i	0

Table 3.8: Values of $q_{k\ell}$ with $Z_6 = 0$. Case (v): $c_{12} = \mathrm{Im}(Z_5\, e^{-2i\theta_{23}}) = Z_1 - A^2/v^2 = 0$.

k	q_{k1}	q_{k2}
1	0	1
2	$-c_{13}$	is_{13}
3	s_{13}	ic_{13}
4	i	0

Table 3.9: Values of $q_{k\ell}$ with $Z_6 = 0$. Case (vi): $s_{12} = \mathrm{Im}(Z_5\, e^{-2i\theta_{23}}) = Z_1 - A^2/v^2 = 0$.

k	q_{k1}	q_{k2}
1	c_{13}	$-is_{13}$
2	0	1
3	s_{13}	ic_{13}
4	i	0

$$m_{A^0}^2 = Y_2 + \tfrac{1}{2}[Z_3 + Z_4 - \varepsilon_{57}|Z_5|]\; v^2\,, \tag{3.74}$$

Table 3.10: The CP quantum numbers of the neutral scalars with $Z_6 = 0$. Case (*iv*): $s_{13} = \mathrm{Im}(Z_5\, e^{-2i\theta_{23}}) = Z_1 - A^2/v^2 - \mathrm{Re}(Z_5\, e^{-2i\theta_{23}}) = 0$.

CP Eigenstates	$\mathrm{Re}(Z_7\, e^{-i\theta_{23}}) =$ $\mathrm{Re}(\rho^Q e^{i\theta_{23}}) = 0$	$\mathrm{Im}(Z_7\, e^{-i\theta_{23}}) =$ $\mathrm{Im}(\rho^Q e^{i\theta_{23}}) = 0$
$c_{12}h_1 + s_{12}h_2$	$+1$	$+1$
$-s_{12}h_1 + c_{12}h_2$	-1	$+1$
h_3	$+1$	-1

Table 3.11: The CP quantum numbers of the neutral scalars with $Z_6 = 0$. Case (*v*): $c_{12} = \mathrm{Im}(Z_5\, e^{-2i\theta_{23}}) = Z_1 - A^2/v^2 = 0$.

CP Eigenstates	$\mathrm{Re}(Z_7\, e^{-i\theta_{23}}) =$ $\mathrm{Re}(\rho^Q e^{i\theta_{23}}) = 0$	$\mathrm{Im}(Z_7\, e^{-i\theta_{23}}) =$ $\mathrm{Im}(\rho^Q e^{i\theta_{23}}) = 0$
h_1	-1	$+1$
$c_{13}h_2 - s_{13}h_3$	$+1$	$+1$
$s_{13}h_1 + c_{13}h_3$	$+1$	-1

Table 3.12: The CP quantum numbers of the neutral scalars with $Z_6 = 0$. Case (*vi*): $s_{12} = \mathrm{Im}(Z_5\, e^{-2i\theta_{23}}) = Z_1 - A^2/v^2 = 0$.

CP Eigenstates	$\mathrm{Re}(Z_7\, e^{-i\theta_{23}}) =$ $\mathrm{Re}(\rho^Q e^{i\theta_{23}}) = 0$	$\mathrm{Im}(Z_7\, e^{-i\theta_{23}}) =$ $\mathrm{Im}(\rho^Q e^{i\theta_{23}}) = 0$
$c_{13}h_1 + s_{13}h_3$	$+1$	$+1$
h_2	-1	$+1$
$-s_{13}h_1 + c_{13}h_3$	$+1$	-1

and one of the CP-even field has mass

$$m_{H^0}^2 = Y_2 + \tfrac{1}{2}[Z_3 + Z_4 + \varepsilon_{57}|Z_5|]\ v^2\,. \tag{3.75}$$

as in eqs. (3.63) and (3.64). The mass of the other CP-even field is always one of the degenerate fields can be expressed in all three cases as follows:

$$m_{h^0}^2 = Z_1 v^2\,. \tag{3.76}$$

(We have not imposed $m_{h^0}^2 < m_{H^0}^2$). Note that eqs. (3.74), (3.75) and (3.76) hold for all six cases (*i*)–(*vi*). The neutral scalar field with mass $m_{h^0}^2 = Z_1 v^2$ has exact Standard Model couplings. As in the previous section, if $Z_7 = 0$, ε_{57} can be replaced by ε_{5Q}.

3.2.3 Special Case: The CP-Conserving Limit when $Z_6 = 0$ and $Z_7 = 0$

In this section, we consider a 2HDM with $Y_3 = Z_6 = Z_7 = 0$.[12] For the moment, let us also assume that $\rho^Q = 0$.

---

[12] Technically, since the potential minimum conditions require $Y_3 = -\frac{1}{2}Z_6 v^2$, explicitly setting $Y_3 = 0$ is redundant.

This model is automatically CP conserving (since one can choose Z_5 to be real and positive without loss of generality). Normally, in a CP-invariant 2HDM, starting from a real Higgs basis one can get to any possible generic real basis with an O(2) transformation. However, in the model under study here, there exists a particular U(2) transformation that is not an O(2) transformation, which has the effect of changing the sign of Z_5. This corresponds to redefining the second Higgs field by multiplication by i, ie,

$$H_2 \to iH_2\,. \tag{3.77}$$

The relevant U(2) matrix is $diag(1, i)$. In Appendix A of [60], the possible definitions of time reversal invariance are discussed. In particular, it was argued that the definition of T is unique if the possible transformations from the real Higgs basis to any real generic basis is O(2). It was shown that T was not unique if the latter transformation group was $O(2) \otimes \mathbb{D}$ where $\mathbb{D}$ is a nontrivial discrete group. Applying this to the model where $Y_3 = Z_6 = Z_7 = 0$, we identify the relevant group as $O(2) \otimes \mathbb{Z}_2$, where $\mathbb{Z}_2$ is the discrete group consisting of the identity and the transformation that changes the sign of Z_5. Thus, we conclude that for the $Y_3 = Z_6 = Z_7 = 0$ model, there are two inequivalent definitions of T, or equivalently two definitions of CP (since the model is CPT invariant). These two symmetries correspond to the two possible transformations defined in eq. (3.10), where the plus sign applies to $\mathrm{Im}(Z_6 e^{-i\bar{\theta}_{23}}) = \mathrm{Im}(Z_7 e^{-i\bar{\theta}_{23}}) = \mathrm{Im}[\rho^Q e^{i\bar{\theta}_{23}}] = 0$ and the negative sign to $\mathrm{Re}(Z_6 e^{-i\bar{\theta}_{23}}) = \mathrm{Re}(Z_7 e^{-i\bar{\theta}_{23}}) = \mathrm{Re}[\rho^Q e^{i\bar{\theta}_{23}}] = 0$.

In this model, the Higgs/gauge boson interactions are insufficient to identify the CP-odd field. We know from the results of the previous section that one of the neutral scalars has couplings exactly identical to the Standard Model Higgs, with $m^2_{h^0} = Z_1 v^2$. We also know that the two remaining neutral fields have opposite CP. However, since $\rho^Q = 0$ the interactions cannot distinguish which one is CP-even and which one is CP-odd; the two inequivalent CP symmetries are both conserved.

Now let us suppose $\rho^Q \neq 0$. In section 3.2.1, we showed how to identify the CP-odd field based on the value of ε_{5Q}. Thus, a non-zero value of ρ^Q has the effect of picking out the definition of CP that is respected by the Yukawa interactions. In fact, a non-zero value of any one of Z_6, Z_7 or ρ^Q identifies the respected CP symmetry; if any two are non-zero, CP violation arises unless the relative phases obey eq. (3.28).

3.3 The Custodial Limit of the 2HDM

The subject of custodial symmetry in the 2HDM doublet model has been addressed by Pomarol and Vega [108], in the context of two cases which they label "case I" and "case II," *not to be confused with the two cases of CP-conservation* defined in equations (3.3) and (3.4), to which they have no correlation. We reproduce some of their work here in order to clarify the significance of their two cases. Specifically, we show in this section that the only difference between their "case I" and "case II" lies in the conditions they impose on the vevs of the scalar fields, which is not a physically measurable distinction.

First we will ignore the coupling of the Higgs doublets to fermions and just consider the scalar potential. To replicate "case I" of [108], we construct two 2×2 matrices whose columns are made up of the Higgs doublet fields in the generic basis:

$$M_1 \equiv (\tilde{\Phi}_1 \, , \Phi_1), \qquad M_2 \equiv (\tilde{\Phi}_2 \, , \Phi_2), \tag{3.78}$$

where $\tilde{\Phi} \equiv i\sigma_2 \Phi^*$. These matrices transform as

$$M_i \to L \ M_i \ R^\dagger \tag{3.79}$$

under global $SU(2)_L \times SU(2)_R$ transformations. In order that a custodial $SU(2)_V$ symmetry be preserved after electroweak symmetry breaking, $< M_i >$ must be proportional to the identity matrix:

$$< M_i >= \frac{1}{\sqrt{2}} \begin{pmatrix} v_i^* & 0 \\ 0 & v_i \end{pmatrix} \propto 1 \quad \Rightarrow \quad v_i^* = v_i, . \tag{3.80}$$

Since $L = R$,

$$< M_i > \to L < M_i > R^\dagger = \frac{v_i}{\sqrt{2}} L R^\dagger =< M_i > \ . \tag{3.81}$$

Note that

$$\mathrm{Tr}[M_i^\dagger M_j] \ = \ \mathrm{Tr}\left[\begin{pmatrix} \Phi_i^0 & -\Phi_i^+ \\ \Phi_i^- & \Phi_i^{0*} \end{pmatrix} \begin{pmatrix} \Phi_j^{0*} & \Phi_j^+ \\ -\Phi_j^- & \Phi_j^0 \end{pmatrix} \right] = \Phi_i^\dagger \Phi_j + \Phi_j^\dagger \Phi_i. \tag{3.82}$$

Thus,

$$\begin{aligned} \mathrm{Tr}[M_1^\dagger M_1] &= 2\Phi_1^\dagger \Phi_1 \\ \mathrm{Tr}[M_2^\dagger M_2] &= 2\Phi_2^\dagger \Phi_2 \\ \mathrm{Tr}[M_1^\dagger M_2] &= \Phi_2^\dagger \Phi_1 + h.c. \end{aligned} \tag{3.83}$$

Using the expressions in eq. (3.83) to construct a $SU(2)_L \times SU(2)_R$ symmetric scalar potential, we reproduce the following result of [108]:

$$\begin{aligned} \mathcal{V} \ = \ & \tfrac{1}{2} m_{11}^2 \mathrm{Tr}[M_1^\dagger M_1] + \tfrac{1}{2} m_{22}^2 \mathrm{Tr}[M_2^\dagger M_2] - m_{12}^2 \mathrm{Tr}[M_1^\dagger M_2] + \frac{1}{4}\lambda_1 \left(\mathrm{Tr}[M_1^\dagger M_1] \right)^2 \\ & + \frac{1}{4}\lambda_2 \left(\mathrm{Tr}[M_2^\dagger M_2] \right)^2 + \tfrac{1}{4}\lambda_3 \mathrm{Tr}[M_1^\dagger M_1] \mathrm{Tr}[M_2^\dagger M_2] + \tfrac{1}{2}\lambda \left(\mathrm{Tr}[M_1^\dagger M_2] \right)^2 \\ & + \tfrac{1}{4} \left(\lambda_6 \mathrm{Tr}[M_1^\dagger M_1] + \lambda_7 \mathrm{Tr}[M_2^\dagger M_2] \right) \mathrm{Tr}[M_1^\dagger M_2]. \end{aligned} \tag{3.84}$$

Comparing eq. (3.84) to the most general form of the scalar potential in eq. (A.1), we find that custodial symmetry imposes the following restrictions on the coefficients of the scalar potential:

$$m_{12}^* = m_{12} \, , \ \lambda_{5,6,7}^* = \lambda_{5,6,7} \, , \ \lambda = \lambda_4 = \lambda_5 \, . \tag{3.85}$$

Let us now convert from the generic basis to the Higgs basis. Plugging $\Phi_a = H_1\widehat{v}_a + H_2\widehat{v}_{\bar{b}}^*\epsilon_{ba}$ into eq. (3.83) yields

$$\begin{aligned}
\mathrm{Tr}[M_1^\dagger M_1] &= 2\left(|\widehat{v}_1|^2 H_1^\dagger H_1 + |\widehat{v}_2|^2 H_2^\dagger H_2 - \widehat{v}_1\widehat{v}_2 H_2^\dagger H_1 - \widehat{v}_1^*\widehat{v}_2^* H_1^\dagger H_2\right) \\
\mathrm{Tr}[M_2^\dagger M_2] &= 2\left(|\widehat{v}_2|^2 H_1^\dagger H_1 + |\widehat{v}_1|^2 H_2^\dagger H_2 + \widehat{v}_1\widehat{v}_2 H_2^\dagger H_1 + \widehat{v}_1^*\widehat{v}_2^* H_1^\dagger H_2\right) \\
\mathrm{Tr}[M_1^\dagger M_2] &= \widehat{v}_2^*\widehat{v}_1 H_1^\dagger H_1 - \widehat{v}_1\widehat{v}_2^* H_2^\dagger H_2 + \left(\widehat{v}_1^2 H_2^\dagger H_1 - \widehat{v}_2^{*^2} H_1^\dagger H_2 + h.c.\right) . \qquad (3.86)
\end{aligned}$$

Subsituting equations (3.86) into eq. (3.84) and grouping like terms yields the following:

$$\begin{aligned}
\mathcal{V} &= Y_1 H_1^\dagger H_1 + Y_2 H_2^\dagger H_2 + [Y_3 H_1^\dagger H_2 + \mathrm{h.c.}] \\
&\quad + \tfrac{1}{2}Z_1(H_1^\dagger H_1)^2 + \tfrac{1}{2}Z_2(H_2^\dagger H_2)^2 + Z_3(H_1^\dagger H_1)(H_2^\dagger H_2) + Z_4(H_1^\dagger H_2)(H_2^\dagger H_1) \\
&\quad + \left\{\tfrac{1}{2}Z_5(H_1^\dagger H_2)^2 + \left[Z_6(H_1^\dagger H_1) + Z_7(H_2^\dagger H_2)\right] H_1^\dagger H_2 + \mathrm{h.c.}\right\} , \qquad (3.87)
\end{aligned}$$

where

$$\begin{aligned}
Y_1 &= m_1^2|\widehat{v}_1|^2 + m_2^2|\widehat{v}_2|^2 - m_{12}^2\left(\widehat{v}_2^*\widehat{v}_1 + c.c.\right) , \\
Y_2 &= m_1^2|\widehat{v}_2|^2 + m_2^2|\widehat{v}_1|^2 + m_{12}^2\left(\widehat{v}_2^*\widehat{v}_1 + c.c.\right) , \\
Y_3 &= m_{12}^2(\widehat{v}_2^{*2} - \widehat{v}_1^{*2}) + \widehat{v}_2^*\widehat{v}_1^*(m_2^2 - m_1^2) , \\
Z_1 &= 2\left(\lambda_1|\widehat{v}_1|^4 + \lambda_2|\widehat{v}_2|^4 + \lambda_3|\widehat{v}_1|^2|\widehat{v}_2|^2\right) + \lambda\left(\widehat{v}_2^*\widehat{v}_1 + c.c.\right)^2 \\
&\quad +(\lambda_6|\widehat{v}_1|^2 + \lambda_7|\widehat{v}_2|^2)\left(\widehat{v}_2^*\widehat{v}_1 + c.c.\right) , \\
Z_2 &= 2\left(\lambda_1|\widehat{v}_2|^4 + \lambda_2|\widehat{v}_1|^4 + \lambda_3|\widehat{v}_1|^2|\widehat{v}_2|^2\right) + \lambda\left(\widehat{v}_2^*\widehat{v}_1 + c.c.\right)^2 \\
&\quad -(\lambda_6|\widehat{v}_2|^2 + \lambda_7|\widehat{v}_1|^2)\left(\widehat{v}_2^*\widehat{v}_1 + c.c.\right) , \\
Z_3 &= 2(\lambda_1 + \lambda_2)|\widehat{v}_1|^2|\widehat{v}_2|^2 + \lambda_3(|\widehat{v}_1|^4 + |\widehat{v}_2|^4) - \lambda\left(\widehat{v}_2^*\widehat{v}_1 + c.c.\right)^2 \\
&\quad +\tfrac{1}{2}(\lambda_6 - \lambda_7)\left(\widehat{v}_2^*\widehat{v}_1 + c.c.\right)\left(|\widehat{v}_2|^2 - |\widehat{v}_1|^2\right) , \\
Z_4 &= 2(\lambda_1 + \lambda_2 - \lambda_3)|\widehat{v}_1|^2|\widehat{v}_2|^2 + \lambda|\widehat{v}_1 - \widehat{v}_2|^2|\widehat{v}_1 + \widehat{v}_2|^2 \\
&\quad +\tfrac{1}{2}(\lambda_6 - \lambda_7)\left[\widehat{v}_2\widehat{v}_1\left(\widehat{v}_2^{*2} - \widehat{v}_1^{*2}\right) + c.c.\right] , \\
Z_5 &= 2(\lambda_1 + \lambda_2 - \lambda_3)(\widehat{v}_2^*\widehat{v}_1^*)^2 + \lambda(\widehat{v}_1^{*2} - \widehat{v}_2^{*2})^2 + (\lambda_6 - \lambda_7)\widehat{v}_2^*\widehat{v}_1^*(\widehat{v}_2^{*2} - \widehat{v}_1^{*2}) , \\
Z_6 &= \left[\lambda_2|\widehat{v}_2|^2 - \lambda_1|\widehat{v}_1|^2 + \lambda_3(|\widehat{v}_1|^2 - |\widehat{v}_2|^2)\right]\widehat{v}_1^*\widehat{v}_2^* + \lambda\left(\widehat{v}_2^*\widehat{v}_1 + c.c.\right)\left(\widehat{v}_1^{*2} - \widehat{v}_2^{*2}\right) \\
&\quad +\tfrac{1}{2}\left(\lambda_6|\widehat{v}_1|^2 + \lambda_7|\widehat{v}_2|^2\right)\left(\widehat{v}_1^{*2} - \widehat{v}_2^{*2}\right) + \tfrac{1}{2}\left(\lambda_7 - \lambda_6\right)\widehat{v}_1^*\widehat{v}_2^*\left(\widehat{v}_2^*\widehat{v}_1 + c.c.\right) , \\
Z_7 &= \left[\lambda_2|\widehat{v}_1|^2 - \lambda_1|\widehat{v}_2|^2 + \lambda_3(|\widehat{v}_2|^2 - |\widehat{v}_1|^2)\right]\widehat{v}_1^*\widehat{v}_2^* - \lambda\left(\widehat{v}_2^*\widehat{v}_1 + c.c.\right)\left(\widehat{v}_1^{*2} - \widehat{v}_2^{*2}\right) \\
&\quad +\tfrac{1}{2}\left(\lambda_6|\widehat{v}_2|^2 + \lambda_7|\widehat{v}_1|^2\right)\left(\widehat{v}_1^{*2} - \widehat{v}_2^{*2}\right) - \tfrac{1}{2}\left(\lambda_7 - \lambda_6\right)\widehat{v}_1^*\widehat{v}_2^*\left(\widehat{v}_2^*\widehat{v}_1 + c.c.\right) . \qquad (3.88)
\end{aligned}$$

At this point, only Y_1, Y_2, Z_1, Z_2, Z_3, and Z_4 are manifestly real. Now, however, we can apply the custodial symmetry condition from eq. (3.81), and use the fact that all of the m^2 and λ parameters are real. Then we find

$$\begin{aligned}
&Y_3,\ Z_6,\ Z_7 \in \mathbb{R} , \\
&\quad Z_4 = Z_5 = 2(\lambda_1 + \lambda_2 - \lambda_3)\widehat{v}_1^2\widehat{v}_2^2 + \lambda(\widehat{v}_1^2 - \widehat{v}_2^2)^2 + (\lambda_6 - \lambda_7)\widehat{v}_2\widehat{v}_1(\widehat{v}_2^2 - \widehat{v}_1^2) \in \mathbb{R} . \qquad (3.89)
\end{aligned}$$

Since we now have a Lagrangian written in terms of "real basis" parameters, one can say that the condition for custodial invariance of the scalar potential is

$$Z_4 = Z_5 \text{ in the real basis.} \tag{3.90}$$

To check that we do not expect any additional relations among the Z parameters, we can compare the number of degrees of freedom in eq. (3.88) and eq. (3.84). In the most general CP-violating 2HDM, one starts with 6 real and 4 complex parameters in the scalar potential, plus v. From these 15 degrees of freedom, 3 are removed by applying the scalar minimum conditions and one corresponds to an overall phase, which is not physically significant. From these 11 physical degrees of freedom, the conditions of CP conservation [eq. (3.1)] remove three, but there is no longer an overall phase to subtract, so there are 9 independent degrees of freedom. The condition of custodial symmetry removes an additional degree of freedom, leaving eight for the custodially-symmetric potential in eqs. (3.84) and (3.88).

We will now replicate "case II" of Pomarol and Vega and show that it produces exactly the same condition on the parameters of the Lagrangian as eq. (3.90). The alternative to the matrices in eq. (3.78) is the following:

$$M_{21} \equiv (\tilde{\Phi}_2\,,\Phi_1)\,, \tag{3.91}$$

which transforms as

$$M_{21} \to L\ M_{21}\ R^\dagger\,, \tag{3.92}$$

under $SU(2)_L \times SU(2)_R$. Preserving a custodial $SU(2)_V$ symmetry after EWSB requires that the vev of M_{21} be proportional to the identity matrix:

$$< M_{21} >= \frac{1}{\sqrt{2}} \begin{pmatrix} v_2^* & 0 \\ 0 & v_1 \end{pmatrix} \propto \mathbb{1} \Rightarrow v_1^* = v_2\,, \tag{3.93}$$

since for $L = R$,

$$< M_{21} > \to L < M_{21} > R^\dagger = \frac{v_2}{\sqrt{2}} L R^\dagger = < M_{21} > \ . \tag{3.94}$$

From this matrix we construct the following:

$$\begin{aligned}
\mathrm{Tr}[M_{21}^\dagger M_{21}] &= \mathrm{Tr}\left[\begin{pmatrix} \Phi_2^0 & -\Phi_2^+ \\ \Phi_1^- & \Phi_1^{0*} \end{pmatrix}\begin{pmatrix} \Phi_2^{0*} & \Phi_1^+ \\ -\Phi_2^- & \Phi_1^0 \end{pmatrix}\right] = \Phi_1^\dagger\Phi_1 + \Phi_2^\dagger\Phi_2\,, \\
\det[M_{21}^\dagger] &= \det\begin{pmatrix} \Phi_2^0 & -\Phi_2^+ \\ \Phi_1^- & \Phi_1^{0*} \end{pmatrix} = \Phi_1^\dagger\Phi_2\,, \\
\det[M_{21}^\dagger M_{21}] &= \det\left[\begin{pmatrix} \Phi_2^0 & -\Phi_2^+ \\ \Phi_1^- & \Phi_1^{0*} \end{pmatrix}\begin{pmatrix} \Phi_2^{0*} & \Phi_1^+ \\ -\Phi_2^- & \Phi_1^0 \end{pmatrix}\right] \\
&= (\Phi_1^\dagger\Phi_2)(\Phi_2^\dagger\Phi_1)\,.
\end{aligned} \tag{3.95}$$

With the expressions in eq. (3.95), one constructs the following $SU(2)_L \times SU(2)_R$ symmetric scalar potential, again replicating the results in [108]:

$$\begin{aligned}
\mathcal{V} &= m^2\mathrm{Tr}[M_{21}^\dagger M_{21}] - \left(m_{12}^2 \det[M_{21}] + h.c.\right) + \lambda \mathrm{Tr}[M_{21}^\dagger M_{21}]^2 + \lambda_4 \det[M_{21}^\dagger M_{21}] \\
&\quad + \tfrac{1}{2}\left(\lambda_5 \det[M_{21}^\dagger]^2 + \lambda' \det[M_{21}^\dagger]\mathrm{Tr}[M_{21}^\dagger M_{21}] + h.c.\right) . \qquad (3.96)
\end{aligned}$$

Hermiticity implies $m^2, \lambda, \lambda_4 \in \mathbb{R}$. As before, we convert to the Higgs basis:

$$\begin{aligned}
\mathrm{Tr}[M_{21}^\dagger M_{21}] &= H_1^\dagger H_1 + H_2^\dagger H_2 \\
\det[M_{21}^\dagger] &= \widehat{v}_1^*\widehat{v}_2 \left(H_1^\dagger H_1 - H_2^\dagger H_2\right) - \widehat{v}_2^2 H_2^\dagger H_1 + \widehat{v}_1^{*2} H_1^\dagger H_2 \\
\det[M_{21}^\dagger M_{21}] &= \left[-\widehat{v}_2^2\widehat{v}_1^2 (H_2^\dagger H_1)^2 + (|\widehat{v}_1|^2 - |\widehat{v}_2|^2)\widehat{v}_1^*\widehat{v}_2^* \left(H_1^\dagger H_1 - H_2^\dagger H_2\right) H_1^\dagger H_2 + h.c.\right] \\
&\quad + |\widehat{v}_2|^2|\widehat{v}_1|^2 \left(H_1^\dagger H_1 - H_2^\dagger H_2\right)^2 + \left(|\widehat{v}_1|^4 + |\widehat{v}_2|^4\right) H_1^\dagger H_2 H_2^\dagger H_1 . \qquad (3.97)
\end{aligned}$$

Subsituting equations (3.97) into eq. (3.96), we again achieve eq. (3.87), but now with different coefficients:

$$\begin{aligned}
Y_1 &= m^2 - m_{12}^2 \widehat{v}_1^* \widehat{v}_2 - m_{12}^{*2} \widehat{v}_1 \widehat{v}_2^* , \\
Y_2 &= m^2 + m_{12}^2 \widehat{v}_1 \widehat{v}_2^* + m_{12}^{*2} \widehat{v}_1^* \widehat{v}_2 , \\
Y_3 &= -m_{12}^2 \widehat{v}_1^{*2} + \widehat{v}_2^{*2} m_{12}^{*2} , \\
Z_1 &= 2\lambda + 2\lambda_4 |\widehat{v}_1|^2 |\widehat{v}_2|^2 + \left[\lambda_5 (\widehat{v}_1^* \widehat{v}_2)^2 + \lambda' \widehat{v}_1^* \widehat{v}_2 + c.c.\right] , \\
Z_2 &= 2\lambda + 2\lambda_4 |\widehat{v}_1|^2 |\widehat{v}_2|^2 + \left[\lambda_5 (\widehat{v}_1^* \widehat{v}_2)^2 - \lambda' \widehat{v}_1^* \widehat{v}_2 + c.c.\right] , \\
Z_3 &= 2\lambda - 2\lambda_4 |\widehat{v}_1|^2 |\widehat{v}_2|^2 - \left[\lambda_5 (\widehat{v}_1^* \widehat{v}_2)^2 + c.c.\right] , \\
Z_4 &= \lambda_4 \left(|\widehat{v}_1|^4 + |\widehat{v}_2|^4\right) - \left[\lambda_5 (\widehat{v}_1^* \widehat{v}_2)^2 + c.c.\right] , \\
Z_5 &= -2\lambda_4 \widehat{v}_1^2 \widehat{v}_2^2 + \left[\lambda_5 \widehat{v}_1^{*4} + \lambda_5^* \widehat{v}_2^{*4}\right] , \\
Z_6 &= \lambda_4 (|\widehat{v}_1|^2 - |\widehat{v}_2|^2) \widehat{v}_2^* \widehat{v}_1^* + \left(\lambda_5 \widehat{v}_1^{*3} \widehat{v}_2 - \lambda_5^* \widehat{v}_2^{*3} \widehat{v}_1 + \lambda' \widehat{v}_1^{*2} - \lambda'^* \widehat{v}_2^{*2}\right) , \\
Z_7 &= \lambda_4 (|\widehat{v}_2|^2 - |\widehat{v}_1|^2) \widehat{v}_2^* \widehat{v}_1^* + \left(-\lambda_5 \widehat{v}_1^{*3} \widehat{v}_2 + \lambda_5^* \widehat{v}_2^{*3} \widehat{v}_1 + \lambda' \widehat{v}_1^{*2} - \lambda'^* \widehat{v}_2^{*2}\right) . \qquad (3.98)
\end{aligned}$$

As usual, Y_1, Y_2, Z_1, Z_2, Z_3, and Z_4 are manifestly real. Applying eq. (3.93) and the fact that m^2, λ_4 and λ are real, we find that in the custodial limit,

$$\begin{gathered}
iY_3,\ iZ_6,\ iZ_7 \in \mathbb{R} , \\
Z_4 = -Z_5 = 2\lambda_4 |\widehat{v}_1|^4 - \lambda_5 \widehat{v}_2^4 - \lambda_5^* \widehat{v}_1^4 \in \mathbb{R} . \qquad (3.99)
\end{gathered}$$

Again we expect no further relations, since we have reduced the number of independent degrees of freedom in eq. (3.98) to 8, the same as in eq. (3.97). Since the phase of H_2 is not physically meaningful, we are free to transform $H_2 \to iH_2$, which leads to $(Y_3,\ Z_6,\ Z_7) \to -i(Y_3,\ Z_6,\ Z_7)$ and $Z_5 \to -Z_5$, putting all of the parameters in the real basis. Furthermore, since the sign of Z_5 reverses, one can now make the statement that the condition for custodial invariance is "$Z_4 = Z_5$ in the real basis,"[13] as in the previous case. This transformation has no effect on the vevs $< \Phi_i >$, as one can see from eq. (2.11).

Thus, we conclude that it is not possible to distinguish physically between these two "cases" of custodial symmetry–the only difference between them is the condition imposed on the vevs. The relationship between the vevs depends on the choice of basis, and thus cannot be physically measurable.[14]

3.3.1 The Basis-Independent Condition for Custodial Symmetry in the Scalar Sector

It is possible generalize the two implementations of custodial symmetry presented in the previous section by constructing a $SU(2)_L \times SU(2)_R$ invariant scalar potential using Higgs basis fields, which avoids having to impose conditions on the vevs. For the purposes of this section, we will take all scalar couplings to be in the real basis, so that

$$Y_i \in \mathbb{R}, \quad Z_i \in \mathbb{R} \qquad \forall\, i\,. \tag{3.100}$$

This requirement removes the freedom to redefine the phase of H_2 except by an overall sign (which would not change the results presented here). Let us define[15]

$$\mathbb{M}_1 \equiv (\tilde{H}_1\,, H_1), \quad \mathbb{M}_2 \equiv (e^{-i\theta_{23}}\tilde{H}_2\,, e^{i\theta_{23}}H_2)\,, \tag{3.101}$$

where the transformation under a global $SU(2)_L \times SU(2)_R$ is defined as usual:

$$\mathbb{M}_i \to L\ \mathbb{M}_i\ R^\dagger\,. \tag{3.102}$$

In section 3.3 we explicitly required that $< \mathbb{M}_i >$ be proportional to the identity matrix so that the custodial symmetry is preserved after EWSB. Here, since we are writing the fields in the Higgs basis, we do not have impose any specific conditions on the vevs, since $< \mathbb{M}_i >$ are automatically proportional to the unit matrix. Now the $SU(2)_L \times SU(2)_R$

---

[13] This statement is meaningful only if the sign of Z_5 in the real basis is a physical observable. If either $Z_6 \neq 0$ or $Z_7 \neq 0$, the operation $H_2 \to \pm i H_2$ transforms the couplings out of the real basis while changing the sign of Z_5, (as noted above). Hence in this case the sign of Z_5 in the real basis is meaningful. If $Z_6 = Z_7 = 0$, then $H_2 \to \pm i H_2$ changes the sign of Z_5 while preserving the real basis. In this case the sign of Z_5 is not a meaningful and custodial symmetry implies that $Z_4 = |Z_5|$ in the real basis.

[14] In ref. [67], it was shown that for custodial symmetry in the quark-scalar sector of specialized versions of the 2HDM, the two ways to implement custodial symmetry can be distinguished based on the presence of the A^0GG effective interactions in the $E \gg m_W$ limit. However, in their discussion, constraints on the Yukawa couplings effectively select a "preferred" basis.

[15] In the special case of $Z_6 = 0, c_{13} = 0$, and $\text{Im}[Z_5\, e^{2i(\theta_{12}-\theta_{23})}] = 0$, eq. (3.101) applies with the substitution $\theta_{23} \to \theta_{23} - \theta_{12}$. In other mass-degenerate cases that we have treated here, in principle a different combination of mixing angles would play the role of θ_{23}.

invariant potential can be written as follows:

$$\begin{aligned}
\mathcal{V} \;&=\; \tfrac{1}{2}Y_1\mathrm{Tr}[\mathbb{M}_1^\dagger\mathbb{M}_1]+\tfrac{1}{2}Y_2\mathrm{Tr}[\mathbb{M}_2^\dagger\mathbb{M}_2]+Y_3e^{-i\theta_{23}}\mathrm{Tr}[\mathbb{M}_1^\dagger\mathbb{M}_2]+\frac{1}{8}Z_1\left(\mathrm{Tr}[\mathbb{M}_1^\dagger\mathbb{M}_1]\right)^2 \\
&\quad+\frac{1}{8}Z_2\left(\mathrm{Tr}[\mathbb{M}_2^\dagger\mathbb{M}_2]\right)^2+\tfrac{1}{4}Z_3\mathrm{Tr}[\mathbb{M}_1^\dagger\mathbb{M}_1]\mathrm{Tr}[\mathbb{M}_2^\dagger\mathbb{M}_2]+\tfrac{1}{2}\lambda\left(\mathrm{Tr}[\mathbb{M}_1^\dagger\mathbb{M}_2]\right)^2 \\
&\quad+\tfrac{1}{2}\left(Z_6e^{-i\theta_{23}}\mathrm{Tr}[\mathbb{M}_1^\dagger\mathbb{M}_1]+Z_7e^{-i\theta_{23}}\mathrm{Tr}[\mathbb{M}_2^\dagger\mathbb{M}_2]\right)\mathrm{Tr}[\mathbb{M}_1^\dagger\mathbb{M}_2]\,, \qquad (3.103)
\end{aligned}$$

where the coefficients have been adjusted to anticipate comparison with the standard form of the scalar potential. Eq. (3.103) is equivalent to

$$\begin{aligned}
\mathcal{V} \;&=\; Y_1H_1^\dagger H_1+Y_2{H_2}^\dagger H_2+[Y_3H_1^\dagger H_2+\mathrm{h.c.}] \\
&\quad+\tfrac{1}{2}Z_1(H_1^\dagger H_1)^2+\tfrac{1}{2}Z_2({H_2}^\dagger H_2)^2+Z_3(H_1^\dagger H_1)({H_2}^\dagger H_2)+Z_4(H_1^\dagger H_2)({H_2}^\dagger H_1) \\
&\quad+\left\{\tfrac{1}{2}Z_5(H_1^\dagger H_2)^2+\left[Z_6(H_1^\dagger H_1)+Z_7({H_2}^\dagger H_2)\right]H_1^\dagger H_2+\mathrm{h.c.}\right\}\,, \qquad (3.104)
\end{aligned}$$

with the condition

$$Z_4=Z_5e^{-2i\theta_{23}}\,,\quad \mathrm{Im}(Z_6e^{-i\theta_{23}})=0\,,\quad \mathrm{Im}(Z_7\,e^{-i\theta_{23}})=0\,. \qquad (3.105)$$

Note that $\mathrm{Im}[Z_5e^{-2i\theta_{23}}]=0$ since Z_4 is manifestly real. Eq. (3.104) is consistent with the results derived previously in a specific basis. One can rewrite eq. (3.105) so that θ_{23} does appear explicitly. Custodial symmetry requires that $Z_5Z_6^{*2}$, $Z_5Z_7^{*2}$, and $Z_5(\rho^Q)^2$ be real-valued, so eq. (3.105) is equivalent to

$$\begin{aligned}
Z_4 &= \pm\varepsilon_{56}|Z_5| &&\text{if}\, Z_6\neq 0\,, \\
Z_4 &= \pm\varepsilon_{57}|Z_5| &&\text{if}\, Z_6=0\,,\; Z_7\neq 0\,, \\
Z_4 &= \pm\varepsilon_{5Q}|Z_5| &&\text{if}\, Z_6=Z_7=0\,,\; \rho^Q\neq 0\,, \qquad (3.106)
\end{aligned}$$

where the positive sign corresponds to $\mathrm{Im}(Z_6e^{-i\theta_{23}})=\mathrm{Im}(Z_7\,e^{-i\theta_{23}})=\mathrm{Im}(\rho^Qe^{i\theta_{23}})=0$ and the negative sign to $\mathrm{Re}(Z_6e^{-i\theta_{23}})=\mathrm{Re}(Z_7\,e^{-i\theta_{23}})=\mathrm{Re}(\rho^Qe^{i\theta_{23}})=0$. If $Z_6=Z_7=\rho^Q=0$, eq. (3.105) has two solutions, which in the real basis can be written

$$Z_4=\pm|Z_5|\,, \qquad (3.107)$$

reflecting the two possible definitions of CP symmetry which arise in this case. Since both CP symmetries are conserved, the condition in eq. (3.107) is indeterminate.

3.3.2 Degeneracy in the Custodial Limit

Finally, one important consequence of custodial symmetry in the Higgs sector is that the charged Higgs boson is always degenerate with one of the neutral Higgs fields. Since the custodial limit is CP-conserving, we can write the following expressions for the CP-odd mass from eq. (3.58):

$$\begin{aligned}
m_{A^0}^2 \;&=\; Y_2+\tfrac{1}{2}[Z_3+Z_4-\mathrm{Re}(Z_5\,e^{-2i\theta_{23}})]v^2\,, \\
&\quad\;\text{for }\;\mathrm{Im}(Z_6e^{-i\theta_{23}})=\mathrm{Im}(Z_7\,e^{-i\theta_{23}})=\mathrm{Im}(\rho^Qe^{i\theta_{23}})=0\,. \qquad (3.108)
\end{aligned}$$

Note that this is the only possible value of $m^2_{A^0}$ since the condition for custodial symmetry in eq. (3.105) eliminates the possibility that eq. (3.59) is correct (as long as either Z_6 or Z_7 is non-zero). Thus, the condition $Z_4 = \mathrm{Re}(Z_5\, e^{-2i\theta_{23}})$ of eq. (3.106) implies a degeneracy between the CP-odd Higgs and the charged Higgs, ie,

$$m^2_{A^0} = m^2_{H^\pm} = Y_2 + \tfrac{1}{2} Z_3 v^2\,. \tag{3.109}$$

This observation agrees with that of [108]. However, in the case of $Z_6 = Z_7 = 0$, there is another possibility for the mass of the CP-odd field from eq. (3.59),

$$m^2_{A^0} = Y_2 + \tfrac{1}{2}[Z_3 + Z_4 + \mathrm{Re}(Z_5\, e^{-2i\theta_{23}})]v^2 \quad \text{for } \mathrm{Re}(\rho^Q e^{i\theta_{23}}) = 0\,. \tag{3.110}$$

This unique case arises because $\rho^Q e^{i\theta_{23}}$ can be either imaginary or real, unlike $Z_{6,7}e^{-i\theta_{23}}$ which must be real for the scalar potential to be custodially symmetric. (As we will see in section 3.3.4, imposing custodial symmetry on the Higgs-Quark sector does not require $\rho^Q e^{i\theta_{23}}$ to be real.) In this case it is the CP-even field H^0 that is degenerate with the charged Higgs boson:

$$m^2_{H^0} = m^2_{H^\pm} = Y_2 + \tfrac{1}{2} Z_3 v^2\,. \tag{3.111}$$

These two possibilities for degeneracy arise from the two possible definitions of CP mentioned in section 3.2.3. This phenomenon of a degenerate charged Higgs boson and CP-even Higgs boson was described for the case $Z_6 = Z_7 = 0$ in [46], in the context of a "twisted" custodial symmetry. In fact, this scenario can be analyzed without the authors' "twisting" formalism, as we have seen here. Finally, we consider the case $Z_6 = Z_7 = \rho^Q = 0$. With nothing to select out one of the two possible definitions of CP, it cannot be determined whether it is the CP-even or CP-odd neutral Higgs boson that is degenerate. We emphasize, in contrast to the authors of [46], that these two possbilities in eqs. (3.109) and (3.111) arise as a result of the two inequivalent definitions of the CP transformation, which exist (independently of the custodial limit) at the special point of parameter space where $Z_6 = Z_7 = \rho^Q = 0$. This is the only case in which the mass-degeneracy $m_{H^\pm} = m_{H^0}$ corresponds to a custodial symmetry.

3.3.3 Basis-Dependent Formulations of Custodial Symmetry in the Higgs-Quark Sector

Let us now extend the two ways to impose $SU(2)_L \times SU(2)_R$ invariance discussed in the previous section to the Higgs-Quark sector. As before, in the final analysis these two cases defined by Pomarol and Vega will be related by basis-transformations, and do not represent the most general way of implementing the symmetry. In Pomarol and Vega's "case I," equation (2.75) becomes

$$-\mathscr{L}_{\mathrm{Y}} = \eta_1 \overline{\mathcal{Q}_L}\; M_1 \begin{pmatrix} \mathcal{U}_R \\ D_R \end{pmatrix} + \eta_2 \overline{\mathcal{Q}_L}\; M_2 \begin{pmatrix} \mathcal{U}_R \\ D_R \end{pmatrix} + \text{h.c.}\,, \tag{3.112}$$

with M_i defined in eq. (3.78). This Lagrangian is manifestly invariant under $SU(2)_L \times SU(2)_R$ given the transformations

$$\begin{array}{rcl} M_i & \rightarrow & L M_i R^\dagger \,, \\ \begin{pmatrix} \mathcal{U}_R \\ D_R \end{pmatrix} & \rightarrow & R \begin{pmatrix} \mathcal{U}_R \\ D_R \end{pmatrix} , \\ \overline{\mathcal{Q}_L} & \rightarrow & \overline{\mathcal{Q}_L} L^\dagger \,. \end{array} \tag{3.113}$$

Comparing with eq. (2.75), one finds the conditions

$$\eta_1^U = \eta_1^{D\dagger} \,, \qquad \eta_2^U = \eta_2^{D\dagger} \,. \tag{3.114}$$

With this constraint, the parameters κ^Q and ρ^Q defined in eq. (2.78) become

$$\begin{array}{llll} \kappa^U & = v_1^* \eta_1^U + v_2^* \eta_2^U \,, & \kappa^{D\dagger} & = v_1 \eta_1^U + v_2 \eta_2^U \,, \\ \rho^U & = -v_2 \eta_1^U + v_1 \eta_2^U \,, & \rho^{D\dagger} & = -v_2^* \eta_1^U + v_1^* \eta_2^U \,. \end{array} \tag{3.115}$$

Since the v_i are real in this case [equation (3.81)], we find $\kappa^U = \kappa^{D\dagger}$, or simply $M_U = M_D$. Similarly, $\rho^{D\dagger} = \rho^U$. The parameters ρ^Q are not quite basis-independent, since they pick up a factor of det(U) under a U(2) transformation. However, from the previous section, we know that we are in the real basis. Thus, one state unambiguous conditions for custodial symmetry in the Yukawa sector:

$$M_U = M_D, \;\; \rho^U = \rho^{D\dagger} \text{ in the real basis.} \tag{3.116}$$

Let us now do the same for Pomarol and Vega's "case II." We construct

$$-\mathscr{L}_{\mathrm{Y}} = \eta_{21} \overline{\mathcal{Q}_L} \; M_{21} \begin{pmatrix} \mathcal{U}_R \\ D_R \end{pmatrix} + \eta_{12} \overline{\mathcal{Q}_L} \; M_{12} \begin{pmatrix} \mathcal{U}_R \\ D_R \end{pmatrix} + \text{h.c.} \,, \tag{3.117}$$

with M_{21} defined in eq. (3.91), and a similar matrix defined as

$$M_{12} \equiv \tilde{\Phi}_1^* \Phi_2 \,. \tag{3.118}$$

This Lagrangian is invariant under $SU(2)_L \times SU(2)_R$ given

$$\begin{array}{rcl} M_{12} & \rightarrow & L M_{12} R^\dagger \,, \\ M_{21} & \rightarrow & L M_{21} R^\dagger \,. \end{array} \tag{3.119}$$

Comparing eq. (3.117) with eq. (2.75), one finds the conditions

$$\eta_1^U = \eta_2^{D\dagger} \,, \qquad \eta_2^U = \eta_1^{D\dagger} \,. \tag{3.120}$$

Substituting equations (3.120) and (3.93) into eq. (2.78) yields

$$\begin{array}{llll} \kappa^U & = v_1^* \eta_1^U + v_2^* \eta_2^U \,, & \kappa^{D\dagger} & = v_2^* \eta_2^U + v_1^* \eta_1^U \,, \\ \rho^U & = -v_2 \eta_1^U + v_1 \eta_2^U \,, & \rho^{D\dagger} & = v_2 \eta_1^U - v_1 \eta_2^U \,. \end{array} \tag{3.121}$$

Thus, one again finds $M_U = M_D$. We also have $\rho^{D\dagger} = -\rho^U$. From the previous section, we know that to express these quantities in the real basis, we need to make the transformation $H_2 \to iH_2$, which is equivalent to

$$\rho^U \to i\rho^U \,, \qquad\qquad \rho^{D\dagger} \to -i\rho^{D\dagger}. \tag{3.122}$$

Thus, the conditions for custodial invariance of the Yukawa sector are given by eq. (3.116), as in the previous case.

3.3.4 The Basis-Independent Custodially-Symmetric Higgs-Quark Lagrangian

Now we will determine, in analogy to the previous section, the form of the $SU(2)_L \times SU(2)_R$-invariant Higgs-Quark Lagrangian, assuming that the fields and scalar couplings are in the real basis. Let us rewrite eq. (2.77) as follows:

$$-\mathscr{L}_{\rm Y} = \overline{\mathcal{Q}_L}(\tilde{H}_1\kappa^U + \tilde{H}_2\rho^U)\mathcal{U}_R + \overline{\mathcal{Q}_L}(H_1{\kappa^D}^\dagger + H_2{\rho^D}^\dagger)D_R + \text{h.c.}\,. \tag{3.123}$$

Then the desired lagrangian is of the form

$$-\mathscr{L}_{\rm Y} = \kappa\overline{\mathcal{Q}_L}\ \mathbb{M}_1 \begin{pmatrix} \mathcal{U}_R \\ D_R \end{pmatrix} + \rho\overline{\mathcal{Q}_L}\ \mathbb{M}_2 \begin{pmatrix} \mathcal{U}_R \\ D_R \end{pmatrix} + \text{h.c.}\,, \tag{3.124}$$

with $\mathbb{M}_1 \equiv \tilde{H}_1\ H_1$ and $\mathbb{M}_2 \equiv (e^{-i\theta_{23}}\tilde{H}_2\,, e^{i\theta_{23}}H_2)$ as in eq. (3.101). Comparing to eq. (3.123), one finds that the condition for invariance under $SU(2)_L \times SU(2)_R$ is

$$\kappa^U = \kappa^{D\dagger}, \qquad e^{-i\theta_{23}}\rho^U = e^{i\theta_{23}}\rho^{D\dagger}\,. \tag{3.125}$$

Again, one can recover Pomarol and Vega's two cases by choosing to constrain the vevs of the Higgs fields. However, if there is no special symmetry imposed on the Yukawa sector (as we are assuming here), then there is nothing to select a "preferred" basis. There are an infinite number of unphysical basis choices which can be made, but one can always write down unambiguous conditions either by going to real basis, or by using invariant combinations as we have done in this section.

Chapter 4

Phenomenology of the 2HDM

4.1 The Oblique Parameters S, T and U

The parameters S, T, and U are independent UV-finite combinations of radiative corrections to gauge boson vacuum polarization diagrams (aka "oblique" corrections). T is related to the parameter ρ by $\rho - 1 = \alpha T$. They are calculated from the transverse part of the gauge boson two-point function:

$$\begin{aligned}
S &\equiv -\frac{16\pi c_W}{s_W g^2}\frac{d}{dq^2}\Pi_{W^3B}(q^2)|_{q^2=0} && (4.1)\\
\alpha T &\equiv \frac{1}{m_W^2}\left[\Pi_{W^1W^1}(0) - \Pi_{W^3W^3}(0)\right], && (4.2)\\
U &\equiv \frac{16\pi}{g^2}\frac{d}{dq^2}\left[\Pi_{W^1W^1}(q^2) - \Pi_{W^3W^3}(q^2)\right]|_{q^2=0}\,. && (4.3)
\end{aligned}$$

Here, $\Pi_{ij}(q^2)$ is defined by

$$i\Pi_{ij}^{\mu\nu} = ig^{\mu\nu}\Pi_{ij}(q^2) + (q^\mu q^\nu \text{ terms})\,. \tag{4.4}$$

Some care must be taken as this differs from the convention of [8, 103, 104], and others, who pull out factors of g^2, etc.

The "linear expansion approximation" is often used in the literature, which extracts the dependence of the functions on q^2, as in refs. [103] and [104]:

$$\Pi_{ij}(q^2) = A_{ij}(0) + q^2 F_{ij}(q^2)\,. \tag{4.5}$$

For the case that the scale of the new physics is much greater than m_Z, one can take the momentum scale q^2 to be of order m_Z^2, and define the oblique parameters as follows [64]:

$$\begin{aligned}
\frac{g^2}{16\pi}S &\equiv c_W^2\left[F_{ZZ}(m_Z^2) - F_{\gamma\gamma}(m_Z^2) + \left(\frac{2s_W^2-1}{s_W c_W}\right)F_{Z\gamma}(m_Z^2)\right], && (4.6)\\
\alpha T &\equiv \frac{A_{WW}(0)}{m_W^2} - \frac{A_{ZZ}(0)}{m_Z^2} && (4.7)\\
\frac{g^2}{16\pi}(S+U) &\equiv F_{WW}(m_W^2) - F_{\gamma\gamma}(m_W^2) - \frac{c_W}{s_W}F_{Z\gamma}(m_W^2)\,. && (4.8)
\end{aligned}$$

The combination $S+U$ rather than U has been taken for calculational simplicity. S, T and U are defined relative to the Standard Model, so that extending the Higgs sector has the effect of shifting the parameters away from zero.[1] Thus, in the calculations that follow, contributions from Standard Model processes have been subtracted out. The Higgs mass used in this "reference" Standard Model (m_ϕ) will be left arbitrary for the moment.

The result for S is calculated in Appendix E.1 and found to be:

$$\begin{aligned} S &= \frac{1}{\pi m_Z^2}\left[q_{k1}^2\mathcal{B}_{22}(m_Z^2;m_Z^2,m_k^2) - \mathcal{B}_{22}(m_Z^2;m_Z^2,m_\phi^2) - m_Z^2 q_{k1}^2\mathcal{B}_0(m_Z^2;m_Z^2,m_k^2)\right. \\ &\quad + m_Z^2\mathcal{B}_0(m_Z^2;m_Z^2,m_\phi^2) + q_{11}^2\mathcal{B}_{22}(m_Z^2;m_2^2,m_3^2) + q_{21}^2\mathcal{B}_{22}(m_Z^2;m_1^2,m_3^2) \\ &\quad \left. + q_{31}^2\mathcal{B}_{22}(m_Z^2;m_1^2,m_2^2) - \mathcal{B}_{22}(m_Z^2;m_{H^\pm}^2,m_{H^\pm}^2)\right] . \end{aligned} \tag{4.9}$$

Here $m_{H^\pm}$ represents the mass of the charged Higgs $H^\pm$, as usual, and m_k the masses of the neutral Higgs h_k ($k=1,2,3$). Repeated indices are summed over. The notation $\mathcal{B}_{22}$ and $\mathcal{B}_0$ was introduced in [64]:

$$\mathcal{B}_{22}(q^2;m_1^2,m_2^2) \equiv B_{22}(q^2;m_1^2,m_2^2) - B_{22}(0;m_1^2,m_2^2)\,, \tag{4.10}$$

$$\mathcal{B}_0(q^2;m_1^2,m_2^2) \equiv B_0(q^2;m_1^2,m_2^2) - B_0(0;m_1^2,m_2^2)\,. \tag{4.11}$$

The functions B_{22} and B_0 are defined in ref. [101] and come from the evaluation of two-point integrals. They can be evaluated using the following formulae of ref. [64]:

$$\begin{aligned} B_{22}(q^2;m_1^2,m_2^2) &= \frac{1}{4}(\Delta+1)[m_1^2+m_2^2-\frac{1}{3}q^2] - \tfrac{1}{2}\int_0^1 dx X\ln(X-i\epsilon)\,, \\ B_0(q^2;m_1^2,m_2^2) &= \Delta - \int_0^1 dx\ln(X-i\epsilon)\,, \\ A_0(m^2) &= m^2(\Delta+1-\ln m^2)\,, \end{aligned} \tag{4.12}$$

where $X \equiv m_1^2 x + m_2^2(1-x) - q^2 x(1-x)$ and $\Delta \equiv \frac{2}{4-d} + \ln(4\pi) - \gamma$, in d space-time dimensions.

The calculation of T and $S+U$ is also undertaken in the Appendix. The result is (summing over repeated indices)

---

[1] Contributions to the oblique parameters from the Higgs-fermion Yukawa couplings will not be included here, since they arise from diagrams of order two-loops and higher.

$$\begin{aligned}
T &= \frac{1}{16\pi m_W^2 s_W^2}\left\{|q_{k2}|^2 F(m_{H^\pm}^2, m_k^2) - q_{21}^2 F(m_1^2, m_3^2) - q_{11}^2 F(m_2^2, m_3^2)\right.\\
&\quad -q_{31}^2 F(m_1^2, m_2^2) + q_{k1}^2[F(m_W^2, m_k^2) - F(m_Z^2, m_k^2)] + 4m_W^2 B_0(0; m_W^2, m_\phi^2)\\
&\quad -4m_Z^2 B_0(0; m_Z^2, m_\phi^2) - 4q_{k1}^2[m_W^2 B_0(0; m_W^2, m_k^2) - m_Z^2 B_0(0; m_Z^2, m_k^2)]\\
&\quad \left. +F(m_Z^2, m_\phi^2) - F(m_W^2, m_\phi^2)\right\},\\
S+U &= \frac{1}{\pi m_W^2}\Big[-q_{k1}^2 m_W^2 \mathcal{B}_0(m_W^2; m_W^2, m_k^2) + m_W^2 \mathcal{B}_0(m_W^2; m_W^2, m_\phi^2)\\
&\quad -\mathcal{B}_{22}(m_W^2; m_W^2, m_\phi^2). + q_{k1}^2 \mathcal{B}_{22}(m_W^2; m_W^2, m_k^2)\\
&\quad +|q_{k2}|^2 \mathcal{B}_{22}(m_W^2; m_{H^\pm}^2, m_k^2) - 2\mathcal{B}_{22}(m_W^2; m_{H^\pm}^2, m_{H^\pm}^2)\Big],
\end{aligned} \tag{4.13}$$

where

$$F(m_1^2, m_2^2) \equiv \tfrac{1}{2}(m_1^2 + m_2^2) - \frac{m_1^2 m_2^2}{m_1^2 - m_2^2}\ln\left(\frac{m_1^2}{m_2^2}\right). \tag{4.14}$$

One can check that

$$F(m^2, m^2) = 0\,. \tag{4.15}$$

This calculation T has been exhibited in a basis-dependent formalism in [51]Grimus:2008nb, and the result is consistent with eq. (4.13).

4.1.1 The Parameter T and the Custodial Limit

The result for T is exhibited in a different form in [108], to emphasize which terms in T arise from the breaking of the custodial symmetry and which arise from the non-custodially-invariant terms in the scalar potential. In the custodial limit, $g' \to 0$ and $m_Z = m_W$. The terms resulting from the gauging of hypercharge, which vanish in this limit, may be rewritten so that they are proportional to g'^2:

$$\begin{aligned}
\alpha T &= \frac{g'^2}{64\pi^2}\sum_{k=1}^{3}\frac{q_{k1}^2}{m_W^2 - m_Z^2}\left\{F(m_k^2, m_Z^2) - F(m_k^2, m_W^2) + F(m_\phi^2, m_W^2) - F(m_\phi^2, m_Z^2)\right.\\
&\qquad +4[m_Z^2 B_0(0; m_Z^2, m_\phi^2) - m_W^2 B_0(0; m_W^2, m_\phi^2)]\\
&\qquad \left. +4[m_W^2 B_0(0; m_W^2, m_k^2) - m_Z^2 B_0(0; m_Z^2, m_k^2)]\right\}\\
&\quad +\frac{g^2}{64\pi^2 m_W^2}\left[\sum_{k=1}^{3}|q_{k2}|^2 F(m_k^2, m_{H^\pm}^2) - q_{11}^2 F(m_2^2, m_3^2) - q_{21}^2 F(m_1^2, m_3^2)\right.\\
&\quad \left. -q_{31}^2 F(m_2^2, m_1^2)\right],
\end{aligned} \tag{4.16}$$

where $\alpha \equiv \frac{e^2}{4\pi}$. Let us now focus on the terms proportional to g^2 in eq. (4.16). One can verify that

$$\sum_{k=1}^{3} |q_{k2}|^2 F(m_k^2, m_{H^\pm}^2) - q_{11}^2 F(m_2^2, m_3^2) - q_{21}^2 F(m_1^2, m_3^2) - q_{31}^2 F(m_2^2, m_1^2) = 0\,, \qquad \text{[custodial limit]} \quad (4.17)$$

due to the degeneracy between one of the neutral fields and the charged Higgs boson. For example, in CP Case I [see Table 3.1], the term displayed in eq. (4.17) is proportional to

$$s_{12}^2 F(m_1^2, m_{H^\pm}^2) + c_{12}^2 F(m_2^2, m_{H^\pm}^2) + F(m_3^2, m_{H^\pm}^2) - c_{12}^2 F(m_2^2, m_3^2) - s_{12}^2 F(m_1^2, m_3^2)\,. \qquad (4.18)$$

Due to the degeneracy between the charged Higgs boson and the CP-odd field m_3, $F(m_3^2, m_{H^\pm}^2) = 0$ and the rest of the terms cancel. One may check the other cases of CP symmetry in a similar manner. In the special case of $Z_6 = Z_7 = 0$, the charged Higgs may be degenerate with a CP-even field rather than the CP-odd field. One can verify for the six cases of CP symmetry (i)–(vi) that eq. (4.17) vanishes when $m_{H^\pm}^2 = m_{H^0}^2$. [For example, in case (i), this term is proportional to $F(m_{H^0}^2, m_{H^\pm}^2) + F(m_{A^0}^2, m_{H^\pm}^2) - F(m_{H^0}^2, m_{A^0}^2) = 0$.] Thus, the only non-zero contribution to αT arises from the breaking of custodial symmetry [the first part of eq. (4.16)].

4.1.2 S, T, and U in the CP-Conserving Limit

The expressions for S, T, and U in the CP-conserving limit can be calculated using Tables 3.1- 3.3. Note that Case II has been divided into two subcases based on the mass ordering of the neutral scalars, Case IIa ($m_{h^0} < m_{A^0}$) and Case IIb ($m_{h^0} > m_{A^0}$). Plugging in the values of the $q_{k\ell}$ parameters from the tables into equations (4.9) and (4.13), one reproduces the results of ref. [64] (the reference mass m_ϕ has been fixed to be m_{h^0}) :

$$\begin{aligned}
S &= \frac{1}{\pi m_Z^2}\Big\{\sin^2(\beta-\alpha)\mathcal{B}_{22}(m_Z^2;m_{H^0}^2,m_{A^0}^2)+\cos^2(\beta-\alpha)\left[\mathcal{B}_{22}(m_Z^2;m_{h^0}^2,m_{A^0}^2)\right.\\
&\quad+\mathcal{B}_{22}(m_Z^2;m_Z^2,m_{H^0}^2)-\mathcal{B}_{22}(m_Z^2;m_Z^2,m_{h^0}^2)-m_Z^2\mathcal{B}_0(m_Z^2;m_Z^2,m_{H^0}^2)\\
&\quad\left.+m_Z^2\mathcal{B}_0(m_Z^2;m_Z^2,m_{h^0}^2)\right]-\mathcal{B}_{22}(m_Z^2;m_{H^\pm}^2,m_{H^\pm}^2)\Big\},\\
T &= \frac{1}{16\pi s_W^2 m_W^2}\Big\{F(m_{H^\pm}^2,m_{A^0}^2)+\sin^2(\beta-\alpha)[F(m_{H^\pm}^2,m_{H^0}^2)-F(m_{A^0}^2,m_{H^0}^2)]\\
&\quad+\cos^2(\beta-\alpha)[F(m_{H^\pm}^2,m_{h^0}^2)-F(m_{A^0}^2,m_{h^0}^2)+F(m_W^2,m_{H^0}^2)-F(m_W^2,m_{h^0}^2)\\
&\quad-F(m_Z^2,m_{H^0}^2)+F(m_Z^2,m_{h^0}^2)+4m_Z^2B_0(0;m_Z^2,m_{H^0}^2)\\
&\quad-4m_Z^2B_0(0;m_Z^2,m_{h^0}^2)-4m_W^2B_0(0;m_W^2,m_{H^0}^2)+4m_W^2B_0(0;m_W^2,m_{h^0}^2)]\Big\},\\
S+U &= \frac{1}{\pi m_W^2}\Big\{\mathcal{B}_{22}(m_Z^2;m_{H^\pm}^2,m_{A^0}^2)-2\mathcal{B}_{22}(m_Z^2;m_{H^\pm}^2,m_{H^\pm}^2)\\
&\quad+\sin^2(\beta-\alpha)\mathcal{B}_{22}(m_W^2;m_{H^\pm}^2,m_{H^0}^2)+\cos^2(\beta-\alpha)[\mathcal{B}_{22}(m_W^2;m_{h^0}^2,m_{H^\pm}^2)\\
&\quad+\mathcal{B}_{22}(m_W^2;m_W^2,m_{H^0}^2)-\mathcal{B}_{22}(m_W^2;m_W^2,m_{h^0}^2)\\
&\quad+m_W^2\mathcal{B}_0(m_W^2;m_W^2,m_{h^0}^2)-m_W^2\mathcal{B}_0(m_W^2;m_W^2,m_{H^0}^2)]\Big\} \qquad (4.19)
\end{aligned}$$

The angle $\beta-\alpha$ is defined in eq. (3.38). Note that these expressions hold for both cases of CP conservation defined in section 3.1.1.

4.2 Numerical Analysis

The experimental determinations of S, T and U are as follows [6]:

$$\begin{aligned}
S &= -0.10\pm .10,\\
T &= -0.08\pm .11,\\
U &= 0.15\pm .11, \qquad (4.20)
\end{aligned}$$

for the value of the Standard Model Higgs mass $m_\phi = 117$ GeV. The values in eq. (4.20) have the Standard Model contributions subtracted out, so that they reflect the deviation from the SM prediction.

Fixing $U=0$, as required by some models, changes the experimental limits on S and T slightly due to correlations between the parameters. The relevant constraints in the $U=0$ scenario for $m_\phi = 117$ GeV are shown in ref. [6] to be consistent with no deviation from the Standard Model:

$$S=-0.04\pm .09,\quad T=0.02\pm .09 \qquad (4.21)$$

For $m_\phi = 300$ GeV, the corresponding values are

$$S = -0.07 \pm .09, \quad T = 0.09 \pm .09 \tag{4.22}$$

These limits indicate that new physics contributions to the oblique parameters are tightly constrained. Their significance for the parameters of the 2HDM is the subject of this section.

The parameters of the 2HDM which are constrained by S, T, and U can be taken to be Z_1, Z_3, $Z_3 + Z_4$, $Z_5\, e^{-2i\theta_{23}}$, $Z_6\, e^{-i\theta_{23}}$ and Y_2, since these 6 quantities determine the physical Higgs masses [see eqs. (2.41) and (2.23)] and the invariant functions $q_{k\ell}$ [see Appendix C]. At this point, it will be assumed that the lightest neutral Higgs mass (m_1) will be interpreted as the reference mass m_ϕ.

The procedure used here to study the effect of the 2HDM on the oblique parameters was to choose random values of the six parameters in the space allowed by the unitarity bounds calculated in Appendix F, subject to the additional requirement that m_1 falls within 15 GeV of 117 GeV, since m_ϕ was taken to be 117 GeV in the PDG analysis. Then the Higgs masses and $q_{k\ell}$ are calculated numerically and inserted into eq. (4.9) and eq. (4.13) to obtain S, T, and U for each point in parameter space.

It was found that the 2HDM consistently produces values of U within .02 of zero. This fact has important ramifications for comparison with experiment, since the limits in eq. (4.21) must be used to constrain S and T, rather than the generic limits in eq. (4.20). Scanning the parameter space and comparing with the allowed "ellipse" in $S - T$ space produces the results shown in Fig. 4.1. From the scatterplot it is evident that the values of S produced are greater than indicated by the central value of $-.04$, although the discrepancy is not statistically significant. Since the values of T cover a broad range, the experimental value of T is easily accomodated.

Meanwhile, the interplay between the constraints from perturbative unitarity and the experimental limits on the oblique parameters is illustrated in Figs. 4.2(a) and (b), which display S and T as a function of the charged Higgs mass $m_{H^\pm}$. In these plots, we have fixed $Y_2 = m_W^2$ so that $m_{H^\pm}$ is limited by the unitarity bound on Z_3. (Recall that $m_{H^\pm}^2 = Y_2 + \frac{1}{2} Z_3 v^2$.) The remaining parameters, shown below, are chosen such that $m_1 = 117$ GeV, to facilitate comparison with eq. (4.21):

$$\begin{aligned} Z_1 = &\quad 0.31\,, & Z_3 + Z_4 = &\quad 24, \\ \mathrm{Re}(Z_6\, e^{-i\theta_{23}}) = &\quad 0.1\,, & \mathrm{Im}(Z_6\, e^{-i\theta_{23}}) = &\quad -1.0, \\ \mathrm{Re}(Z_5\, e^{-2i\theta_{23}}) = &\quad -1.0\,, & \mathrm{Im}(Z_5\, e^{-2i\theta_{23}}) = &\quad 1.0. \end{aligned} \tag{4.23}$$

Fixing the quantity $Z_3 + Z_4$, rather than Z_4 alone, ensures that m_1 (and thus m_ϕ) does not vary as Z_3 is increased. This simplifies the task of using the experimental limits in eq. (4.21). The sharp minimum in Fig. 4.2(b) results from the fact that corrections to T are minimized when the difference between the charged Higgs and heavy neutral Higgs masses is small. This effect will be exhibited in more detail in section 4.2.1.

The values for the Z_i in eq. (4.23) and our choice of $Y_2 = (80.4\ \text{GeV})^2$ generate



(a)

(b)

Figure 4.1: Scatterplots for T as a function of S, with $m_1 = 117 \pm 15$ GeV. The ellipse, representing the 2 σ contour, is adapted from [6]. The second plot shows a close-up of the allowed region.

the following masses for the Higgs particles:

$$\begin{aligned} m_1 &= 117 \text{ GeV}\,, \\ m_2 &= 831 \text{ GeV}\,, \\ m_3 &= 882 \text{ GeV}\,. \end{aligned} \tag{4.24}$$



(a)

(b)

Figure 4.2: S and T shown as a function of $m_{H^\pm}$ (solid line), with Y_2 fixed to be m_W^2. The mass scale is in GeV. The unitarity bound (dashed line) applies to Z_3; all other parameters were picked so as to undersaturate unitarity. The 1σ and 2σ contours (shaded regions) are obtained from eq. (4.21).

The charged Higgs mass can be at most 875 GeV for $Y_2 = m_W^2$, based on the unitarity limit $Z_3 \leq 8\pi$.

This numerical analysis shows that the current experimental limits on the oblique parameters are consistent with contributions to new physics from a second Higgs doublet over a large region of the 2HDM parameter space. However, experimental bounds on the S, T, and U parameters do impose some non-trivial constraints. Our results also suggest that the 2HDM favors a slightly positive value of S and a value of U within .02 of zero, which is consistent with present data within the statistical error. The 2HDM produces a broad range of values for T, which overlap completely with the experimental 2 σ bounds. Finally, we have shown that taken together, the bounds on S, T, and U and the unitarity limits act synergistically to restrict much the 2HDM parameter space.

4.2.1 Splitting Between Neutral and Charged Higgs Masses

One can consider the case where the neutral masses m_1, m_2, and m_3 are light, and the charged Higgs mass heavier ($m_{H^\pm} > m_3$). For a given choice of Z_3, the size of the splitting $\Delta m \equiv m_{H^\pm} - m_3$ is maximized when the other Z_i are small. As we will see in this section, the measured value of T puts a limit on Δm. The following values for the Z_i have been chosen to maximize Δm, while preserving our requirement that $m_1 \approx 117$ GeV at $Y_2 = m_W^2$:

$$\begin{aligned} Z_1 &= 0.25\,, & \mathrm{Re}(Z_6\, e^{-i\theta_{23}}) &= 0.01, \\ Z_3 + Z_4 &= .26\,, & \mathrm{Im}(Z_6\, e^{-i\theta_{23}}) &= 0.01, \\ \mathrm{Re}(Z_5\, e^{-2i\theta_{23}}) &= 0.01\,, & \mathrm{Im}(Z_5\, e^{-2i\theta_{23}}) &= 0.01. \end{aligned} \tag{4.25}$$

With the parameters fixed as in eq. (4.25), the size of the splitting (controlled by Z_3) can easily become large; for example, $\Delta m = 400$ GeV at $Z_3 = 9$. However, one can now impose constraints from the oblique parameters. Using the values of the Z_i in eq. (4.25), S and T as functions of Δm are shown in Figs. 4.3(a) and (b), respectively.

In these plots, as in the previous section, the experimental limits are taken from eq. (4.21), since $U \approx 0$. The graph of T shows that the mass of the charged Higgs can at most be roughly 100 GeV greater than m_3 and still be consistent with the experimental limits, which corresponds to the value $Z_3 = 1.5$.

4.3 Measuring $tan(\beta)$

In sections 2.5 and 2.6, we have written out the entire interaction Lagrangian for the Higgs bosons of the 2HDM. Yet, the famous parameter $\tan\beta$, given by $\tan\beta \equiv v_2/v_1$ in a generic basis [see eq. (2.3)], does not appear in any physical Higgs (or Goldstone) boson coupling. This is rather surprising given the large literature of 2HDM phenomenology in which the parameter $\tan\beta$ is ubiquitous. For example, numerous methods have been proposed for measuring $\tan\beta$ at future colliders [42, 81, 15, 62, 19, 36, 31, 84]. In a generic basis, one can also define the relative phase of the two vacuum expectation values, $\xi = \arg\,(v_2 v_1^*)$. However, neither $\tan\beta$ nor ξ are basis-independent. One can remove ξ by rephasing one of the two Higgs doublet fields, and both ξ and $\tan\beta$ can be removed entirely by transforming to the Higgs basis. Thus, in a general 2HDM, $\tan\beta$ is an unphysical parameter with no significance *a priori*.

The true significance of $\tan\beta$ emerges only in specialized versions of the 2HDM, where $\tan\beta$ is promoted to a physical parameter. As noted in section 2.6, the general 2HDM generally predicts FCNCs in conflict with experimental data. One way to avoid this phenomenological problem is to constrain the theoretical structure of the 2HDM. Such constraints often pick out a preferred basis. Relative to that basis, $\tan\beta$ is then a meaningful parameter.

The most common 2HDM constraint is the requirement that some of the Higgs-fermion Yukawa couplings vanish in a "preferred" basis. This leads to the well known



Figure 4.3: S and T shown as a function of the mass splitting between the charged Higgs and the heaviest neutral Higgs. The mass scale is in GeV. The central values (dashed lines) and the 1σ and 2σ contours (shaded regions) are obtained from eq. (4.21). Splitting of more than 100 GeV is disfavored by the constraints from T.

type-I and type-II 2HDMs [69] (henceforth called 2HDM-I and 2HDM-II). In the 2HDM-I, there exists a preferred basis where $\eta_2^U = \eta_2^D = 0$ [63, 69]. In the 2HDM-II, there exists a preferred basis where $\eta_1^U = \eta_2^D = 0$ [37, 69]. These conditions can be enforced by a suitable symmetry. For example, the MSSM possesses a type-II Higgs-fermion interaction, in which case the supersymmetry guarantees that $\eta_1^U = \eta_2^D = 0$. In non-supersymmetric models, appropriate discrete symmetries can be found to enforce the type-I or type-II Higgs-fermion couplings.[2]

The conditions for type-I and type-II Higgs-fermion interactions given above are basis-dependent. But, there is also a basis-independent criterion that was first given in ref. [34]:[3]

---

[2] These discrete symmetries also imply that some of the coefficients of the scalar potential must also vanish in the same preferred basis [63, 37, 69, 87].

$$
\begin{aligned}
\epsilon_{\bar{a}\bar{b}}\eta^D_a\eta^U_b = \epsilon_{ab}\eta^{D\,\dagger}_{\bar{a}}\eta^{U\,\dagger}_{\bar{b}} &= 0\,, \qquad \text{type-I}\,, &(4.26)\\
\delta_{a\bar{b}}\,\eta^{D\,\dagger}_{\bar{a}}\eta^U_b &= 0\,, \qquad \text{type-II}\,. &(4.27)
\end{aligned}
$$

We can now prove that $\tan\beta$ is a physical parameter in the 2HDM-II (a similar analysis holds for the 2HDM-I[4]). In the preferred basis where $\eta^U_1 = \eta^D_2 = 0$, we shall denote: $\widehat{v} = e^{i\eta}(\cos\beta\,, \sin\beta\, e^{i\xi})$ and $\widehat{w} = e^{-i\eta}(-\sin\beta e^{-i\xi}\,, \cos\beta)$. Evaluating $\kappa^Q = \widehat{v}^* \cdot \eta^Q$ and $\rho^Q = \widehat{w}^* \cdot \eta^Q$ in the preferred basis, and recalling that the κ^Q are diagonal real matrices, it follows that:

$$\boldsymbol{I} e^{-i(\xi+2\eta)}\tan\beta = -\rho^{D\,\dagger}(\kappa^D)^{-1} = (\rho^U)^{-1}\kappa^U\,, \tag{4.28}$$

where $\boldsymbol{I}$ is the identity matrix in quark flavor space and $\kappa^Q = \sqrt{2}M_Q/v$ [see eqs. (1.23) and (1.24)]. These two definitions are consistent if $\kappa^D\kappa^U + \rho^{D\,\dagger}\rho^U = 0$ is satisfied. But the latter is equivalent to the type-II condition [which can be verified by inserting eq. (2.79) into eq. (4.27)].

To understand the phase factor that appears in eq. (4.28), we note that only unitary matrices of the form $U = \mathrm{diag}(e^{i\chi_1}\,, e^{i\chi_2})$ that span a U(1)×U(1) subgroup of the flavor-U(2) group preserve the the type-II conditions $\eta^U_1 = \eta^D_2 = 0$ in the preferred basis. Under transformations of this type, $\eta \to \eta + \chi_1$ and $\xi \to \xi + \chi_2 - \chi_1$. Using eq. (2.80), it follows that $\rho^Q \to e^{i(\chi_1+\chi_2)}\rho^Q$. Hence $\rho^Q e^{-i(\xi+2\eta)}$ is invariant with respect to such U(1)×U(1) transformations. We conclude that eq. (4.28) is covariant with respect to transformations that preserve the type-II condition.

The conditions specified in eq. (4.28) are quite restrictive. In particular, they determine the matrices ρ^Q:

$$\rho^D e^{-i(\xi+2\eta)} = \frac{-\sqrt{2}M_D\tan\beta}{v}\,, \qquad \rho^U e^{-i(\xi+2\eta)} = \frac{\sqrt{2}M_U\cot\beta}{v}\,. \tag{4.29}$$

Up to an overall phase, ρ^U and ρ^D are real diagonal matrices with non-negative entries. There is also some interesting information in the phase factors of eq. (4.29). Although the ρ^Q are pseudo-invariants, we have noted below eq. (2.83) that $e^{i\theta_{23}}\rho^Q$ is U(2)-invariant. This means that the phase factor $e^{-i(\theta_{23}+\xi+2\eta)}$ is a physical parameter. Moreover, we can now define $\tan\beta$ as a physical parameter of the 2HDM-II as follows:

$$\tan\beta = \frac{v}{3\sqrt{2}}\left|\mathrm{Tr}\left(\rho^D M_D^{-1}\right)\right|\,, \tag{4.30}$$

where $0 \leq \beta \leq \pi/2$. This is a manifestly basis-independent definition, so $\tan\beta$ is indeed physical.

---

[3] In this paper, we have slightly modified our definition of the Yukawa coupling. What is called η^D in ref. [34] is called $\eta^{D\,\dagger}$ here.

[4] Eq. (4.26) involves pseudo-invariant quantities. Nevertheless, setting these quantities to zero yields a U(2)-invariant condition.

In Higgs studies at future colliders, suppose one encounters phenomena that appear consistent with a 2HDM. It may not be readily apparent that there is any particular structure in the Higgs-fermion interactions. In particular, it could be that eq. (4.28) is simply false. Here we present a model-independent version of the $\tan\beta$ parameter. For simplicity, we assume that the Yukawa couplings of the Higgs bosons to the third generation fermions dominate, in which case we can ignore the effects of the first two generations.[5] In a one-generation model, one can introduce three $\tan\beta$-like parameters

$$\tan\beta_b \equiv \frac{v}{\sqrt{2}}\frac{|\rho^D|}{m_b}\,, \qquad \tan\beta_t \equiv \frac{\sqrt{2}}{v}\frac{m_t}{|\rho^U|}\,, \qquad \tan\beta_\tau \equiv \frac{v}{\sqrt{2}}\frac{|\rho^E|}{m_\tau}\,, \tag{4.31}$$

where $\tan\beta_\tau$ is analogous to $\tan\beta_d$ and depends on the third generation Higgs-lepton interaction. In a type-II model, one indeed has $\tan\beta_b = \tan\beta_t = \tan\beta_\tau = \tan\beta$. In the more general (type-III) 2HDM, there is no reason for the three parameters above to coincide. However, these three parameters are indeed U(2)-invariant quantities, and thus correspond to physical observables that can be measured in the laboratory. The interpretation of these parameters is straightforward. In the Higgs basis, up and down-type quarks interact with both Higgs doublets. But, clearly there exists some basis (*i.e.*, a rotation by an angle β_t from the Higgs basis) for which only one of the two up-type quark Yukawa couplings is non-vanishing. This defines the physical angle β_t. The interpretation of the other two angles is similar.

Since the phase of $e^{i\theta_{23}}\rho^Q$ is a physical parameter, one can generalize eq. (4.31) by defining

$$e^{i(\theta_{23}-\chi_b)}\tan\beta_b \equiv \frac{v}{\sqrt{2}}\frac{\rho^{D\,*}}{m_b}\,, \qquad e^{i(\theta_{23}-\chi_t)}\tan\beta_t \equiv \frac{\sqrt{2}}{v}\frac{m_t}{\rho^U}\,, \tag{4.32}$$

and similarly for $\tan\beta_\tau$. Thus, in addition to three $\tan\beta$-like parameters, there are three independent physical phases χ_b, χ_t and χ_τ that could in principle be deduced from experiment. Of course, in the 2HDM-II, one must have $\beta_b = \beta_t = \beta_\tau$ and $\chi_b = \chi_t = \chi_\tau$.

A similar analysis can be presented for the case of the 2HDM-I. In this case, one is led to define slightly different $\tan\beta$-like physical parameters. But, these would be related to those defined in eq. (4.31) in a simple way. A particular choice could be motivated if one has evidence that that either the type-I or type-II conditions are approximately satisfied.

We conclude this section by illustrating the utility of this approach in the case of the MSSM. This example has already been presented in ref. [34] in the case of a CP-conserving Higgs sector. We briefly explain how that analysis is generalized in the case of a CP-violating Higgs sector. The MSSM Higgs sector is a CP-conserving type-II 2HDM in the limit of exact supersymmetry. However, when supersymmetry breaking effects are taken into account, loop corrections to the Higgs potential and the Higgs-fermion interactions can lead to both CP-violating effects in the Higgs sector, and the (radiative) generation of the Higgs-fermion Yukawa couplings that are absent in the type-II limit. In particular, in the approximation that supersymmetric masses are significantly larger than m_Z, the effective Lagrangian that describes the coupling of the Higgs bosons to the third generation quarks is given (in the notation of [26]) by

---

[5] This is probably not a bad assumption, since κ^Q is proportional to the quark mass matrix M_Q.

$$-\mathcal{L}_{\rm eff} = (h_b+\delta h_b)(\overline{q}_L\Phi_1)b_R+(h_t+\delta h_t)(\overline{q}_L\widetilde{\Phi}_2)t_R+\Delta h_b\,(\overline{q}_L\Phi_2)b_R+\Delta h_t\,(\overline{q}_L\widetilde{\Phi}_1)t_R+\text{h.c.}\,, \tag{4.33}$$

where $\overline{q}_L \equiv (\overline{u}_L\,,\,\overline{d}_L)$. Note that the terms proportional to Δh_b and Δh_t, which are absent in the tree-level MSSM, are generated at one-loop due to supersymmetry-breaking effects Thus, we identify $\eta^D = ((h_b+\delta h_b)^*\,,\,\Delta h_b^*)$ and $\eta^U = (\Delta h_t\,,\,h_t+\delta h_t)$. The tree-level MSSM is CP-conserving, and $\xi = 0$ in the supersymmetric basis. At one-loop, CP-violating effects can shift ξ away from zero, and we shall denote this quantity by $\Delta\xi$.[6] Evaluating $\kappa^Q = \widehat{v}^*\cdot\eta^Q$ and $\rho^Q = \widehat{w}^*\cdot\eta^Q$ as we did above eq. (4.28),

$$\begin{aligned}
e^{i\eta}\kappa^D &= c_\beta(h_b+\delta h_b)^* + e^{-i\Delta\xi}s_\beta(\Delta h_b)^*\,,\\
e^{-i\eta}\rho^D &= -e^{i\Delta\xi}s_\beta(h_b+\delta h_b)^* + c_\beta(\Delta h_b)^*,\\
e^{i\eta}\kappa^U &= c_\beta\Delta h_t + e^{-i\Delta\xi}s_\beta(h_t+\delta h_t)\,,\\
e^{-i\eta}\rho^U &= -e^{i\Delta\xi}s_\beta\Delta h_t + c_\beta(h_t+\delta h_t)\,.
\end{aligned} \tag{4.34}$$

By definition, the κ^Q are real and non-negative, and related to the top and bottom quark masses via eqs. (1.23) and (1.24). Thus, the tree-level relations between m_b, m_t and h_b, h_t respectively are modified [105, 32, 27, 68, 72]:[7]

$$m_b = \frac{v\kappa^D}{\sqrt{2}} = \frac{vc_\beta h_b}{\sqrt{2}}\left[1+\text{Re}\left(\frac{\delta h_b}{h_b}+\frac{\Delta h_b}{h_b}e^{i\Delta\xi}\tan\beta\right)\right] \equiv \frac{vc_\beta h_b}{\sqrt{2}}\left[1+\text{Re}(\Delta_b)\right]\,, \tag{4.35}$$

$$m_t = \frac{v\kappa^U}{\sqrt{2}} = \frac{vs_\beta h_t}{\sqrt{2}}\left[1+\text{Re}\left(\frac{\delta h_t}{h_t}+\frac{\Delta h_t}{h_t}e^{i\Delta\xi}\cot\beta\right)\right] \equiv \frac{vs_\beta h_t}{\sqrt{2}}\left[1+\text{Re}(\Delta_t)\right]\,, \tag{4.36}$$

which define the complex quantities Δ_b and Δ_t.[8]Eq. (4.32) then yields:

$$\tan\beta_b \;=\; \left|\frac{-e^{-i\Delta\xi}s_\beta(h_b+\delta h_b)+c_\beta\Delta h_b}{c_\beta(h_b+\delta h_b)+e^{i\Delta\xi}s_\beta\Delta h_b}\right|\,, \qquad \chi_b = \theta_{23}+\psi_b+\eta\,, \tag{4.37}$$

$$\tan\beta_t \;=\; \left|\frac{c_\beta\Delta h_t+e^{-i\Delta\xi}s_\beta(h_t+\delta h_t)}{-\Delta h_t s_\beta e^{i\Delta\xi}+c_\beta(h_t+\delta h_t)}\right|\,, \qquad \chi_t = \theta_{23}+\psi_t+\eta\,, \tag{4.38}$$

---

[6] In practice, one would rephase the fields after computing the radiative corrections. But, since we are advocating basis-independent methods in this paper, there is no need for us to do this.

[7] If one of the Higgs fields is rephased in order to remove the phase $\Delta\xi$, then one simultaneously rephases $\Delta h_{b,t}$ such that the quantities $\Delta h_{b,t}e^{i\Delta\xi}$ are invariant with respect to the rephasing. In particular, h_b and h_t are not rephased, since these tree-level quantities are always real and positive and proportional to the tree-level values of m_b and m_t, respectively.

[8] In deriving eqs. (4.35) and (4.36), we computed $\kappa^Q = |\kappa^Q|$ by expanding up to linear order in the one-loop quantities $\Delta h_{b,t}$ and $\delta h_{b,t}$. Explicit expressions for Δ_b and Δ_t in terms of supersymmetric masses and parameters, and references to the original literature can be found in ref. [26].

where $\psi_{t,b} \equiv \arg(e^{-i\eta}\rho^{U,D})$. Expanding the numerators and denominators above and dropping terms of quadratic order in the one-loop quantities, we end up with

$$\tan\beta_b = \frac{\tan\beta}{1+\mathrm{Re}\,\Delta_b}\left[1+\frac{1}{s_\beta^2}\mathrm{Re}\left(\frac{\delta h_b}{h_b}-c_\beta^2\Delta_b\right)\right], \tag{4.39}$$

$$\cot\beta_t = \frac{\cot\beta}{1+\mathrm{Re}\,\Delta_t}\left[1+\mathrm{Re}\left(\Delta_t-\frac{1}{c_\beta s_\beta}\frac{\Delta h_t}{h_t}e^{i\Delta\xi}\right)\right]. \tag{4.40}$$

We have chosen to write $\tan\beta_b/\tan\beta$ in terms of Δ_b and $\delta h_b/h_b$, and $\cot\beta_t/\cot\beta$ in terms of Δ_t and $\Delta h_t/h_t$ in order to emphasize the large $\tan\beta$ behavior of the deviations of these quantities from one. In particular, keeping only the leading $\tan\beta$-enhanced corrections, eqs. (4.35) and (4.36) imply that[9]

$$\Delta_b \simeq e^{i\Delta\xi}\,\frac{\Delta h_b}{h_b}\,\tan\beta\,, \qquad \Delta_t \simeq \frac{\delta h_t}{h_t}\,. \tag{4.41}$$

That is, the complex quantity Δ_b is $\tan\beta$-enhanced. In typical models at large $\tan\beta$, the quantity $|\Delta_b|$ can be of order 0.1 or larger and of either sign. Thus, keeping only the one-loop corrections that are $\tan\beta$-enhanced,[10]

$$\tan\beta_b \simeq \frac{\tan\beta}{1+\mathrm{Re}\,\Delta_b}\,, \qquad \cot\beta_t \simeq \cot\beta\left[1-\tan\beta\,\mathrm{Re}\left(\frac{\Delta h_t}{h_t}e^{i\Delta\xi}\right)\right]. \tag{4.42}$$

Thus, we have expressed the basis-independent quantities $\tan\beta_b$ and $\tan\beta_t$ in terms of parameters that appear in the natural basis of the MSSM Higgs sector. Indeed, we find that $\tan\beta_b \neq \tan\beta_t$ as a consequence of supersymmetry-breaking loop-effects.

---

[9] Because the one-loop corrections δh_b, Δh_b, δh_t and Δh_t depend only on Yukawa and gauge couplings and the supersymmetric particle masses, they contain no hidden $\tan\beta$ enhancements or suppressions [25].

[10] In ref. [34] the one-loop $\tan\beta$-enhanced correction to $\cot\beta_t$ was incorrectly omitted.

Chapter 5

Conclusion

The basis-independent formalism will allow multiple Higgs bosons to be interpreted in experiments before the underlying dynamics (MSSM, etc.) are understood. Once the physical observables (ρ^Q, scalar masses, etc.) are measured, they can then be related to physical parameters such as the $\tan\beta$-like parameters to identify whether the scalars are consistent with a specific model, such as supersymmetric model. At this point, we have a complete theoretical description of the scalar, boson, and Yukawa sectors of the basis-independent 2HDM in both the CP-conserving and CP-violating cases. We have defined a mass matrix for neutral Higgs particles in terms of basis-independent scalar couplings [Y_2, Z_1, Z_3 ,Z_4, $Z_5 e^{-2i\theta_{23}}$, and $Z_6 e^{-i\theta_{23}}$.] Since Z_7 does not appear in the mass matrix, it is possible to have CP violation in the scalar sector of the theory (arising from Z_7 terms) even in the absence of mixing betweeen CP-even and CP-odd Higgs eigenstates. When the physical Higgs bosons are CP eigenstates (regardless of whether CP violation appears in the Lagrangian), there exist three ways to implement the mixing of the neutral Higgs particles, which we have called cases I, IIa, and IIb. These cases, which can be physically distinguished based on the invariant $Z_5 e^{-2i\theta_{23}}$, correspond to different orderings of the neutral Higgs fields in the mass matrix. We have also described the mixing angles in the special cases of $Z_6 = 0$ and $Z_6 = Z_7 = 0$. For $Z_6 = 0$, we find that barring degeneracy in the neutral scalars, the mass matrix breaks up into 1×1 and 2×2 blocks, as it does in the CP-conserving cases. However, the neutral scalars may still have indefinite CP quantum numbers, since CP violation may arise from the interactions involving Z_7. For $Z_6 = 0$ there are three cases that can be distinguished, labeled cases (*i*),(*ii*) and (*iii*) in this document, which can again be related to the different patterns of mass ordering. For completeness, we also describe the situations in which $Z_6 = 0$ and the neutral scalars are doubly degenerate. In special case of $Z_6 = Z_7 = 0$, we find that there are two physically distinguishable cases corresponding to different definitions of a time reversal (T) transformation.

The decoupling limit and the custodial limit have been developed in this basis-independent formalism, and the phenomenology of the oblique parameters and $\tan\beta$-like parameters have been defined in such a way as to be physically measurable without imposing symmetries or other assumptions on the scalar sector. The oblique parameters of the CP-violating 2HDM were analyzed numerically and found to be consistent with the experimental values within statistical error when plausible values of the scalar couplings were postulated (based on unitary limits). However, since our analysis shows that the 2HDM favors specific ranges for S and U, it may be possible to rule out the model in the future if experimental limits change. (Our analysis suggests that the 2HDM can accommodate equally well both positive and negative values of T, so that parameter is unlikely to falsify the model.) We also made use of the oblique parameters to constrain the mass difference between the neutral Higgs bosons and the charged Higgs boson. Other phenomenological impications of the 2HDM, such as FCNCs, remain open to further research.

Appendix A

The 2HDM scalar potential in a generic basis

Let Φ_1 and Φ_2 denote two complex hypercharge-one, $\mathrm{SU}(2)_{\mathrm{L}}$ doublets of scalar fields. The most general gauge-invariant scalar potential is given by

$$\begin{aligned}\mathcal{V} \;=\;& m_{11}^2\Phi_1^\dagger\Phi_1+m_{22}^2\Phi_2^\dagger\Phi_2-[m_{12}^2\Phi_1^\dagger\Phi_2+\mathrm{h.c.}]\\&+\tfrac{1}{2}\lambda_1(\Phi_1^\dagger\Phi_1)^2+\tfrac{1}{2}\lambda_2(\Phi_2^\dagger\Phi_2)^2+\lambda_3(\Phi_1^\dagger\Phi_1)(\Phi_2^\dagger\Phi_2)+\lambda_4(\Phi_1^\dagger\Phi_2)(\Phi_2^\dagger\Phi_1)\\&+\left\{\tfrac{1}{2}\lambda_5(\Phi_1^\dagger\Phi_2)^2+\big[\lambda_6(\Phi_1^\dagger\Phi_1)+\lambda_7(\Phi_2^\dagger\Phi_2)\big]\Phi_1^\dagger\Phi_2+\mathrm{h.c.}\right\}\,,\end{aligned}\tag{A.1}$$

where m_{11}^2, m_{22}^2, and $\lambda_1,\cdots,\lambda_4$ are real parameters. In general, m_{12}^2, λ_5, λ_6 and λ_7 are complex. The form of eq. (A.1) holds for any generic choice of Φ_1–Φ_2 basis, whereas the coefficients m_{ij}^2 and λ_i are basis-dependent quantities. Matching eq. (A.1) to the U(2)-covariant form of eq. (2.2), we identify:

$$Y_{11}=m_{11}^2\,,\qquad Y_{12}=Y_{21}^*=-m_{12}^2\,,\qquad Y_{22}=m_{22}^2\,,\tag{A.2}$$

and

$$\begin{aligned}&Z_{1111}=\lambda_1\,, && Z_{2222}=\lambda_2\,,\\&Z_{1122}=Z_{2211}=\lambda_3\,, && Z_{1221}=Z_{2112}=\lambda_4\,,\\&Z_{1212}=\lambda_5\,, && Z_{2121}=\lambda_5^*\,,\\&Z_{1112}=Z_{1211}=\lambda_6\,, && Z_{1121}=Z_{2111}=\lambda_6^*\,,\\&Z_{2212}=Z_{1222}=\lambda_7\,, && Z_{2221}=Z_{2122}=\lambda_7^*\,.\end{aligned}\tag{A.3}$$

Explicit formulae for the coefficients of the Higgs basis scalar potential in terms of the corresponding coefficients of eq. (A.1) in a generic basis can be found in ref. [34].

Appendix B

The neutral Higgs boson squared-mass matrix in a generic basis

Starting from eq. (2.2), one can obtain the neutral Higgs squared-mass matrix from the quadratic part of the scalar potential:

$$\mathcal{V}_{\rm mass} = \frac{1}{2} \begin{pmatrix} \Phi_a^0 & \Phi_{\bar{b}}^{0\,\dagger} \end{pmatrix} \mathscr{M}^2 \begin{pmatrix} \Phi_{\bar{c}}^{0\,\dagger} \\ \Phi_d^0 \end{pmatrix} . \tag{B.1}$$

Thus, $\mathscr{M}^2$ is given by the following matrix of second derivatives:

$$\mathscr{M}^2 = \left. \begin{pmatrix} \dfrac{\partial^2 \mathcal{V}}{\partial \Phi_a^0 \partial \Phi_{\bar{c}}^{0\,\dagger}} & \dfrac{\partial^2 \mathcal{V}}{\partial \Phi_a^0 \partial \Phi_d^0} \\[2ex] \dfrac{\partial^2 \mathcal{V}}{\partial \Phi_{\bar{b}}^{0\,\dagger} \partial \Phi_{\bar{c}}^{0\,\dagger}} & \dfrac{\partial^2 \mathcal{V}}{\partial \Phi_{\bar{b}}^{0\,\dagger} \partial \Phi_d^0} \end{pmatrix} \right|_{\Phi_a^0 = v_a} , \tag{B.2}$$

where $v_a \equiv v\widehat{v}_a/\sqrt{2}$ and $\widehat{v}_{\bar{a}}^*\widehat{v}_a = 1$. With $\mathcal{V}$ given by eq. (2.2), one finds:

$$\mathscr{M}^2 = \begin{pmatrix} (Y_{a\bar{c}})^* + \frac{1}{2}v^2\left[(Z_{a\bar{c}f\bar{e}} + Z_{f\bar{c}a\bar{e}})\,\widehat{v}_e\widehat{v}_{\bar{f}}^*\right]^* & \frac{1}{4}v^2(Z_{e\bar{a}f\bar{d}} + Z_{e\bar{d}f\bar{a}})\widehat{v}_{\bar{e}}^*\widehat{v}_{\bar{f}}^* \\[2ex] \frac{1}{4}v^2(Z_{b\bar{e}c\bar{f}} + Z_{c\bar{e}b\bar{f}})\widehat{v}_e\widehat{v}_f & Y_{b\bar{d}} + \frac{1}{2}v^2(Z_{e\bar{f}b\bar{d}} + Z_{e\bar{d}b\bar{f}})\widehat{v}_{\bar{e}}^*\widehat{v}_f \end{pmatrix} . \tag{B.3}$$

In deriving this result, we used the hermiticity properties of Y and Z to rewrite the upper left hand block so that the indices appear in the standard order for matrix multiplication in eq. (B.1). In addition, we employed:

$$\frac{\partial \Phi_e^0}{\partial \Phi_a^0} = \delta_{e\bar{a}} \,, \qquad \frac{\partial \Phi_{\bar{f}}^{0\,\dagger}}{\partial \Phi_{\bar{b}}^{0\,\dagger}} = \delta_{b\bar{f}} \,. \tag{B.4}$$

It is convenient to express the squared-mass matrix in terms of (pseudo)-invariants. To do this, we note that we can expand an hermitian second-ranked tensor [which satisfies $A_{a\bar{b}} = (A_{b\bar{a}})^*$] in terms of the eigenvectors of $V_{a\bar{b}} \equiv \widehat{v}_a \widehat{v}^*_{\bar{b}}$:

$$A_{a\bar{b}} = \mathrm{Tr}(VA)V_{a\bar{b}} + \mathrm{Tr}(WA)W_{a\bar{b}} + \left[(\widehat{v}^*_{\bar{c}}\widehat{w}_d A_{c\bar{d}})\widehat{v}_a \widehat{w}^*_{\bar{b}} + (\widehat{w}^*_{\bar{c}}\widehat{v}_d A_{c\bar{d}})\widehat{w}_a \widehat{v}^*_{\bar{b}}\right] , \tag{B.5}$$

where $W_{a\bar{b}} \equiv \widehat{w}_a \widehat{w}^*_{\bar{b}} = \delta_{a\bar{b}} - V_{a\bar{b}}$. Likewise, we can expand a second-ranked symmetric tensor with two unbarred (or two barred indices), *e.g.*,

$$A_{ab} = (\widehat{v}^*_{\bar{c}}\widehat{v}^*_{\bar{d}}A_{cd})\widehat{v}_a\widehat{v}_b + (\widehat{w}^*_{\bar{c}}\widehat{w}^*_{\bar{d}}A_{cd})\widehat{w}_a\widehat{w}_b + (\widehat{v}^*_{\bar{c}}\widehat{w}^*_{\bar{d}}A_{cd})(\widehat{v}_a\widehat{w}_b + \widehat{w}_a\widehat{v}_b) . \tag{B.6}$$

We can therefore rewrite the upper and lower right hand 2×2 blocks of the squared-mass matrix [eq. (B.3)] respectively as:

$$[\mathscr{M}^2]_{\bar{a}\bar{d}} = \tfrac{1}{2}v^2\left[Z_1 v^*_{\bar{a}} v^*_{\bar{d}} + Z_5 w^*_{\bar{a}} w^*_{\bar{d}} + Z_6(\widehat{v}^*_{\bar{a}}\widehat{w}^*_{\bar{d}} + \widehat{w}^*_{\bar{a}}\widehat{v}^*_{\bar{d}})\right] , \tag{B.7}$$

$$\begin{aligned}[\mathscr{M}^2]_{b\bar{d}} = {} & (Y_1 + Z_1 v^2)V_{b\bar{d}} + [Y_2 + \tfrac{1}{2}(Z_3 + Z_4)v^2]W_{b\bar{d}} \\ & + [(Y_3 + Z_6 v^2)\widehat{v}_b\widehat{w}^*_{\bar{d}} + (Y^*_3 + Z^*_6 v^2)\widehat{w}_b\widehat{v}^*_{\bar{d}}] .\end{aligned} \tag{B.8}$$

The upper and lower left hand blocks are then given by the hermitian adjoints of the lower and upper right hand blocks, respectively. Note that eq. (B.8) can be simplified further by eliminating Y_1 and Y_3 using the scalar potential minimum conditions [eq. (2.21)].

Let us apply this result to the Higgs bases, where $\widehat{v} = (1\,,\,0)$ and $\widehat{w} = (0\,,\,1)$. After imposing the scalar potential minimum conditions,

$$\mathscr{M}^2 = \frac{1}{2}v^2 \begin{pmatrix} Z_1 & Z^*_6 & Z_1 & Z_6 \\ Z_6 & Z_3 + Z_4 + 2Y_2/v^2 & Z_6 & Z_5 \\ Z_1 & Z^*_6 & Z_1 & Z_6 \\ Z^*_6 & Z^*_5 & Z^*_6 & Z_3 + Z_4 + 2Y_2/v^2 \end{pmatrix} . \tag{B.9}$$

The massless Goldstone boson eigenvector

$$G^0 = \frac{-i}{\sqrt{2}}\begin{pmatrix} 1 \\ 0 \\ -1 \\ 0 \end{pmatrix} , \tag{B.10}$$

can be determined by inspection [the normalization factor is chosen for consistency with eq. (2.22)]. Thus, we can perform a (unitary) similarity transformation on $\mathscr{M}^2$ to remove the Goldstone boson from the neutral Higgs squared-mass matrix. Explicitly, with the unitary matrix

$$V = \frac{1}{\sqrt{2}}\begin{pmatrix} 1 & 0 & 0 & -i \\ 0 & 1 & -i & 0 \\ 1 & 0 & 0 & i \\ 0 & 1 & i & 0 \end{pmatrix} , \tag{B.11}$$

it follows from eq. (B.9) that:

$$V^\dagger \mathscr{M}^2 V = \begin{pmatrix} \mathcal{M} & 0 \\ 0 & 0 \end{pmatrix}, \tag{B.12}$$

where $\mathcal{M}$ is the 3×3 neutral Higgs squared-mass matrix in the φ_1^0–φ_2^0–a^0 basis obtained in eq. (2.24).

We diagonalize $\mathcal{M}$ as described in section 2.4. The corresponding diagonalization of $\mathscr{M}^2$ is given by:

$$\mathcal{D}\mathscr{M}^2\mathcal{D}^\dagger \equiv \begin{pmatrix} R & 0 \\ 0 & 1 \end{pmatrix} V^\dagger \mathscr{M}^2 V \begin{pmatrix} R^T & 0 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} \mathcal{M}_D & 0 \\ 0 & 0 \end{pmatrix}, \tag{B.13}$$

where $\mathcal{M}_D = \text{diag}\,(m_1^2\,,\, m_2^2\,,\, m_3^2)$ and m_k is the mass of the neutral Higgs mass-eigenstate h_k. The diagonalizing matrix $\mathcal{D}$ is given by:

$$\mathcal{D} \equiv \begin{pmatrix} R & 0 \\ 0 & 1 \end{pmatrix} V^\dagger = \frac{1}{\sqrt{2}} \begin{pmatrix} d_{11} & d_{12} & d_{11}^* & d_{12}^* \\ d_{21} & d_{22} & d_{21}^* & d_{22}^* \\ d_{31} & d_{32} & d_{31}^* & d_{32}^* \\ d_{41} & d_{42} & d_{41}^* & d_{42}^* \end{pmatrix}, \tag{B.14}$$

where

$$\begin{array}{lll} d_{11} &=& c_{13}c_{12}\,, \qquad d_{21} = c_{13}s_{12}\,, \qquad d_{31} = s_{13}\,, \qquad d_{41} = i\,, \\ d_{12} &=& -s_{123}e^{-i\theta_{23}}\,, \quad d_{22} = c_{123}e^{-i\theta_{23}}\,, \quad d_{32} = ic_{13}e^{-i\theta_{23}}\,, \; d_{42} = 0\,, \end{array} \tag{B.15}$$

with the c_{ij} and s_{ij} defined in eq. (2.28) and

$$c_{123} \equiv c_{12} - is_{12}s_{13}\,, \qquad s_{123} \equiv s_{12} + ic_{12}s_{13}\,. \tag{B.16}$$

Note that $\mathcal{D}$ is a unitary matrix and $\det\mathcal{D} = 1$. Unitarity implies that:

$$\text{Re}\,(d_{k1}d_{\ell 1}^* + d_{k2}d_{\ell 2}^*) = \delta_{k\ell}\,, \tag{B.17}$$

$$\tfrac{1}{2}\sum_{k=1}^4 |d_{k1}|^2 = \tfrac{1}{2}\sum_{k=1}^4 |d_{k2}|^2 = 1\,, \qquad \sum_{k=1}^4 d_{k2}^2 = \sum_{k=1}^4 d_{k1}d_{k2} = 0\,. \tag{B.18}$$

Noting that $d_{41} = i$ and $d_{42} = 0$ [and using eq. (B.22)], these equations reduce to eqs. (2.36) and (2.37) given in section 2.4. In addition, det $\mathcal{D} = -i\det RW = 1$, where RW is given in eq. (2.33). This yields an additional constraint on the $d_{k\ell}$ [*c.f.* eq. (2.35)].

The matrix $\mathcal{D}$ converts the neutral Higgs basis fields into the neutral Higgs mass-eigenstates:

$$\begin{pmatrix} h_1 \\ h_2 \\ h_3 \\ G^0 \end{pmatrix} = \mathcal{D} \begin{pmatrix} \overline{H}_1^{0\,\dagger} \\ H_2^{0\,\dagger} \\ \overline{H}_1^0 \\ H_2^0 \end{pmatrix}, \tag{B.19}$$

where $\overline{H}_1^0 \equiv H_1^0 - v/\sqrt{2}$.

The mass-eigenstate fields do not depend on the choice of basis. Using the fact that H_1 is invariant and H_2 is pseudo-invariant with respect to flavor-U(2) transformations, eq. (B.19) implies that the d_{k1} are invariants whereas the d_{k2} are pseudo-invariants with the same transformation law as H_2 [eq. (2.20)]. One can also check this directly from eq. (B.13), using the fact that the physical Higgs masses must be basis-independent. These results then imply that θ_{12} and θ_{13} are invariant whereas $e^{i\theta_{23}}$ is a pseudo-invariant, *i.e.*, $e^{i\theta_{23}} \to (\det U)^{-1} e^{i\theta_{23}}$ under an arbitrary flavor-U(2) transformation U.

Finally, using the results of this appendix, we can eliminate the Higgs basis fields entirely and obtain the diagonalizing matrix that converts the neutral Higgs fields in the generic basis into the neutral Higgs mass-eigenstates:

$$\begin{pmatrix} h_1 \\ h_2 \\ h_3 \\ G^0 \end{pmatrix} = \mathcal{D} \begin{pmatrix} \overline{\Phi}^{0\,\dagger}_{\bar{b}} \widehat{U}^{\dagger}_{b\bar{a}} \\ \widehat{U}_{a\bar{b}} \overline{\Phi}^0_b \end{pmatrix} , \tag{B.20}$$

where $\overline{\Phi}^0_a \equiv \Phi^0_a - v\widehat{v}_a/\sqrt{2}$ and $\widehat{U}$ is the matrix that converts the generic basis fields into the Higgs basis fields [see eq. (2.10)]. Eq. (B.20) then yields:

$$h_k = \frac{1}{\sqrt{2}} \left[\overline{\Phi}^{0\,\dagger}_{\bar{a}} (d_{k1}\widehat{v}_a + d_{k2}\widehat{w}_a) + (d^*_{k1}\widehat{v}^*_{\bar{a}} + d^*_{k2}\widehat{w}^*_{\bar{a}}) \overline{\Phi}^0_a \right] , \tag{B.21}$$

where $h_4 \equiv G^0$. Note that the U(2)-invariance of the h_k imply that the d_{k1} are invariants and the d_{k2} are pseudo-invariants that transform oppositely to $\widehat{w}$ as $d_{k2} \to (\det U) d_{k2}$ in agreement with the previous results above. Indeed, it is useful to define:

$$d_{k1} \equiv q_{k1} \,, \qquad \text{and} \qquad d_{k2} \equiv q_{k2} e^{-i\theta_{23}} \,, \tag{B.22}$$

where *all* the $q_{k\ell}$ are U(2)-invariant [see eq. (2.39)]. In particular, $\widehat{w}_a e^{-i\theta_{23}}$ is a proper vector with respect to flavor-U(2) transformations. Hence,

$$h_k = \frac{1}{\sqrt{2}} \left[\overline{\Phi}^{0\,\dagger}_{\bar{a}} (q_{k1}\widehat{v}_a + q_{k2}\widehat{w}_a e^{-i\theta_{23}}) + (q^*_{k1}\widehat{v}^*_{\bar{a}} + q^*_{k2}\widehat{w}^*_{\bar{a}} e^{i\theta_{23}}) \overline{\Phi}^0_a \right] , \tag{B.23}$$

provides an invariant expression for the neutral Higgs mass-eigenstates.

Appendix C

Explicit formulae for the neutral Higgs masses and mixing angles

To obtain expressions for the neutral Higgs masses and mixing angles, we insert eq. (2.47) into eq. (2.2), and expand out the resulting expression, keeping only terms that are linear and quadratic in the fields. Using eqs. (2.17) and (2.18), one can express the resulting expression in terms of the invariants (Y_1, Y_2 and $Z_{1,2,3,4}$) and pseudo-invariants (Y_3, $Z_{5,6,7}$). The terms linear in the fields vanish if the potential minimum conditions [eq. (2.21)] are satisfied. We then eliminate Y_1 and Y_3 from the expressions of the quadratic terms. The result is:

$$\begin{aligned}\mathcal{V}_2 &= H^+H^-(Y_2+\frac{v^2 Z_3}{2})+\frac{v^2}{2}h_j h_k\Big\{Z_1\mathrm{Re}(q_{j1})\mathrm{Re}(q_{k1})+[\frac{Z_3+Z_4}{2}+Y_2/v^2]\mathrm{Re}(q_{j2}q^*_{k2})\\ &\qquad +\tfrac{1}{2}\mathrm{Re}(Z_5 q_{j2}q_{k2}\,e^{-2i\theta_{23}})+\mathrm{Re}(q_{j1})\mathrm{Re}(Z_6 q_{k2}\,e^{-i\theta_{23}})+\mathrm{Re}(q_{k1})\mathrm{Re}(Z_6 q_{j2}\,e^{-i\theta_{23}})\Big\}\\ &= m^2_{H^\pm}H^+H^-+\tfrac{1}{2}\sum_k m_k^2(h_k)^2+\tfrac{1}{2}v^2\sum_{j\neq k}C_{jk}h_j h_k\,. \end{aligned} \tag{C.1}$$

In eq. (C.1), there is an implicit sum over $j,k=1,\ldots,4$ (with $h_4\equiv G^0$), where the C_{kj} are given by[1]

$$C_{23}c_{12}-C_{13}s_{12} = s_{13}\,\mathrm{Re}(Z_6\,e^{-i\theta_{23}})-\tfrac{1}{2}c_{13}\,\mathrm{Im}(Z_5\,e^{-2i\theta_{23}})\,, \tag{C.2}$$

$$C_{23}s_{12}+C_{13}c_{12} = \tfrac{1}{2}(Z_1-A^2/v^2)\sin 2\theta_{13}-\cos 2\theta_{13}\,\mathrm{Im}(Z_6\,e^{-i\theta_{23}})\,, \tag{C.3}$$

$$\begin{aligned}C_{12} &= c_{12}s_{12}\Big[c_{12}^2(Z_1-A)-\mathrm{Re}(Z_5\,e^{-2i\theta_{23}})+2s_{12}c_{12}\mathrm{Im}(Z_6\,e^{-i\theta_{23}})\Big]\\ &\qquad (c_{12}^2-s_{12}^2)\Big[\tfrac{1}{2}s_{13}\mathrm{Im}(Z_5\,e^{-2i\theta_{23}})+c_{13}\mathrm{Re}(Z_6\,e^{-i\theta_{23}})\Big]\,.\end{aligned} \tag{C.4}$$

---

[1] For convenience we provide linear combinations of C_{23} and C_{13}, but the explicit forms can be obtained by a trivial calculation.

However, since $q_{41} = i$ and $q_{42} = 0$, it is clear that there are no terms in eq. (C.1) involving G^0. Hence, we may restrict the sum to run over $j, k = 1, 2, 3$. The charged Higgs mass obtained above confirms the result quoted in eq. (2.23). The neutral Higgs boson masses are given by:

$$m_k^2 = |q_{k2}|^2 A^2 + v^2 \left[q_{k1}^2 Z_1 + \mathrm{Re}(q_{k2})\,\mathrm{Re}(q_{k2} Z_5\, e^{-2i\theta_{23}}) + 2q_{k1}\mathrm{Re}(q_{k2} Z_6\, e^{-i\theta_{23}})\right] , \tag{C.5}$$

where A^2 is defined in eq. (2.42). It is often convenient to assume that $m_1 \leq m_2 \leq m_3$.

Note that the right-hand side of eq. (C.5) is manifestly U(2)-invariant. Moreover, by using eqs. (2.36) and (2.37), one finds that the sum of the three neutral Higgs boson squared-masses is given by

$$\mathrm{Tr}\ \mathcal{M} = \sum_k m_k^2 = 2Y_2 + (Z_1 + Z_3 + Z_4)v^2 , \tag{C.6}$$

as expected. A more explicit form for the neutral Higgs squared-masses than the one obtained in eq. (C.5) would require the solution of the cubic characteristic equation [eq. (2.25)]. Although an analytic solution can be found, it is too complicated to be of much use (a numerical evaluation is more practical).

For the neutral scalar states h_i to correspond to physical mass-eigenstates, the coefficients C_{jk} of eqs. (C.2), (C.3) and (C.4) must vanish. Since C_{jk} is symmetric under the interchange of its indices, the conditions $C_{jk} = 0$ yield three independent equations that determine the two mixing angles θ_{12} and θ_{13} and an invariant combination of θ_{23} and the phase of Z_6 (or Z_5). These three invariant angles are defined modulo π once a definite convention is established for the signs of neutral Higgs mass-eigenstate fields (as discussed at the end of section 2.4). Unique solutions for the invariant angles within this domain are obtained after a mass ordering for the three neutral Higgs bosons is specified (except at certain singular points of the 2HDM parameter space as noted in footnote 8).

To determine explicit formulae for the invariant angles, we shall initially assume that $Z_6 \equiv |Z_6| e^{i\theta_6} \neq 0$ and define the invariant angles ϕ and θ_{56}:

$$\begin{aligned} \phi &\equiv \theta_6 - \theta_{23} , \\ \theta_{56} &\equiv \theta_5 - \theta_6 , \end{aligned} \qquad \text{where} \qquad \begin{cases} \theta_6 \equiv \arg Z_6 , \\ \theta_5 \equiv \frac{1}{2} \arg Z_5 . \end{cases} \tag{C.7}$$

The factor of 1/2 in the definition of θ_5 has been inserted for convenience. As discussed in section 2.4, we can fix the conventions for the overall signs of the h_k fields by restricting the domain of θ_{12}, θ_{13} and ϕ to the region:

$$-\pi/2 \leq \theta_{12}\, , \, \theta_{13} < \pi/2 , \qquad 0 \leq \phi < \pi . \tag{C.8}$$

Setting $C_{13} = C_{23} = 0$ in eqs. (C.2) and (C.3) yields:

$$\tan\theta_{13} \quad = \quad \frac{\mathrm{Im}(Z_5\, e^{-2i\theta_{23}})}{2\,\mathrm{Re}(Z_6\, e^{-i\theta_{23}})} , \tag{C.9}$$

$$\tan 2\theta_{13} \quad = \quad \frac{2\,\mathrm{Im}(Z_6\, e^{-i\theta_{23}})}{Z_1 - A^2/v^2} . \tag{C.10}$$

Using the well known identity $\tan 2\theta_{13} = 2\tan\theta_{13}/(1-\tan^2\theta_{13})$, one can use eqs. (C.9) and (C.10) to eliminate θ_{13} and obtain an equation for ϕ.[2] The resulting equation for ϕ has more than one solution. Plugging a given solution for ϕ back into eq. (C.9) yields a corresponding solution for θ_{13}. Note that if $(\theta_{13}\,,\,\phi)$ is a solution to eqs. (C.9) and (C.10), then so is $(-\theta_{13}\,,\,\phi\pm\pi)$, in agreement with eqs. (2.50) and (2.52). By restricting to the domain of θ_{13} and ϕ specified by eq. (C.8), only one of these two solutions survives. However, multiple solutions to eqs. (C.9) and (C.10) still exist within the allowed domain, which correspond to different choices for the mass ordering of the three neutral Higgs fields. By imposing a particular mass ordering, a unique solution is selected [see eqs. (C.22) and (C.26)].

Finally, having obtained ϕ and $\tan\theta_{13}$, we use $C_{12}=0$ in eq. (C.4) to compute θ_{12}, with the result

$$\tan 2\theta_{12} = \frac{s_{13}\,\mathrm{Im}(Z_5\,e^{-2i\theta_{23}}) + 2c_{13}\,\mathrm{Re}(Z_6\,e^{-i\theta_{23}})}{c_{13}^2\,(A^2/v^2 - Z_1) + \mathrm{Re}(Z_5\,e^{-2i\theta_{23}}) - 2s_{13}c_{13}\,\mathrm{Im}(Z_6\,e^{-i\theta_{23}})}\,. \tag{C.11}$$

We can simplify the above result by using eqs. (C.9) and (C.10) to solve for $\mathrm{Im}(Z_5\,e^{-2i\theta_{23}})$ and $\mathrm{Im}(Z_6\,e^{-i\theta_{23}})$ and eliminate these factors from eq. (C.11). The end result is:

$$\tan 2\theta_{12} = \frac{2\cos 2\theta_{13}\,\mathrm{Re}(Z_6\,e^{-i\theta_{23}})}{c_{13}\left[c_{13}^2(A^2/v^2 - Z_1) + \cos 2\theta_{13}\,\mathrm{Re}(Z_5\,e^{-2i\theta_{23}})\right]}\,. \tag{C.12}$$

Note that if θ_{12} is a solution to eq. (C.12), then $\theta_{12}\pm\pi/2$ is also a solution. That is, eq. (C.12) yields two solutions for θ_{12} in the allowed domain [eq. (C.8)], which correspond to the two possible mass orderings of h_1 and h_2 as shown below eq. (C.25).

The neutral Higgs boson masses were given in eq. (C.5). With the help of eqs. (C.9), (C.10) and (C.12), one can can express these masses in terms of Z_1, Z_6 and the invariant angles:

$$m_1^2 = \left[Z_1 - \frac{s_{12}}{c_{12}c_{13}}\mathrm{Re}(Z_6\,e^{-i\theta_{23}}) + \frac{s_{13}}{c_{13}}\mathrm{Im}(Z_6\,e^{-i\theta_{23}})\right]v^2\,, \tag{C.13}$$

$$m_2^2 = \left[Z_1 + \frac{c_{12}}{s_{12}c_{13}}\mathrm{Re}(Z_6\,e^{-i\theta_{23}}) + \frac{s_{13}}{c_{13}}\mathrm{Im}(Z_6\,e^{-i\theta_{23}})\right]v^2\,, \tag{C.14}$$

$$m_3^2 = \left[Z_1 - \frac{c_{13}}{s_{13}}\mathrm{Im}(Z_6\,e^{-i\theta_{23}})\right]v^2\,. \tag{C.15}$$

For the subsequent analysis, it is useful to invert eqs. (C.13)–(C.15) and solve for Z_1, $\mathrm{Re}(Z_6\,e^{-i\theta_{23}})$ and $\mathrm{Im}(Z_6\,e^{-i\theta_{23}})$:

$$Z_1 v^2 = m_1^2 c_{12}^2 c_{13}^2 + m_2^2 s_{12}^2 c_{13}^2 + m_3^2 s_{13}^2\,, \tag{C.16}$$

$$\mathrm{Re}(Z_6\,e^{-i\theta_{23}})\,v^2 = c_{13}s_{12}c_{12}(m_2^2 - m_1^2)\,, \tag{C.17}$$

$$\mathrm{Im}(Z_6\,e^{-i\theta_{23}})\,v^2 = s_{13}c_{13}(c_{12}^2 m_1^2 + s_{12}^2 m_2^2 - m_3^2)\,. \tag{C.18}$$

---

[2] Recall that the quantity A^2 [eq. (2.42)] depends on ϕ via $\mathrm{Re}(Z_5\,e^{-2i\theta_{23}}) = |Z_5|\cos 2(\theta_{56}+\phi)$.

In addition, eqs. (C.10) and (C.12) can be used to express $\mathrm{Re}(Z_5\,e^{-i\theta_{23}})$ in terms of Z_6:

$$\mathrm{Re}(Z_5\,e^{-2i\theta_{23}}) = \frac{c_{13}}{s_{13}}\mathrm{Im}(Z_6\,e^{-i\theta_{23}}) + \frac{c_{12}^2 - s_{12}^2}{c_{13}s_{12}c_{12}}\,\mathrm{Re}(Z_6\,e^{-i\theta_{23}})\,. \tag{C.19}$$

Inserting eqs. (C.17) and (C.18) into eqs. (C.9) and (C.19) then yields expressions for $\mathrm{Im}(Z_5\,e^{-i\theta_{23}})$ and $\mathrm{Re}(Z_5\,e^{-i\theta_{23}})$ in terms of the invariant angles and the neutral Higgs masses. The above results can be used to derive an expression for

$$\begin{aligned}\mathrm{Im}(Z_5^*Z_6^2) \;&=\; 2\,\mathrm{Re}(Z_5 e^{-2i\theta_{23}})\,\mathrm{Re}(Z_6\,e^{-i\theta_{23}})\,\mathrm{Im}(Z_6\,e^{-i\theta_{23}}) \\ &\qquad - \mathrm{Im}(Z_5 e^{-2i\theta_{23}})\left\{[\mathrm{Re}(Z_6\,e^{-i\theta_{23}})]^2 - [\mathrm{Im}(Z_6\,e^{-i\theta_{23}})]^2\right\}\,.\end{aligned} \tag{C.20}$$

Using eq. (C.9) and eqs. (C.17)–(C.19), one can simplify the right hand side of eq. (C.20) to obtain:

$$\mathrm{Im}(Z_5^*Z_6^2)\,v^6 = 2s_{13}c_{13}^2s_{12}c_{12}\,(m_2^2 - m_1^2)(m_3^2 - m_1^2)(m_3^2 - m_2^2)\,. \tag{C.21}$$

Eq. (C.21) was first derived in ref. [88]; it is equivalent to a result initially obtained in ref. [98]. In particular, if any two of the neutral Higgs masses are degenerate, then $\mathrm{Im}(Z_5^*Z_6^2) = 0$, in which case one can always find a basis in which the pseudo-invariants Z_5 and Z_6 are simultaneously real. The neutral scalar squared-mass matrix [eq. (2.24)] then breaks up into a block diagonal form consisting of a 2×2 block and a 1×1 block. The diagonalization of the 2×2 block has a simple analytic form, and the neutral scalar mixing can be treated more simply by introducing one invariant mixing angle instead of the three needed in the general case. Note that $\mathrm{Im}(Z_5^*Z_6^2) = 0$ is a necessary (although not sufficient) requirement for a CP-conserving Higgs sector, as discussed in Chapter 3.1. For the remainder of this Appendix, we shall assume that the neutral Higgs boson masses are non-degenerate.

In order to facilitate the discussion of the CP-conserving limit and the decoupling limit of the 2HDM (which are treated in Chapter 3.1 and Appendix D, respectively), it is useful to derive a number of additional relations for the invariant angles. First, we employ eqs. (C.13)–(C.15) to eliminate θ_{12} and ϕ and obtain a single equation for θ_{13}:

$$s_{13}^2 = \frac{(Z_1v^2 - m_1^2)(Z_1v^2 - m_2^2) + |Z_6|^2v^4}{(m_3^2 - m_1^2)(m_3^2 - m_2^2)}\,. \tag{C.22}$$

Eq. (C.22) determines c_{13} (in the convention where $c_{13} \geq 0$). The sign of s_{13} is determined from eq. (C.15), which can be rewritten as:

$$\sin\phi = \frac{(Z_1v^2 - m_3^2)\tan\theta_{13}}{|Z_6|v^2}\,. \tag{C.23}$$

Since $\sin\phi \geq 0$ in the angular domain specified by eq. (C.8), it follows that the sign of s_{13} is equal to the sign of the quantity $Z_1v^2 - m_3^2$. In particular, if m_3^2 is the largest eigenvalue of $\widetilde{\mathcal{M}}$ [eq. (2.24)], then it must be greater than the largest diagonal element of $\widetilde{\mathcal{M}}$. That is, $Z_1v^2 - m_3^2 < 0$ if $m_3 > m_{1,2}$, in which case $s_{13} \leq 0$.

However, eq. (C.23) does not fix the sign of $\cos\phi$. To determine this sign, we can use eq. (C.9) to eliminate θ_{13} from eq. (C.23). Consequently, one obtains a single equation for ϕ:

$$\tan 2\phi = \frac{\mathrm{Im}(Z_5^* Z_6^2)}{\mathrm{Re}(Z_5^* Z_6^2) + \dfrac{|Z_6|^4 v^2}{m_3^2 - Z_1 v^2}}\,. \tag{C.24}$$

Given $\sin\phi \geq 0$ and $\tan 2\phi$ in the region $0 \leq \phi < \pi$, one can uniquely determine the value of ϕ (and hence the sign of $\cos\phi$). Thus, for a fixed ordering of the neutral Higgs masses, eqs. (C.22)–(C.24) provide a unique solution for (θ_{13}, ϕ) in the domain $-\pi/2 \leq \theta_{13} < \pi/2$ and $0 \leq \phi < \pi$.

Next, we note that eq. (C.17) can be rewritten as:

$$\sin 2\theta_{12} = \frac{2\,|Z_6|\,v^2\cos\phi}{c_{13}(m_2^2 - m_1^2)}\,. \tag{C.25}$$

As advertised below eq. (C.12), the mass ordering of m_1 and m_2 fixes the sign of $\sin 2\theta_{12}$. In particular, in the angular domain of eq. (C.8), $m_2 > m_1$ implies that $s_{12}\cos\phi \geq 0$. The sign of s_{12} is then fixed after using eq. (C.21) to infer that $\sin 2\theta_{56}\cos\phi \geq 0$ for $m_3 > m_2 > m_1$.

An alternative expression for θ_{12} can be obtained by combining eqs. (C.16) and (C.22): which yields:

$$c_{13}^2 s_{12}^2 = \frac{(Z_1 v^2 - m_1^2)(m_3^2 - Z_1 v^2) - |Z_6|^2 v^4}{(m_2^2 - m_1^2)(m_3^2 - m_2^2)}\,. \tag{C.26}$$

Note the similarity of the expressions given by eqs. (C.22) and (C.26); both these results play an important role in determining the conditions that govern the decoupling limit.

A simpler form for $\tan^2\theta_{13}$ can also be obtained by combining eqs. (C.10) and (C.23):

$$\tan^2\theta_{13} = \frac{m_3^2 - A^2}{m_3^2 - Z_1 v^2}\,. \tag{C.27}$$

Finally, one can derive an expression for $m_2^2 - m_3^2$, after eliminating $\mathrm{Im}(Z_6 e^{-i\theta_{23}})$ in favor of $\mathrm{Re}(Z_5 e^{-2i\theta_{23}})$ using eq. (C.19):

$$m_2^2 - m_3^2 = \frac{v^2}{c_{13}^2}\left[\mathrm{Re}(Z_5 e^{-2i\theta_{23}}) + \frac{c_{13}(s_{12}^2 - c_{12}^2 s_{13}^2)}{s_{12}c_{12}}\mathrm{Re}(Z_6 e^{-i\theta_{23}})\right]\,. \tag{C.28}$$

The expressions for the differences of squared-masses [eqs. (C.25) and (C.28)] take on rather simple forms in the CP-conserving limit.

In this discussion we have assumed that $Z_6 \neq 0$. The $Z_6 = 0$ case is treated separately in section 3.2.

Appendix D

The Decoupling Limit of the 2HDM

One can consider the case in which all but one Higgs have masses at some high scale Λ. The effective low energy theory is a one-Higgs-doublet model that corresponds to the Higgs sector of the Standard Model; the heavier scalar particles are therefore “decoupled” from the low energy theory [59, 65]. The light neutral (SM-like) Higgs has a mass of order the electroweak scale: $m_1 \sim v$. This decoupling limit corresponds to $Y_2 \gg v^2$ and $|Z_i| \lesssim \mathcal{O}(1)\ \forall\ i$. In the first section of this chapter, I present the basis-independent description of the 2HDM that was derived in [66]. Section D.2 describes the impact of CP-violation on the effective dimension-6 operators in the decoupling limit of the 2HDM.

D.1 The Decoupling Condition in the Basis-Independent Formalism

We shall order the neutral scalar masses according to $m_1 < m_{2,3}$ and define the invariant Higgs mixing angles accordingly. Thus, we expect one light CP-even Higgs boson, h_1, with couplings identical (up to small corrections) to those of the Standard Model (SM) Higgs boson. Using the fact that m_1^2, $|Z_i|v^2 \ll m_2^2$, m_3^2, $m_{H^\pm}^2$ in the decoupling limit, eqs. (C.14) and (C.15) yield:

$$|s_{12}| \lesssim \mathcal{O}\left(\frac{v^2}{m_2^2}\right) \ll 1\,, \qquad\qquad |s_{13}| \lesssim \mathcal{O}\left(\frac{v^2}{m_3^2}\right) \ll 1\,, \tag{D.1}$$

and eq. (C.24) imples that $\tan 2\phi + \tan 2\theta_{56} \ll 1$, where $\theta_{56} \equiv \frac{1}{2}\arg Z_5 - \arg Z_6$. This latter inequality is equivalent to:

$$\mathrm{Im}(Z_5\, e^{-2i\theta_{23}}) \lesssim \mathcal{O}\left(\frac{v^2}{m_3^2}\right) \ll 1\,. \tag{D.2}$$

Note that eq. (D.2) is also satisfied if $\theta_{23} \to \theta_{23} + \pi/2$. These two respective solutions (modulo π) correspond to the two possible mass orderings of h_2 and h_3.

One can explicitly verify the assumed mass hierarchy of the Higgs bosons in the decoupling limit. Using eqs. (C.13) and (D.1), it follows that $m_1^2 = Z_1 v^2$, with corrections $\lesssim \mathcal{O}(v^4/m_{2,3}^2)$. Eq. (C.27) yields $m_3^2 = A^2$, with corrections $\lesssim \mathcal{O}(v^2)$, and eq. (C.28) yields $m_3^2 - m_2^2 \lesssim \mathcal{O}(v^2)$. Finally, eqs. (2.23) and (2.42) imply that $m_{H^\pm}^2 - m_3^2 \lesssim \mathcal{O}(v^2)$. That is, $m_1 \ll m_2 \simeq m_3 \simeq m_{H^\pm}$.

The values of the $q_{k\ell}$ in the exact decoupling limit, where

$$s_{12} = s_{13} = \operatorname{Im}(Z_5\, e^{-2i\theta_{23}}) = 0\,, \tag{D.3}$$

are tabulated in Table D.1.

Table D.1: The U(2)-invariant quantities $q_{k\ell}$ in the exact decoupling limit.

k	q_{k1}	q_{k2}
1	1	0
2	0	1
3	0	i
4	i	0

It is a simple exercise to insert the values of the $q_{k\ell}$ in the exact decoupling limit into the Higgs couplings of sections 2.5 and 2.6. The couplings of $h_1 \equiv h$ are then given by:

$$\begin{aligned}\mathscr{L}_{\rm h} = {} & \tfrac{1}{2}(\partial_\mu h)^2 - \tfrac{1}{2}Z_1 v^2 h^2 - \tfrac{1}{2}vZ_1 h^3 - \tfrac{1}{8}vZ_1 h^4 + \left(gm_W W_\mu^+ W^{\mu -} + \frac{g}{2c_W} m_Z Z_\mu Z^\mu\right) h \\ & + \left[\frac{g^2}{4} W_\mu^+ W^{\mu -} + \frac{g^2}{8c_W^2} Z_\mu Z^\mu\right] h^2 + \left\{\left(\frac{eg}{2} A^\mu W_\mu^+ - \frac{g^2 s_W^2}{2c_W} Z^\mu W_\mu^+\right) G^- h + \text{h.c.}\right\} \\ & -\tfrac{1}{2} ig\left[W_\mu^+ G^- \overset{\leftrightarrow}{\partial^\mu} h + \text{h.c.}\right] + \frac{g}{2c_W} Z^\mu G^0 \overset{\leftrightarrow}{\partial_\mu} h + \frac{1}{v}\overline{D} M_D D h + \frac{1}{v}\overline{U} M_U U h\,. \end{aligned} \tag{D.4}$$

This is precisely the SM Higgs Lagrangian. Even in the most general CP-violating 2HDM, the interactions of the h in the decoupling limit are CP-conserving and diagonal in quark flavor space. CP-violating and flavor non-diagonal effects in the Higgs interactions are suppressed by factors of $\mathcal{O}(v^2/m_{2,3}^2)$. In contrast to the SM-like Higgs boson h, the interactions of the heavy neutral Higgs bosons (h_2 and h_3) and the charged Higgs bosons ($H^\pm$) exhibit both CP-violating and quark flavor non-diagonal couplings (proportional to the ρ^Q) in the decoupling limit. In particular, whereas eq. (D.2) implies that $\sin 2(\theta_5 - \theta_{23}) \ll 1$, the CP-violating invariant quantities $\sin(\theta_6 - \theta_{23})$ and $\sin(\theta_7 - \theta_{23})$ [*c.f.* eq. (3.2)] need not be small in the most general 2HDM.

One can understand the origin of the decoupling conditions [eqs. (D.1) and (D.2)] as follows. First, using eq. (C.5), we see that we can decouple h_2 and h_3 (and $H^\pm$) by taking $A^2 \gg v^2$ while sending $q_{12} \to 0$. Thus, in the convention in which the mass ordering of the three neutral Higgs states is $m_1 \leq m_2 \leq m_3$, it follows that the exact decoupling limit is formally achieved when $A^2 \to \infty$ and $|q_{12}|^2 = s_{12}^2 + c_{12}^2 s_{13}^2 = 0$, which implies that $s_{12} = s_{13} = 0$. Inserting these results into eq. (2.44) yields $\widetilde{R} = I$, where I is the 3×3 identity matrix. Consequently, $\widetilde{\mathcal{M}}$ [see eqs. (2.41)–(2.43)] must be diagonal up to corrections of $\mathcal{O}(v^2/A^2)$. However, because eq. (2.41) is dominated in the decoupling limit by its 22 and 33 elements (which are approximately degenerate), it follows that the 23 element must vanish exactly in leading order. Thus, in the exact decoupling limit, $\text{Im}(Z_5\, e^{-2i\theta_{23}}) = 0$. Note that this latter constraint is consistent with eq. (C.9), as $\theta_{13} = 0$ in the decoupling limit.

For further details and a more comprehensive treatment of the decoupling limit, see ref. [59].

D.2 Dimension 6 Operators in the Decoupling Limit

One can consider the case in which all but one Higgs have masses at some high scale Λ. The light neutral (SM-like) Higgs has a mass of order the electroweak scale: $m_1 \sim v$. In this limit,

$$\begin{aligned} s_{12} &\sim s_{13} \sim & \mathcal{O}\left(\frac{v^2}{\Lambda^2}\right), \\ c_{12} &\sim c_{13} \sim & 1, \end{aligned} \tag{D.5}$$

as discussed in [66]. In this scenario one can imagine that one has integrated out the higher mass fields, leaving the Standard Model as the effective low-energy theory. In a linear model of electroweak symmetry breaking, deviations from the Standard Model can be realized in two dimension 6 operators of the effective Lagrangian:

$$\mathcal{L}_{eff} = \mathcal{L}_{SM} + \frac{1}{v^2}(S\ \mathcal{O}_S + T\ \mathcal{O}_T),$$

where S and T are of order $\mathcal{O}(\frac{v^2}{\Lambda^2})$.[1] The parameter U, which corresponds to a dimension 8 operator and is of order $\mathcal{O}(\frac{v^4}{\Lambda^4})$, will be negligible here [53]. Thus, in a decoupling scenario, contributions to S and T from an extended Higgs sector are constrained by requiring that they not spoil the rough agreement with the Standard Model demonstrated in equations (4.21) – (4.22). Using a notation similar to that of Grojean et al. [54], one can define the operators as:

$$\mathcal{O}_S = \frac{\alpha}{4 s_W c_W} \mathcal{O}_{WB}, \quad \mathcal{O}_T = -2\alpha \mathcal{O}_h, \tag{D.6}$$

where

$$\mathcal{O}_{WB} = \Phi^\dagger \sigma^a \Phi W^a_{\mu\nu} B^{\mu\nu}, \quad \mathcal{O}_h = |\Phi^\dagger D_\mu \Phi|^2. \tag{D.7}$$

The covariant derivative is given by $D_\mu = \delta_\mu - \frac{ig}{2}\sigma^a W^a_\mu - ig' Y B_\mu$, and the field strengths by $W^a_{\mu\nu} = \delta_\mu W^a_\nu - \delta_\nu W^a_\mu + g\epsilon_{abc} W^b_\mu W^c_\nu$ and $B_{\mu\nu} = \delta_\mu B_\nu - \delta_\nu B_\mu$. $\mathcal{O}_{WB}$ and $\mathcal{O}_h$ are related to two well-known CP-conserving operators (O_{WB} and $O_\varphi^{(3)}$) from the work of Buchmüller and Wyler [23].

---

[1] If the new physics involves additional vectors, such as gauge bosons, two additional parameters (Y and W) associated with dimension 6 operators are added to the effective Lagrangian—see ref. [14].

Before calculating S and T in this limit, it is useful to derive relations between the scalar masses. This is done in Appendix B, with the results

$$\begin{aligned} m_3^2 &\equiv \Lambda^2, \\ m_2^2 &= \Lambda^2 + \mathrm{Re}(Z_5 e^{-2i\theta_{23}})v^2, \\ m_{H^\pm}^2 &= \Lambda^2 + \tfrac{1}{2}[\mathrm{Re}(Z_5 e^{-2i\theta_{23}}) - Z_4]v^2. \end{aligned} \tag{D.8}$$

These relations allow S to be expanded in the decoupling limit in powers of $\frac{v^2}{\Lambda^2}$.

Now one can calculate S in the decoupling limit. It is convenient to set the reference point m_ϕ equal to m_1. Thus, one can write the general expression for S as follows:

$$\begin{aligned} S = \frac{1}{\pi m_Z^2}\Big\{ & q_{31}^2\left[\mathcal{B}_{22}(m_Z^2;m_Z^2,m_3^2) + \mathcal{B}_{22}(m_Z^2;m_1^2,m_2^2) - m_Z^2\mathcal{B}_0(m_Z^2;m_Z^2,m_3^2)\right] \\ & + q_{21}^2\left[\mathcal{B}_{22}(m_Z^2;m_Z^2,m_2^2) + \mathcal{B}_{22}(m_Z^2;m_1^2,m_3^2) - m_Z^2\mathcal{B}_0(m_Z^2;m_Z^2,m_2^2)\right] \\ & - (q_{21}^2 + q_{31}^2)\left[\mathcal{B}_{22}(m_Z^2;m_Z^2,m_1^2) - m_Z^2\mathcal{B}_0(m_Z^2;m_Z^2,m_1^2)\right] \\ & + q_{11}^2\mathcal{B}_{22}(m_Z^2;m_2^2,m_3^2) - \mathcal{B}_{22}(m_Z^2;m_{H^\pm}^2,m_{H^\pm}^2)\Big\}, \end{aligned} \tag{D.9}$$

where the identity $\sum_{k=1}^3 q_{k1}^2 = 1$ has been used. In the decoupling limit, $q_{31} \sim q_{21} \sim \mathcal{O}\left(\frac{v^2}{\Lambda^2}\right)$, and $q_{11} \sim 1$; so S becomes

$$\begin{aligned} S = \frac{1}{\pi m_Z^2}\Big\{ & \mathcal{B}_{22}(m_Z^2;m_2^2,m_3^2) - \mathcal{B}_{22}(m_Z^2;m_{H^\pm}^2,m_{H^\pm}^2) + \mathcal{O}\left(\frac{v^4}{\Lambda^4}\right)\Big[\mathcal{B}_{22}(m_Z^2;m_Z^2,m_3^2) \\ & + \mathcal{B}_{22}(m_Z^2;m_1^2,m_2^2) - m_Z^2\mathcal{B}_0(m_Z^2;m_Z^2,m_3^2) + \mathcal{B}_{22}(m_Z^2;m_Z^2,m_2^2) + \mathcal{B}_{22}(m_Z^2;m_1^2,m_3^2) \\ & - m_Z^2\mathcal{B}_0(m_Z^2;m_Z^2,m_2^2) + \mathcal{B}_{22}(m_Z^2;m_Z^2,m_1^2) - m_Z^2\mathcal{B}_0(m_Z^2;m_Z^2,m_1^2)\Big]\Big\}. \end{aligned} \tag{D.10}$$

The expression above can be evaluated with equations (D.8) and the following rules, where the order parameter y is defined by $y \equiv \frac{v^2}{\Lambda^2}$, and a is order 1:

$$\begin{aligned} \mathcal{B}_{22}(m_Z^2;\Lambda^2 + av^2,\Lambda^2) &= -\frac{m_Z^2}{12}\left[\Delta - \ln\Lambda^2 - \frac{a}{2}y + \frac{1}{10}\frac{m_Z^2}{v^2}y + \mathcal{O}(y^2)\right], \\ \mathcal{B}_{22}(m_Z^2;\Lambda^2 + av^2,\Lambda^2 + av^2) &= -\frac{m_Z^2}{12}\left[\Delta - \ln\Lambda^2 - ay + \frac{1}{10}\frac{m_Z^2}{v^2}y + \mathcal{O}(y^2)\right], \\ \mathcal{B}_{22}(m_Z^2;\Lambda^2 + av^2,v^2) &= -\frac{m_Z^2}{12}\left[\Delta + \frac{5}{6} - \ln\Lambda^2 + \mathcal{O}(y)\right], \\ m_Z^2\mathcal{B}_0(m_Z^2;m_Z^2,\Lambda^2 + av^2) &= \frac{v^2}{2}y + \mathcal{O}(y^2). \end{aligned} \tag{D.11}$$

Thus one finds that $S = \frac{1}{\pi m_Z^2}\left[\frac{m_Z^2}{12}(\frac{m_3^2+m_2^2}{2v^2} - \frac{m_{H^\pm}^2}{v^2})y + m_Z^2\mathcal{O}\left(y^2 \ln y\right)\right]$. Since $y^2 \ln y \ll y$ for $y \ll 1$, the result may be written

$$S \approx \frac{1}{24\pi}\frac{m_3^2 + m_2^2 - 2m_{H^\pm}^2}{m_3^2}. \tag{D.12}$$

Hence, the limits on S given in equations (4.21) – (4.22) constrain the splitting between the neutral heavy Higgs massses and the charged Higgs mass to be small. This is equivalent to constraining differences of the scalar couplings, as one can see from using eq. (D.8) to rewrite eq. (D.12):

$$S \approx \frac{1}{24\pi}\frac{Z_4 v^2}{m_3^2}. \tag{D.13}$$

One can evaluate T in a similar manner, using the expansion

$$F(\Lambda^2 + av^2, \Lambda^2 + bv^2) \approx \Lambda^2 y^2 \frac{(a-b)^2}{6}. \tag{D.14}$$

The expression for T in eq. (4.13) can be written

$$\begin{aligned} T \;=\; & \frac{1}{16\pi s_W^2 m_W^2}\Big\{|q_{22}|^2 F(m_{H^\pm}^2, m_2^2) + |q_{32}|^2 F(m_{H^\pm}^2, m_3^2) - q_{11}^2 F(m_2^2, m_3^2) \\ & +(q_{31}^2 + q_{21}^2)\left[F(m_{H^\pm}^2, m_1^2) - F(m_W^2, m_1^2) + F(m_Z^2, m_1^2) + 4m_W^2 B_0(0; m_W^2, m_1^2)\right. \\ & \left. -4m_Z^2 B_0(0; m_Z^2, m_1^2)\right] + q_{31}^2\left[F(m_W^2, m_3^2) - F(m_1^2, m_2^2) - F(m_Z^2, m_3^2)\right. \\ & \left. -4m_W^2 B_0(0; m_W^2, m_3^2) + 4m_Z^2 B_0(0; m_Z^2, m_3^2)\right] \\ & +q_{21}^2\left[F(m_W^2, m_2^2) - F(m_1^2, m_3^2) - F(m_Z^2, m_2^2)\right. \\ & \left. -4m_W^2 B_0(0; m_W^2, m_2^2) + 4m_Z^2 B_0(0; m_Z^2, m_2^2)\right]\Big\}. \end{aligned} \tag{D.15}$$

Noting that $|q_{22}|^2 \sim |q_{32}|^2 \sim q_{11}^2 \sim 1$, T can be expanded in the decoupling limit as

$$T = \frac{1}{16\pi s_W^2 m_W^2}\left[F(m_{H^\pm}^2, m_2^2) + F(m_{H^\pm}^2, m_3^2) - F(m_2^2, m_3^2) + q_{31}^2\mathcal{O}(v^2 \ln y) + q_{21}^2\mathcal{O}(v^2 \ln y)\right]. \tag{D.16}$$

As in the previous calculation, the answer contains terms of $\mathcal{O}(y)$ and terms of $\mathcal{O}(y^2 \ln y)$. Neglecting the latter, and using eq. (D.14), one obtains the following:

$$T \approx \frac{(m_{H^\pm}^2 - m_3^2)(m_{H^\pm}^2 - m_2^2)}{48\pi s_W^2 m_W^2 m_3^2}, \tag{D.17}$$

or

$$T \approx \frac{\left[Z_4 - \mathrm{Re}(Z_5 e^{-2i\theta_{23}})\right]\left[Z_4 + \mathrm{Re}(Z_5 e^{-2i\theta_{23}})\right] v^4}{192\pi s_W^2 m_W^2 m_3^2}. \tag{D.18}$$

As a check on this calculation, one can use eqs. (D.12) and (D.17) to calculate the effect of the MSSM Higgs sector on S and T in the decoupling limit. In the supersymmetric limit CP is conserved, so m_2 becomes m_{H^0} and m_3 becomes m_{A^0}. The MSSM mass relations can be approximated in the decoupling limit by the following:

$$\begin{aligned} m^2_{H^\pm} &= m^2_{A^0} + m^2_W, \\ m^2_{H^0} &= m^2_{A^0} + m^2_Z \sin^2 2\beta + \mathcal{O}(v^2/m^2_{A^0}). \end{aligned} \tag{D.19}$$

Substituting eq. (D.19) into (D.12) and (D.17) gives

$$\begin{aligned} S(MSSM - Higgs) &\approx \frac{m_Z^2(\sin^2 2\beta - 2c_W^2)}{24\pi m^2_{A^0}}, \\ T(MSSM - Higgs) &\approx \frac{m_Z^2(c_W^2 - \sin^2 2\beta)}{48\pi s_W^2 m^2_{A^0}}, \end{aligned} \tag{D.20}$$

which agree with the results in [64].

D.3 The Lack of CP-Violating Effects in Dimension 6 Operators

One notes that despite allowing for the physical Higgs fields to be mixings of CP-eigenstates, no new phenomena related to CP-violation have emerged in this calculation. In fact, although there exist CP-violating operators of dimension 6, the following discussion will show that they do not contribute to S or T.

The CP-violating dimension 6 operators, as listed in ref. [23], are:

$$\mathcal{O}_{\Phi\tilde{G}} = (\Phi^\dagger\Phi)\tilde{G}^{A\mu\nu}G^A_{\mu\nu}, \tag{D.21}$$

$$\mathcal{O}_{\Phi\tilde{W}} = (\Phi^\dagger\Phi)\tilde{W}^{a\mu\nu}W^a_{\mu\nu}, \tag{D.22}$$

$$\mathcal{O}_{\Phi\tilde{B}} = (\Phi^\dagger\Phi)\tilde{B}^{\mu\nu}B_{\mu\nu}, \tag{D.23}$$

$$\mathcal{O}_{\tilde{W}B} = (\Phi^\dagger\sigma^a\Phi)\tilde{W}^{a\mu\nu}B_{\mu\nu}, \tag{D.24}$$

where $\tilde{V}^{\mu\nu} \equiv \frac{1}{2}\epsilon^{\mu\nu\rho\sigma}V_{\rho\sigma}$.

The operators in equations (D.21) – (D.23) cannot contribute to T [see eq. (4.2)]; there is no CP-violating equivalent of $\mathcal{O}_h$. However, one might ask whether the operator $\mathcal{O}_{\tilde{W}B}$, which is the CP-violating analogue of $\mathcal{O}_{WB}$, contributes to S. Starting with eq. (D.24), one obtains the following:

$$\begin{aligned} \mathcal{O}_{\tilde{W}B} &= \frac{1}{2}(\Phi^\dagger\sigma^a\Phi)\epsilon^{\mu\nu\rho\sigma}W^a_{\rho\sigma}B_{\mu\nu} \\ &= \frac{1}{2}(\Phi^\dagger\sigma^a\Phi)\epsilon^{\mu\nu\rho\sigma}(\delta_\rho W^a_\sigma - \delta_\sigma W^a_\rho + g\epsilon_{abc}W^b_\mu W^c_\nu)(\delta_\mu B_\nu - \delta_\nu B_\mu). \end{aligned} \tag{D.25}$$

After some manipulation of indices and replacing Φ with $\frac{1}{\sqrt{2}}\left(\begin{smallmatrix}0\\v\end{smallmatrix}\right)$, one finds that the contribution to $i\Pi^{\mu\nu}_{W^3B}(q^2)$ would be $-iv^2\epsilon^{\sigma\nu\mu\rho}q_\rho q_\sigma$. As this has no part proportional to $g^{\mu\nu}q^2$, one concludes that the operator $\mathcal{O}_{\tilde{W}B}$ makes no contribution to S [see equations

(4.1) and (4.4)]. Other CP-violating operators (of higher dimension) are suggested in [93] and [8]. However, in the decoupling scenario, operators above dimension 6 are suppressed by powers of $\frac{v^2}{\Lambda^2}$ [53], and thus will not produce measurable effects on oblique corrections.

Appendix E

Derivation of Basis-Independent Conditions for CP-invariance of Scalar Potential

In this section we derive the conditions for an explicitly CP-conserving scalar potential. Let us start by applying a CP-transformation of the scalar potential in eq. (2.2), *i.e.* $\mathcal{CP}\ \mathcal{V}\ \mathcal{CP}^{-1}$. Beginning with the quadratic ($Y_{a\bar{b}}$) term and using the definition in eq. (3.10), we note that

$$\mathcal{CP}\ \Phi^\dagger_{\bar{a}}\Phi_b\ \mathcal{CP}^{-1} = (\widehat{v}^*_{\bar{a}}\widehat{v}^*_{\bar{c}} \pm e^{2i\theta_{23}}\,\widehat{w}^*_{\bar{a}}\widehat{w}^*_{\bar{c}})(\widehat{v}_b\widehat{v}_d \pm e^{-2i\theta_{23}}\,\widehat{w}_b\widehat{w}_d)\Phi^T_c\,\Phi^*_{\bar{d}}\,. \tag{E.1}$$

Using $\Phi^T_c\Phi^*_{\bar{d}} = \Phi^\dagger_{\bar{d}}\Phi_c$, the first term in eq. (2.2) becomes

$$\begin{aligned}\mathcal{CP}\ \mathcal{V}_{quad}\ \mathcal{CP}^{-1} &= Y_{a\bar{b}}(\widehat{v}^*_{\bar{a}}\widehat{v}^*_{\bar{c}}\widehat{v}_b\widehat{v}_d + \widehat{w}^*_{\bar{a}}\widehat{w}^*_{\bar{c}}\widehat{w}_b\widehat{w}_d \pm e^{2i\theta_{23}}\,\widehat{w}^*_{\bar{a}}\widehat{w}^*_{\bar{c}}\widehat{v}_b\widehat{v}_d \\ &\qquad \pm e^{-2i\theta_{23}}\widehat{v}^*_{\bar{a}}\widehat{v}^*_{\bar{c}}\widehat{w}_b\widehat{w}_d)\Phi^\dagger_{\bar{d}}\Phi_c \\ &= (Y_1 + Y_2 \pm e^{2i\theta_{23}}Y^*_3\widehat{v}_d\widehat{w}^*_{\bar{c}} \pm e^{-2i\theta_{23}}Y_3\widehat{w}_d\widehat{v}^*_{\bar{c}}\Phi^\dagger_{\bar{d}}\Phi_c\,.\end{aligned} \tag{E.2}$$

Applying eq. (2.11), and using the orthogonality relation $\widehat{v}_a\widehat{w}^*_{\bar{a}} = 0$, this becomes

$$\begin{aligned}\mathcal{CP}\ \mathcal{V}_{quad}\ \mathcal{CP}^{-1} &= Y_1 + Y_2 \pm e^{-2i\theta_{23}}Y_3\widehat{v}_c\widehat{v}^*_{\bar{c}}\widehat{w}_d\widehat{w}^*_{\bar{d}}H^\dagger_2H_1 \pm e^{2i\theta_{23}}Y^*_3\widehat{v}_c\widehat{v}^*_{\bar{c}}\widehat{w}_d\widehat{w}^*_{\bar{d}}H^\dagger_1H_2 \\ &= Y_1 + Y_2 \pm (e^{-2i\theta_{23}}Y_3H^\dagger_2H_1 + h.c.)\,.\end{aligned} \tag{E.3}$$

Now requiring the Lagrangian to be invariant under CP, ie $\mathcal{CP}\ \mathcal{V}_{quad}\ \mathcal{CP}^{-1} = \mathcal{V}_{quad}$, yields the following condition:

$$Y_1 + Y_2 \pm (e^{-2i\theta_{23}}Y_3H^\dagger_2H_1 + h.c.) = Y_1 + Y_2 + (Y_3H^\dagger_1H_2 + h.c.)\,, \tag{E.4}$$

or

$$e^{-i\theta_{23}}Y_3 = \pm Y^*_3e^{i\theta_{23}}\,. \tag{E.5}$$

The analogous calculation for the quartic term in eq. (2.2) yields similar conditions for $Z_6e^{-i\theta_{23}}$ and $Z_7e^{-i\theta_{23}}$:

$$e^{-i\theta_{23}}Z_{6,7} = \pm Z^*_{6,7}e^{i\theta_{23}}\,. \tag{E.6}$$

Meanwhile, for the Z_5 part of the potential, the relevant term after doing a CP-transformation appears as

$$\mathcal{CP}\ \mathcal{V}\ \mathcal{CP}^{-1}\ \ni \tfrac{1}{2}\left[e^{-4i\theta_{23}}Z_5(H_2^\dagger H_1)^2 + h.c.\right]\ , \tag{E.7}$$

which leads to the condition

$$Z_5^* e^{2i\theta_{23}} = Z_5 e^{-2i\theta_{23}}\,. \tag{E.8}$$

One can now calculate, for example,

$$\begin{aligned} Z_5^* Z_6^2 &= (Z_5 e^{-4i\theta_{23}})(Z_6^{*2}e^{4i\theta_{23}}) = Z_5 Z_6^{*2}, \\ Z_6 Z_7^* &= (\pm Z_6^* e^{2i\theta_{23}})(\mp Z_7 e^{-2i\theta_{23}}) = -Z_6^* Z_7, \\ Z_5^*(Z_6^2+Z_7^2) &= (Z_5 e^{-4i\theta_{23}})(Z_6^{*2}e^{4i\theta_{23}} + Z_7^{*2}e^{4i\theta_{23}}) = Z_5(Z_6^{*2}+Z_7^{*2}), \end{aligned} \tag{E.9}$$

using eq. (E.6) and eq. (E.8). Thus, we replicate the CP conservation conditions of eq. (3.1).

E.1 Calculation of S, T and U

The one-loop corrections to the gauge boson propagators contain 3- and 4-point interactions between gauge bosons and the Higgs bosons of the 2HDM, the form of which can be read off from eqs. (2.65) – (2.66). The resulting Feynman rules in t'Hooft-Feynman gauge are shown in Table E.1. To simplify the Feynman rules, we have made use of eqs. (2.36) and (2.38). The 2HDM contributions to S are shown in Tables E.2 and E.2; Contributions to T and $S+U$ are displayed in Tables E.4, E.5 and E.6, respectively. The reference Standard Model contributions, which are subtracted out from the 2HDM contributions, are shown in Table E.7. The integrals are evaluated as in ref. [85]:

$$\int \frac{d^4k}{(2\pi)^4}\frac{k^\mu k^\nu}{(k^2-m_1^2)((k+q)^2-m_2^2)} = \frac{i}{16\pi^2}g^{\mu\nu}B_{22}(q^2;m_1^2,m_2^2), \tag{E.10}$$

$$\int \frac{d^4k}{(2\pi)^4}\frac{1}{(k^2-m_1^2)((k+q)^2-m_2^2)} = \frac{i}{16\pi^2}B_0(q^2;m_1^2,m_2^2), \tag{E.11}$$

$$\int \frac{d^4k}{(2\pi)^4}\frac{1}{(k^2-m^2)} = \frac{i}{16\pi^2}A_0(m^2). \tag{E.12}$$

The contributions to S from the diagrams in Table E.2 and Table E.7 are compiled according to eq. (4.5) and eq. (4.6), with the following result:

$$\begin{aligned}
\frac{g^2}{16\pi c_W^2}S &\equiv F_{ZZ}^{2H}(m_Z^2) - F_{\gamma\gamma}^{2H}(m_Z^2) - \frac{c_{2W}}{s_W c_W}F_{Z\gamma}^{2H}(m_Z^2) \\
&\quad -F_{ZZ}^{SM}(m_Z^2) + F_{\gamma\gamma}^{SM}(m_Z^2) + \frac{c_{2W}}{s_W c_W}F_{Z\gamma}^{SM}(m_Z^2) \\
&= \frac{g^2}{16\pi^2 m_Z^2}\left[q_{k1}^2\mathcal{B}_{22}(m_Z^2;m_Z^2,m_k^2) - \mathcal{B}_{22}(m_Z^2;m_Z^2,m_\phi^2)\right. \\
&\quad -m_Z^2 q_{k1}^2\mathcal{B}_0(m_Z^2;m_Z^2,m_k^2) + m_Z^2\mathcal{B}_0(m_Z^2;m_Z^2,m_\phi^2) \\
&\quad +q_{11}^2\mathcal{B}_{22}(m_Z^2;m_2^2,m_3^2) + q_{21}^2\mathcal{B}_{22}(m_Z^2;m_1^2,m_3^2) \\
&\quad \left.+q_{31}^2\mathcal{B}_{22}(m_Z^2;m_1^2,m_2^2) - \mathcal{B}_{22}(m_Z^2;m_{H^\pm}^2,m_{H^\pm}^2)\right]. \qquad \text{(E.13)}
\end{aligned}$$

The parameter T can be calculated in a similar manner, using the following relations provided in ref. [64]:

$$4B_{22}(0;m_1^2,m_2^2) = F(m_1^2,m_2^2) + A_0(m_1^2) + A_0(m_2^2)\,, \qquad \text{(E.14)}$$

$$B_0(0;m_1^2,m_2^2) = \frac{A_0(m_1^2) - A_0(m_2^2)}{m_1^2 - m_2^2}\,, \qquad \text{(E.15)}$$

with $F(m_1^2,m_2^2) \equiv \frac{1}{2}(m_1^2 + m_2^2) - \frac{m_1^2 m_2^2}{m_1^2 - m_2^2}\ln\left(\frac{m_1^2}{m_2^2}\right)$ as before. Adding the contributions to T from all the diagrams shown in Tables E.4, E.5 and E.7 yields

$$\begin{aligned}
\alpha T &\equiv \frac{A_{WW}^{2H}(0)}{m_W^2} - \frac{A_{ZZ}^{2H}(0)}{m_Z^2} - \left[\frac{A_{WW}^{SM}(0)}{m_W^2} - \frac{A_{ZZ}^{SM}(0)}{m_Z^2}\right] \\
&= \frac{g^2}{16\pi^2 m_W^2}\left\{|q_{k2}|^2 B_{22}(0;m_{H^\pm}^2,m_k^2) - q_{21}^2 B_{22}(0;m_1^2,m_3^2) - q_{11}^2 B_{22}(0;m_2^2,m_3^2)\right. \\
&\quad -q_{31}^2 B_{22}(0;m_1^2,m_2^2) - \frac{1}{2}A_0(m_{H^\pm}^2) + q_{k1}^2[B_{22}(0;m_W^2,m_k^2) - B_{22}(0;m_Z^2,m_k^2)] \\
&\quad -B_{22}(0;m_W^2,m_\phi^2) - q_{k1}^2[m_W^2 B_0(0;m_W^2,m_k^2) - m_Z^2 B_0(0;m_Z^2,m_k^2)] \\
&\quad \left.+B_{22}(0;m_Z^2,m_\phi^2) + m_W^2 B_0(0;m_W^2,m_\phi^2) - m_Z^2 B_0(0;m_Z^2,m_\phi^2)\right\}. \qquad \text{(E.16)}
\end{aligned}$$

This can be simplified using eq. (E.14) and $\alpha = \frac{g^2 s_W^2}{4\pi}$:

$$\begin{aligned}
T = \quad &\frac{1}{16\pi m_W^2 s_W^2}\left\{|q_{k2}|^2 F(m_{H^\pm}^2,m_k^2) - q_{21}^2 F(m_1^2,m_3^2) - q_{11}^2 F(m_2^2,m_3^2) - q_{31}^2 F(m_1^2,m_2^2)\right. \\
&-F(m_W^2,m_\phi^2) + q_{k1}^2[F(m_W^2,m_k^2) - F(m_Z^2,m_k^2)] + F(m_Z^2,m_\phi^2) \\
&-4q_{k1}^2[m_W^2 B_0(0;m_W^2,m_k^2) - m_Z^2 B_0(0;m_Z^2,m_k^2)] \\
&\left.+4m_W^2 B_0(0;m_W^2,m_\phi^2) - 4m_Z^2 B_0(0;m_Z^2,m_\phi^2)\right\}. \qquad \text{(E.17)}
\end{aligned}$$

Lastly, adding all of the contributions to $S+U$ in Tables E.6 and E.7 gives the following:

$$\begin{aligned}
\frac{g^2}{16\pi}(S+U) &= F_{WW}(m_W^2) - F_{\gamma\gamma}(m_W^2) - \frac{c_W}{s_W}F_{Z\gamma}(m_W^2) \\
&= \frac{g^2}{16\pi^2 m_W^2}\Big[-q_{k1}^2 m_W^2 \mathcal{B}_0(m_W^2; m_W^2, m_k^2) + m_W^2 \mathcal{B}_0(m_W^2; m_W^2, m_\phi^2) \\
&\quad -\mathcal{B}_{22}(m_W^2; m_W^2, m_\phi^2) + q_{k1}^2 \mathcal{B}_{22}(m_W^2; m_W^2, m_k^2) \\
&\quad +|q_{k2}|^2 \mathcal{B}_{22}(m_W^2; m_{H^\pm}^2, m_k^2) - 2\mathcal{B}_{22}(m_W^2; m_{H^\pm}^2, m_{H^\pm}^2)\Big],
\end{aligned} \tag{E.18}$$

or

$$\begin{aligned}
S+U &= \frac{1}{\pi m_W^2}\Big[-q_{k1}^2 m_W^2 \mathcal{B}_0(m_W^2; m_W^2, m_k^2) + m_W^2 \mathcal{B}_0(m_W^2; m_W^2, m_\phi^2) \\
&\quad -\mathcal{B}_{22}(m_W^2; m_W^2, m_\phi^2) + q_{k1}^2 \mathcal{B}_{22}(m_W^2; m_W^2, m_k^2) \\
&\quad +|q_{k2}|^2 \mathcal{B}_{22}(m_W^2; m_{H^\pm}^2, m_k^2) - 2\mathcal{B}_{22}(m_W^2; m_{H^\pm}^2, m_{H^\pm}^2)\Big].
\end{aligned} \tag{E.19}$$

Table E.1: Feynman rules used in the calculation of the oblique parameters.

 $= \frac{ig^2}{2}\mathrm{Re}(\lvert q_{i1}\rvert^2 + \lvert q_{i2}\rvert^2)g^{\mu\nu}$ $= \frac{ig^2}{2}g^{\mu\nu}$	 $= \frac{ig^2}{2c_W^2}\mathrm{Re}(\lvert q_{i1}\rvert^2 + \lvert q_{i2}\rvert^2)g^{\mu\nu}$ $= \frac{ig^2}{2c_W^2}g^{\mu\nu}$
 $= \frac{ig^2}{2}g^{\mu\nu}$	 $= \frac{ig^2}{2c_W^2}c_{2W}^2 g^{\mu\nu}$
 $= iq_{i1}m_W g\ g^{\mu\nu}$	 $= iq_{i1}m_Z g\ g^{\mu\nu}$
 $= -iq_{i2}\frac{g}{2}(p_2+p_1)^\mu$	 $= \mathrm{Im}(q_{32}q_{12}^*)\frac{g}{2c_W}(p_2+p_1)^\mu$ $= q_{21}\frac{g}{2c_W}(p_2+p_1)^\mu$
 $= -iq_{i1}\frac{g}{2}(p_2+p_1)^\mu$	 $= \mathrm{Im}(q_{22}q_{32}^*)\frac{g}{2c_W}(p_2+p_1)^\mu$ $= -q_{11}\frac{g}{2c_W}(p_2+p_1)^\mu$
 $= -i\frac{g}{2}(p_2+p_1)^\mu$	 $= \mathrm{Im}(q_{12}q_{22}^*)\frac{g}{2c_W}(p_2+p_1)^\mu$ $= q_{31}\frac{g}{2c_W}(p_2+p_1)^\mu$
 $= igs_W(p_2+p_1)^\mu$	 $= (c_W^2 - s_W^2)\frac{ig}{2c_W}(p_2+p_1)^\mu$
 $= im_W g\ g^{\mu\nu}$	 $= im_Z g\ g^{\mu\nu}$
	 $= -\frac{g}{2c_W}(p_2+p_1)^\mu$

Table E.2: Diagrams representing the 2HDM contributions to S, part 1.

Contributions to $\Pi_{ZZ}^{2H}(m_Z^2)$	
	$= -\frac{g^2 M_Z^2}{16\pi^2 c_W^2} q_{i1}^2 B_0(m_Z^2; m_Z^2, m_i^2)$
	$= \frac{g^2}{16\pi^2 c_W^2} q_{i1}^2 B_{22}(m_Z^2; m_Z^2, m_i^2)$
	$= \frac{g^2}{16\pi^2 c_W^2} q_{21}^2 B_{22}(m_Z^2; m_1^2, m_3^2)$
	$= \frac{g^2}{16\pi^2 c_W^2} q_{11}^2 B_{22}(m_Z^2; m_2^2, m_3^2)$
	$= \frac{g^2}{16\pi^2 c_W^2} q_{31}^2 B_{22}(m_Z^2; m_1^2, m_2^2)$
	$= \frac{g^2}{16\pi^2 c_W^2} c_{2W}^2 B_{22}(m_Z^2; m_{H^\pm}^2, m_{H^\pm}^2)$

Table E.3: Diagrams representing the 2HDM contributions to S, part 2.

Contributions to $\Pi_{\gamma\gamma}^{2H}(m_Z^2)$ and $\Pi_{Z\gamma}^{2H}(m_Z^2)$	
	$= 4\frac{g^2}{16\pi^2} s_W^2 B_{22}(m_Z^2; m_{H^\pm}^2, m_{H^\pm}^2)$
	$= \frac{2g^2}{16\pi^2 c_W} s_W c_{2W} B_{22}(m_Z^2; m_{H^\pm}^2, m_{H^\pm}^2)$

Table E.4: Diagrams representing the 2HDM contributions to T, part 1.

Contributions to $A_{WW}^{2H}(0)$
 $= -\frac{g^2 m_W^2}{16\pi^2} q_{i1}^2 B_0(0; m_W^2, m_i^2)$
 $= \frac{g^2}{16\pi^2} q_{i1}^2 B_{22}(0; m_W^2, m_i^2)$
 $= \frac{g^2}{16\pi^2} \|q_{12}\|^2 B_{22}(0; m_{H^\pm}^2, m_1^2)$
 $= \frac{g^2}{16\pi^2} \|q_{22}\|^2 B_{22}(0; m_{H^\pm}^2, m_2^2)$
 $= \frac{g^2}{16\pi^2} \|q_{32}\|^2 B_{22}(0; m_{H^\pm}^2, m_3^2)$
 $= -\frac{1}{2} \frac{g^2}{16\pi^2} A_0(m_i^2)$
 $= -\frac{1}{2} \frac{g^2}{16\pi^2} A_0(m_{H^\pm}^2)$

Table E.5: Diagrams representing the 2HDM contributions to T, part 2.

Contributions to $A^{2H}_{ZZ}(0)$



$$= -\frac{g^2 m_Z^2}{16\pi^2 c_W^2} q_{i1}^2 B_0(0; m_Z^2, m_i^2)$$



$$= \frac{g^2}{16\pi^2 c_W^2} q_{i1}^2 B_{22}(0; m_Z^2, m_i^2)$$



$$= \frac{g^2}{16\pi^2 c_W^2} q_{21}^2 B_{22}(0; m_1^2, m_3^2)$$



$$= \frac{g^2}{16\pi^2 c_W^2} q_{11}^2 B_{22}(0; m_2^2, m_3^2)$$



$$= \frac{g^2}{16\pi^2 c_W^2} q_{31}^2 B_{22}(0; m_1^2, m_2^2)$$



$$= -\frac{1}{2}\frac{g^2}{16\pi^2 c_W^2} A_0(m_i^2)$$



$$= -\frac{1}{2}\frac{g^2}{16\pi^2 c_W^2} c_{2W}^2 A_0(m_{H^\pm}^2)$$



$$= \frac{g^2}{16\pi^2 c_W^2} c_{2W}^2 B_{22}(0; m_{H^\pm}^2, m_{H^\pm}^2)$$
$$= \frac{1}{2}\frac{g^2}{16\pi^2 c_W^2} c_{2W}^2 A_0(m_{H^\pm}^2)$$

Table E.6: Diagrams representing the 2HDM contributions to $S + U$.

Contributions to $\Pi^{2H}_{WW}(m_W^2)$



$= -\frac{g^2 m_W^2}{16\pi^2} q_{i1}^2 B_0(m_W^2; m_W^2, m_i^2)$



$= \frac{g^2}{16\pi^2} q_{i1}^2 B_{22}(m_W^2; m_W^2, m_i^2)$



$= \frac{g^2}{16\pi^2} |q_{i2}|^2 B_{22}(m_W^2; m_{H^\pm}^2, m_i^2)$

Contributions to $\Pi^{2H}_{\gamma\gamma}(m_W^2)$ and $\Pi^{2H}_{Z\gamma}(m_W^2)$



$= 4\frac{g^2}{16\pi^2} s_W^2 B_{22}(m_W^2; m_{H^\pm}^2, m_{H^\pm}^2)$



$= 2\frac{g^2 s_W c_{2W}}{16\pi^2 c_W} B_{22}(m_W^2; m_{H^\pm}^2, m_{H^\pm}^2)$

Table E.7: Standard Model contributions to the oblique parameters.

Contributions to $\Pi^{SM}_{WW}(m_W^2)$ and $\Pi^{SM}_{ZZ}(m_Z^2)$	
	$= -\frac{g^2 m_W^2}{16\pi^2} B_0(m_W^2; m_W^2, m_1^2)$
	$= \frac{g^2}{16\pi^2} B_{22}(m_W^2; m_W^2, m_1^2)$
	$= -\frac{g^2 m_Z^2}{16\pi^2 c_W^2} B_0(m_Z^2; m_Z^2, m_1^2)$
	$= \frac{g^2}{16\pi^2 c_W^2} B_{22}(m_Z^2; m_Z^2, m_1^2)$
Contributions to $A^{SM}_{WW}(0)$ and $A^{SM}_{ZZ}(0)$	
	$= -\frac{g^2 m_W^2}{16\pi^2} B_0(0; m_W^2, m_1^2)$
	$= \frac{g^2}{16\pi^2} B_{22}(0; m_W^2, m_1^2)$
	$= -\frac{g^2 m_Z^2}{16\pi^2 c_W^2} B_0(0; m_Z^2, m_1^2)$
	$= \frac{g^2}{16\pi^2 c_W^2} B_{22}(0; m_Z^2, m_1^2)$

Appendix F

Derivation of Tree-Level Unitarity Limits

In section 1.1.4.2, constraints from perturbative unitarity in the Standard Model were reviewed. In particular, an upper bound for the SM Higgs boson mass m_h were derived from the scattering of $W^+W^- \to W^+W^-$. One can use a similar argument to put upper bounds on the magnitudes of the Z_i parameters in the CP-violating 2HDM. The implications of unitarity for the 2HDM has been studied in the context of scattering of gauge bosons and the physical scalars [4, 47, 74, 82, 83, 118]. By putting an upper limit ξ on the amplitude for a process $\varphi_A\varphi_B \to \varphi_C\varphi_D$, one can quantify the constraints from perturbative unitarity as follows:

$$\left|\frac{g_{ABCD}}{16\pi}\right| < \xi. \tag{F.1}$$

Here the parameter ξ will be taken to be $\frac{1}{2}$, as in section 1.1.4.2. We will only consider tree-level scattering here, so only quartic couplings will be involved, namely $W^+W^-W^+W^-$, $W^+W^-H^+H^-$, $(H^+e^{i\theta_{23}})(H^+e^{i\theta_{23}})W^-W^- + \text{h.c.}$, $Z^0Z^0Z^0h_m$, $G^0h_mG^-(H^+e^{i\theta_{23}}) + \text{h.c.}$, $Z^0Z^0H^+H^-$, and $Z^0Z^0W^-(H^+e^{i\theta_{23}})$.

The equivalence theorem (see section 1.1.2) allows one to equate a high energy scattering amplitude involving gauge bosons to the analogous amplitude involving Goldstone bosons[1], by making the replacements $W^\pm \to G^\pm$, $Z^0 \to G^0$. Thus, one can translate limits on the gauge boson/Higgs couplings into limits on the Goldstone/Higgs couplings. The resulting constraints on Z_1, Z_3, Z_3+Z_4, $\text{Re}(Z_5e^{-2i\theta_{23}})$, and $\text{Re}(Z_6e^{-i\theta_{23}})$ can be read off directly from $\mathcal{V}_4$ [eq. (2.60)], as shown in Table F.1.

The CP-violating parameters $\text{Im}(Z_5e^{-2i\theta_{23}})$ and $\text{Im}(Z_6e^{-i\theta_{23}})$ appear in a more convoluted form in the quartic scalar potential. From the interaction

$$\tfrac{1}{2}\text{Im}(q_{m2}Z_6\, e^{-i\theta_{23}})\, G^0G^0G^0h_m\,, \tag{F.2}$$

---

[1] up to a sign, which will not be important here

Table F.1: Calculation of tree-level unitarity limits on the CP-conserving quartic couplings. Combinatoric factors are included to take into account identical particles.

Relevant Term in Scalar Potential	Amplitude	Unitarity bound
$\frac{1}{2}Z_1 G^+G^-G^+G^-$	$\frac{1}{16\pi}(\frac{1}{2}Z_1)\cdot 4$	$\lvert Z_1\rvert < 4\pi$
$\frac{1}{2}Z_3 G^0G^0H^+H^-$	$\frac{1}{16\pi}(\frac{1}{2}Z_3)\cdot 2$	$\lvert Z_3\rvert < 8\pi$
$(Z_3+Z_4)G^+G^-H^+H^-$	$\frac{1}{16\pi}(Z_3+Z_4)$	$\lvert Z_3+Z_4\rvert < 8\pi$
$\frac{1}{2}Z_5 e^{-2i\theta_{23}}H^+H^+e^{2i\theta_{23}}G^-G^- + \text{h.c.}$	$\frac{1}{16\pi}\text{Re}(Z_5 e^{-2i\theta_{23}})\cdot 4$	$\lvert\text{Re}(Z_5 e^{-2i\theta_{23}})\rvert < 2\pi$
$Z_6 e^{-i\theta_{23}}G^0G^0G^-(H^+e^{i\theta_{23}}) + \text{h.c.}$	$\frac{1}{16\pi}\text{Re}(Z_6 e^{-i\theta_{23}})\cdot 4$	$\lvert\text{Re}(Z_6 e^{-i\theta_{23}})\rvert < 2\pi$

and Table 2.1, one can write Feynman rules for $m = 1, 2$:

$$\begin{aligned} g_{G^0G^0G^0h_1} &= \tfrac{1}{2}(-s_{12}\text{Im}[Z_6 e^{-i\theta_{23}}] - c_{12}s_{13}\text{Re}[Z_6 e^{-i\theta_{23}}])\cdot 3!\,, \\ g_{G^0G^0G^0h_2} &= \tfrac{1}{2}(c_{12}\text{Im}[Z_6 e^{-i\theta_{23}}] - s_{12}s_{13}\text{Re}[Z_6 e^{-i\theta_{23}}])\cdot 3!\,. \end{aligned} \tag{F.3}$$

Unitarity requires $|g_{G^0G^0G^0h_m}| < 8\pi$. It is convenient to combine the two limits in quadrature to isolate $\text{Im}(Z_6 e^{-i\theta_{23}})$:

$$\begin{aligned} |g_{G^0G^0G^0h_1}|^2 + |g_{G^0G^0G^0h_2}|^2 &< 64\pi^2\,, \\ \Big[\text{Im}[Z_6 e^{-i\theta_{23}}]\Big]^2 + s_{13}^2\Big[\text{Re}[Z_6 e^{-i\theta_{23}}]\Big]^2 &< \frac{64\pi^2}{9}\,. \end{aligned} \tag{F.4}$$

Since $s_{13}^2\left[\text{Re}(Z_6 e^{-i\theta_{23}})\right]^2$ is real and non-negative, it must be true that $|\text{Im}(Z_6 e^{-i\theta_{23}})| < \frac{8\pi}{3}$. Similarly, one can use the term

$$\tfrac{1}{2}i\,G^0 h_m\Big\{G^-H^+e^{i\theta_{23}}\left[q_{m2}^* Z_4 - q_{m2}Z_5 e^{-2i\theta_{23}}\right] + \text{h.c.}\Big\}\,, \tag{F.5}$$

with $m = 1, 2$ to derive the following:

$$\begin{aligned} g_{G^0G^-(H^+e^{i\theta_{23}})h_1} &= -c_{12}s_{13}Z_4 - s_{12}\text{Im}(Z_5 e^{-2i\theta_{23}}) - c_{12}s_{13}\text{Re}(Z_5 e^{-2i\theta_{23}}), \\ g_{G^0G^-(H^+e^{i\theta_{23}})h_2} &= -s_{12}s_{13}Z_4 + c_{12}\text{Im}(Z_5 e^{-2i\theta_{23}}) - s_{12}s_{13}\text{Re}(Z_5 e^{-2i\theta_{23}}). \end{aligned} \tag{F.6}$$

Adding in quadrature and applying the unitarity bound gives, after some simplification:

$$s_{13}^2\left[Z_4 + \text{Re}(Z_5 e^{-2i\theta_{23}})\right]^2 + \left[\text{Im}(Z_5 e^{-2i\theta_{23}})\right]^2 < 64\pi^2. \tag{F.7}$$

One can conclude that $|\text{Im}(Z_5 e^{-2i\theta_{23}})| < 8\pi$.

Bibliography

[1] Kenji Abe et al. A High-Precision Measurement of the Left-Right Z Boson Cross-Section Asymmetry. *Phys. Rev. Lett.*, 84:5945–5949, 2000.

[2] Koya Abe et al. An Improved Direct Measurement of Leptonic Coupling Asymmetries with Polarized Z Bosons. *Phys. Rev. Lett.*, 86:1162–1166, 2001.

[3] U. Aglietti et al. Tevatron for LHC report: Higgs [hep-ph/0612172]. 2006.

[4] Andrew G. Akeroyd, Abdesslam Arhrib, and El-Mokhtar Naimi. Note on Tree-level Unitarity in the General Two Higgs Doublet Model. *Phys. Lett.*, B490:119–124, 2000.

[5] Guido Altarelli and G. Isidori. Lower limit on the Higgs mass in the standard model: An Update. *Phys. Lett.*, B337:141–144, 1994.

[6] C. Amsler and others [Particle Data Group]. Review of Particle Physics. *Phys. Lett.*, B667:1, 2008.

[7] Greg W. Anderson. New Cosmological Constraints on the Higgs Boson and Top Quark Masses. *Phys. Lett.*, B243:265–270, 1990.

[8] Thomas Appelquist and Guo-Hong Wu. The Electroweak Chiral Lagrangian and New Precision Measurements. *Phys. Rev.*, D48:3235–3241, 1993.

[9] Peter Arnold and Stamatis Vokos. Instability of Hot Electroweak Theory: Bounds on m(H) and M(t). *Phys. Rev.*, D44:3620–3627, 1991.

[10] K. A. Assamagan et al. The Higgs Working Group: Summary Report 2003 [hep-ph/0406152]. 2004.

[11] David Atwood, Laura Reina, and Amarjit Soni. Phenomenology of Two Higgs Doublet Models with Flavor Changing Neutral Currents. *Phys. Rev.*, D55:3156–3176, 1997.

[12] Jonathan Bagger and Carl Schmidt. Equivalence Theorem Redux. *Phys. Rev.*, D41:264, 1990.

[13] R. Barate et al. Search for the Standard Model Higgs Boson at LEP. *Phys. Lett.*, B565:61–75, 2003.

[14] Riccardo Barbieri, Alex Pomarol, Riccardo Rattazzi, and Alessandro Strumia. Electroweak Symmetry Breaking after LEP-1 and LEP-2. *Nucl. Phys.*, B703:127–146, 2004.

[15] Vernon D. Barger, Tao Han, and J. Jiang. Tan beta Determination from Heavy Higgs Boson Production at Linear Colliders. *Phys. Rev.*, D63:075002, 2001.

[16] J. Bernabeu, G. C. Branco, and M. Gronau. CP Restrictions on Quark Mass Matrices. *Phys. Lett.*, B169:243–247, 1986.

[17] D.S. Bernstein. *Matrix Mathematics*. Princeton University Press, Princeton, NJ, 2005.

[18] Thomas Binoth and Adrian Ghinculov. New Developments in the 1/N Expansion and Nonperturbative Higgs Physics. *Nucl. Phys.*, B550:77–98, 1999.

[19] E. Boos et al. Polarisation in Sfermion Ddecays: Determining tan(beta) and Trilinear Couplings. *Eur. Phys. J.*, C30:395–407, 2003.

[20] F. J. Botella and Joao P. Silva. Jarlskog - like Invariants for Theories with Scalars and Fermions. *Phys. Rev.*, D51:3870–3875, 1995.

[21] Vincenzo Branchina and Hugo Faivre. Effective Potential (In)stability and Lower Bounds on the Scalar (Higgs) Mass. *Phys. Rev.*, D72:065017, 2005.

[22] Vincenzo Branchina, Hugo Faivre, and Vincent Pangon. Effective potential and vacuum stability. *J. Phys.*, G36:015006, 2009.

[23] W. Buchmuller and D. Wyler. Effective Lagrangian Analysis of New Interactions and Flavor Conservation. *Nucl. Phys.*, B268:621, 1986.

[24] N. Cabibbo, L. Maiani, G. Parisi, and R. Petronzio. Bounds on the Fermions and Higgs Boson Masses in Grand Unified Theories. *Nucl. Phys.*, B158:295–305, 1979.

[25] Marcela S. Carena, David Garcia, Ulrich Nierste, and Carlos E. M. Wagner. b → s Gamma and Supersymmetry with Large tan(beta). *Phys. Lett.*, B499:141–146, 2001.

[26] Marcela S. Carena and Howard E. Haber. Higgs Boson Theory and Phenomenology. ((V)). *Prog. Part. Nucl. Phys.*, 50:63–152, 2003.

[27] Marcela S. Carena, M. Olechowski, S. Pokorski, and C. E. M. Wagner. Electroweak Symmetry Breaking and Bottom - Top Yukawa Unification. *Nucl. Phys.*, B426:269–300, 1994.

[28] J. A. Casas, J. R. Espinosa, and M. Quiros. Improved Higgs Mass Stability Bound in the Standard Model and implications for supersymmetry. *Phys. Lett.*, B342:171–179, 1995.

[29] D. Chang, W. S. Hou, and W. Y. Keung. Two Loop Contributions of Flavor Changing Neutral Higgs Bosons to mu → e gamma. *Phys. Rev.*, D48:217–224, 1993.

[30] Michael S. Chanowitz and Mary K. Gaillard. The TeV Physics of Strongly Interacting W's and Z's. *Nucl. Phys.*, B261:379, 1985.

[31] S. Y. Choi and others. Determining tan(beta) in tau tau Fusion to SUSY Higgs Bosons at a Photon Collider. *Phys. Lett.*, B606:164–172, 2005.

[32] J. A. Coarasa, Ricardo A. Jimenez, and Joan Sola. Strong Effects on the Hadronic Widths of the Neutral Higgs Bosons in the MSSM. *Phys. Lett.*, B389:312–320, 1996.

[33] John M. Cornwall, David N. Levin, and George Tiktopoulos. Derivation of Gauge Invariance from High-Energy Unitarity Bounds on the s Matrix. *Phys. Rev.*, D10:1145, 1974.

[34] Sacha Davidson and Howard E. Haber. Basis-Independent Methods for the Two-Higgs-Doublet Model. *Phys. Rev.*, D72:035004, 2005.

[35] Nilendra G. Deshpande and Ernest Ma. Pattern of Symmetry Breaking with Two Higgs Doublets. *Phys. Rev.*, D18:2574, 1978.

[36] M. A. Doncheski, Stephen Godfrey, and Shou-hua Zhu. Measurement of tan(beta) in Associated t H+- Production in Gamma Gamma Collisions. *Phys. Rev.*, D68:053001, 2003.

[37] John F. Donoghue and Ling Fong Li. Properties of Charged Higgs Bosons. *Phys. Rev.*, D19:945, 1979.

[38] Loyal Durand, Peter N. Maher, and Kurt Riesselmann. Two Loop Unitarity Constraints on the Higgs Boson Coupling. *Phys. Rev.*, D48:1084–1096, 1993.

[39] Martin B. Einhorn. Speculations on a Strongly Interacting Higgs Sector. *Nucl. Phys.*, B246:75, 1984.

[40] Martin B. Einhorn and D. R. Timothy Jones. The Effective Potential, the Renormalisation Group and Vacuum Stability. *JHEP*, 04:051, 2007.

[41] J. R. Espinosa and M. Quiros. Improved metastability bounds on the standard model Higgs mass. *Phys. Lett.*, B353:257–266, 1995.

[42] Jonathan L. Feng and Takeo Moroi. Determining tan(beta) from the SUSY Higgs Sector at Future e+ e- Colliders. *Phys. Rev.*, D56:5962–5980, 1997.

[43] Zoltan Fodor, Kieran Holland, Julius Kuti, Daniel Nogradi, and Chris Schroeder. New Higgs Physics from the Lattice. *PoS*, LAT2007:056, 2007.

[44] L. Lavoura G.C. Branco and J.P. Silva. *CP Violation.* Oxford University Press, Oxford, England, 1999.

[45] Howard Georgi and Dimitri V. Nanopoulos. Suppression of Flavor Changing Effects from Neutral Spinless Meson Exchange in Gauge Theories. *Phys. Lett.*, B82:95, 1979.

[46] J. M. Gerard and M. Herquet. A Twisted Custodial Symmetry in the Two-Higgs-Doublet Model. *Phys. Rev. Lett.*, 98:251802, 2007.

[47] I. F. Ginzburg and I. P. Ivanov. Tree-level Unitarity Constraints in the 2HDM with CP- Violation [hep-ph/0312374]. 2003.

[48] S. L. Glashow. Partial Symmetries of Weak Interactions. *Nucl. Phys.*, 22:579–588, 1961.

[49] Sheldon L. Glashow and Steven Weinberg. Natural Conservation Laws for Neutral Currents. *Phys. Rev.*, D15:1958, 1977.

[50] Sheldon L. Glashow and Steven Weinberg. Natural Conservation Laws for Neutral Currents. *Phys. Rev.*, D15:1958, 1977.

[51] W. Grimus, L. Lavoura, O. M. Ogreid, and P. Osland. A Precision Constraint on Multi-Higgs-Doublet Models. *J. Phys.*, G35:075001, 2008.

[52] W. Grimus, L. Lavoura, O. M. Ogreid, and P. Osland. The oblique parameters in multi-Higgs-doublet models. *Nucl. Phys.*, B801:81–96, 2008.

[53] Benjamin Grinstein and Mark B. Wise. Operator Analysis for Precision Electroweak Physics. *Phys. Lett.*, B265:326–334, 1991.

[54] Christophe Grojean, Witold Skiba, and John Terning. Disguising the Oblique Parameters. *Phys. Rev.*, D73:075008, 2006.

[55] Bohdan Grzadkowski, John F. Gunion, and Jan Kalinowski. Finding the CP-Violating Higgs Bosons at e+ e- Colliders. *Phys. Rev.*, D60:075011, 1999.

[56] J. F. Gunion, B. Grzadkowski, H. E. Haber, and J. Kalinowski. LEP Limits on CP-Violating Non-Minimal Higgs Sectors. *Phys. Rev. Lett.*, 79:982–985, 1997.

[57] J. F. Gunion and Howard E. Haber. Higgs Bosons in Supersymmetric Models. 1. *Nucl. Phys.*, B272:1, 1986.

[58] J. F. Gunion and Howard E. Haber. Higgs Bosons in Supersymmetric Models. 2. Implications for Phenomenology. *Nucl. Phys.*, B278:449, 1986.

[59] John F. Gunion and Howard E. Haber. The CP-Conserving Two-Higgs-Doublet Model: The Approach to the Decoupling Limit. *Phys. Rev.*, D67:075019, 2003.

[60] John F. Gunion and Howard E. Haber. Conditions for CP-Violation in the General Two-Higgs- Doublet Model. *Phys. Rev.*, D72:095002, 2005.

[61] John F. Gunion, Howard E. Haber, Gordon L. Kane, and Sally Dawson. The Higgs Hunter's Guide. SCIPP-89/13.

[62] John F. Gunion, T. Han, J. Jiang, and A. Sopczak. Determining tan(beta) with Neutral and Charged Higgs Bosons at a Future e+ e- Linear Collider. *Phys. Lett.*, B565:42–60, 2003.

[63] H. E. Haber, Gordon L. Kane, and T. Sterling. The Fermion Mass Scale and Possible Effects of Higgs Bosons on Experimental Observables. *Nucl. Phys.*, B161:493, 1979.

[64] Howard E. Haber. Introductory Low-energy Supersymmetry. *Boulder TASI*, 92:0589-688, 1993.

[65] Howard E. Haber and Yosef Nir. Multiscalar Models with a High-Energy Scale. *Nucl. Phys.*, B335:363, 1990.

[66] Howard E. Haber and Deva O'Neil. Basis-Independent Methods for the Two-Higgs-Doublet Model II: The Significance of tan(beta). *Phys. Rev.*, D74:015018, 2006.

[67] Howard E. Haber and Alex Pomarol. Constraints from Global Symmetries on Radiative Corrections to the Higgs Sector. *Phys. Lett.*, B302:435–441, 1993.

[68] Lawrence J. Hall, Riccardo Rattazzi, and Uri Sarid. The Top Quark Mmass in Supersymmetric SO(10) Unification. *Phys. Rev.*, D50:7048–7065, 1994.

[69] Lawrence J. Hall and Mark B. Wise. Flavor Changing Higgs-Boson Couplings. *Nucl. Phys.*, B187:397, 1981.

[70] Thomas Hambye and Kurt Riesselmann. Matching conditions and Higgs mass upper bounds revisited. *Phys. Rev.*, D55:7255–7262, 1997.

[71] Thomas Hambye and Kurt Riesselmann. SM Higgs Mass Bounds from Theory [hep-ph/9708416]. 1997.

[72] Ralf Hempfling. Yukawa Coupling Unification with Supersymmetric Threshold Corrections. *Phys. Rev.*, D49:6168–6172, 1994.

[73] R.A. Horn and C.R. Johnson. *Matrix Analysis*. Cambridge Univ. Press, Cambridge, UK, 1990.

[74] H. Huffel and G. Pocsik. Unitarity Bounds on Higgs Boson Masses in the Weinberg-Salam Model with Two Higgs Doublets. *Zeit. Phys.*, C8:13, 1981.

[75] P. Q. Hung and G. Isidori. Anatomy of the Higgs Mass Spectrum. *Phys. Lett.*, B402:122–129, 1997.

[76] Gino Isidori, Giovanni Ridolfi, and Alessandro Strumia. On the Metastability of the Standard Model Vacuum. *Nucl. Phys.*, B609:387–409, 2001.

[77] Gino Isidori, Vyacheslav S. Rychkov, Alessandro Strumia, and Nikolaos Tetradis. Gravitational corrections to Standard Model vacuum decay. *Phys. Rev.*, D77:025034, 2008.

[78] S. Jager and S. Willenbrock. Unitarity and the Scale of Fermion Mass Generation. *Phys. Lett.*, B435:139–144, 1998.

[79] C. Jarlskog. A Basis Independent Formulation of the Connection Between Quark Mass Matrices, CP Violation and Experiment. *Z. Phys.*, C29:491–497, 1985.

[80] C. Jarlskog. Commutator of the Quark Mass Matrices in the Standard Electroweak Model and a Measure of Maximal CP Violation. *Phys. Rev. Lett.*, 55:1039, 1985.

[81] Jun-ichi Kamoshita. Impact of Light Higgs Properties on the Determination of tan(beta) and m(susy). *Prog. Theor. Phys.*, 100:773–780, 1998.

[82] Shinya Kanemura, Takahiro Kubota, and Eiichi Takasugi. Lee-Quigg-Thacker Bounds for Higgs Boson Masses in a Two Doublet Model. *Phys. Lett.*, B313:155–160, 1993.

[83] Shinya Kanemura, Yasuhiro Okada, Eibun Senaha, and C. P. Yuan. Higgs Coupling Constants as a Probe of New Physics. *Phys. Rev.*, D70:115002, 2004.

[84] R. Kinnunen, S. Lehti, F. Moortgat, A. Nikitenko, and M. Spira. Measurement of the H/A to tau tau Cross Section and Possible Constraints on tan beta. *Eur. Phys. J.*, C40N5:23–32, 2005.

[85] (ed.) Langacker, P. Precision Tests of the Standard Electroweak Model. Singapore, Singapore: World Scientific (1995) 1008 p. (Advanced series on directions in high energy physics: 14).

[86] W. Langguth and I. Montvay. A Numerical Estimate of the Upper Limit for the Higgs Boson Mass. *Z. Phys.*, C36:725, 1987.

[87] L. Lavoura. Signatures of Discrete Symmetries in the Scalar Sector. *Phys. Rev.*, D50:7089–7092, 1994.

[88] L. Lavoura and Joao P. Silva. Fundamental CP Violating Quantities in a SU(2) x U(1) Model with Many Higgs Doublets. *Phys. Rev.*, D50:4619–4624, 1994.

[89] Benjamin W. Lee, C. Quigg, and H. B. Thacker. The Strength of Weak Interactions at Very High-Energies and the Higgs Boson Mass. *Phys. Rev. Lett.*, 38:883–885, 1977.

[90] Benjamin W. Lee, C. Quigg, and H. B. Thacker. Weak Interactions at Very High-Energies: The Role of the Higgs Boson Mass. *Phys. Rev.*, D16:1519, 1977.

[91] M. Lindner. Implications of Triviality for the Standard Model. *Zeit. Phys.*, C31:295, 1986.

[92] Manfred Lindner, Marc Sher, and Helmut W. Zaglauer. Probing Vacuum Stability Bounds at the Fermilab Collider. *Phys. Lett.*, B228:139, 1989.

[93] Anthony C. Longhitano. Low-Energy Impact of a Heavy Higgs Boson Sector. *Nucl. Phys.*, B188:118, 1981.

[94] M. Luscher and P. Weisz. Is There a Strong Interaction Sector in the Standard Lattice Higgs Model? *Phys. Lett.*, B212:472, 1988.

[95] M. Luscher and P. Weisz. Scaling Laws and Triviality Bounds in the Lattice phi**4 Theory. 2. One Component Model in the Phase with Spontaneous Symmetry Breaking. *Nucl. Phys.*, B295:65, 1988.

[96] William J. Marciano, G. Valencia, and S. Willenbrock. Renormalization Group Improved Unitarity Bounds on the Higgs Boson and Top Quark Masses. *Phys. Rev.*, D40:1725, 1989.

[97] William J. Marciano and Scott S. D. Willenbrock. Radiative Corrections to Heavy Higgs Scalar Production and Decay. *Phys. Rev.*, D37:2509, 1988.

[98] A. Mendez and A. Pomarol. Signals of CP Violation in the Higgs Sector. *Phys. Lett.*, B272:313–318, 1991.

[99] F.D. Murnaghan. *The Unitary and Rotation Groups.* Spartan Books, Washington, DC, 1962.

[100] E. A. Paschos. Diagonal Neutral Currents. *Phys. Rev.*, D15:1966, 1977.

[101] G. Passarino and M. J. G. Veltman. One Loop Corrections for e+ e- Annihilation Into mu+ mu- in the Weinberg Model. *Nucl. Phys.*, B160:151, 1979.

[102] Michael Edward Peskin and Daniel V. Schroeder. *An Introduction to Quantum Field Theory.* Addison-Wesley, Reading, Massachusetts, 1995.

[103] Michael Edward Peskin and Tatsu Takeuchi. A New Constraint on a Strongly Interacting Higgs Sector. *Phys. Rev. Lett.*, 65:964–967, 1990.

[104] Michael Edward Peskin and Tatsu Takeuchi. Estimation of Oblique Electroweak Corrections. *Phys. Rev.*, D46:381–409, 1992.

[105] Damien M. Pierce, Jonathan A. Bagger, Konstantin T. Matchev, and Ren-jie Zhang. Precision Corrections in the Minimal Supersymmetric Standard Model. *Nucl. Phys.*, B491:3–67, 1997.

[106] Apostolos Pilaftsis and Carlos E. M. Wagner. Higgs Bosons in the Minimal Supersymmetric Standard Model with Explicit CP Violation. *Nucl. Phys.*, B553:3–42, 1999.

[107] Yu. F. Pirogov and O. V. Zenin. Two-Loop Renormalization Group Restrictions on the Standard Model and the Fourth Chiral Family. *Eur. Phys. J.*, C10:629–638, 1999.

[108] Alex Pomarol and Roberto Vega. Constraints on CP violation in the Higgs Sector from the Rho Parameter. *Nucl. Phys.*, B413:3–15, 1994.

[109] Jianming Qian. Higgs Searches at the Fermilab Tevatron $p\bar{p}$ Collider. *Fermilab-conf-08-575-e.*

[110] D. Rainwater. Searching for the Higgs Boson [hep-ph/0702124]. 2007.

[111] K. Riesselmann and S. Willenbrock. Ruling Out a Strongly-Interacting Standard Higgs Model. *Phys. Rev.*, D55:311–321, 1997.

[112] Marc Sher. Electroweak Higgs Potentials and Vacuum Stability. *Phys. Rept.*, 179:273–418, 1989.

[113] Marc Sher. Precise Vacuum Stability Bound in the Standard Model. *Phys. Lett.*, B317:159–163, 1993.

[114] Marc Sher and Helmut W. Zaglauer. Cosmic Ray Induced Vacuum Decay in the Standard Model. *Phys. Lett.*, B206:527, 1988.

[115] Andre Sopczak. Status of Higgs Boson Searches at the Tevatron. *Fermilab-conf-09-088-e-ppd-t*, 2009.

[116] J.L. Synge and A. Schild. *Tensor Calculus.* Dover publications, Inc., New York, NY, 1978.

[117] The ALEPH, DELPHI, L3, OPAL, SLD Collaborations, the LEP Electroweak Working Group, the SLD Electroweak and Heavy Flavour Groups. Precision Electroweak Measurements on the Z Resonance. *Phys. Rept.*, 427:257, 2006.

[118] H. Arthur Weldon. The Effects of Multiple Higgs Bosons on Tree Unitarity. *Phys. Rev.*, D30:1547, 1984.